Can a wristband wearable device accurately measure acute stress of novice surgeons during a high-fidelity surgical simulation?

Submission date	Recruitment status	Prospectively registered
20/02/2018	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
24/02/2018	Completed	Results
Last Edited	Condition category	[] Individual participant data
23/02/2018	Other	Record updated in last year

Plain English summary of protocol

Background and study aims

It is known that acute stress has a direct impact on surgical performance and patient safety as well. As it is rather unfeasible to measure surgeon's stress in the operating room, surgical simulation has been proposed as the best alternative, since it is establish way of training that is designed to replicate real-life situations replicating stress, prevent biases and provide objective metrics. As there is no gold standard method to measure stress, several subjective and/or objective methods have been proposed. The rationale for the objective methods lies in the fact that acute stress provokes changes. Therefore changes that are looked at are: Heart rhythm changes as measured by heart rate (HR) or heart rate variability (HRV), electrodermal activity (EDA) levels, thermal activity, blood pressure variability (BVP), and saliva stress biomarkers have been suggested as markers. Using a wearable device that simultaneously captures HR/HRV, BVP, and EDA, in order to evaluate the accuracy of these parameters on stress detection, as well as, to compare them against the most widely method used, the Holter derived HRV could be helpful for measuring stress levels. The aim of this study is to assess the feasibility of a new watch-sized device to noninvasively measure stress parameters in novices during a simulation task in a high fidelity simulator as well as to explore the best stress detector among the recorded parameters. This study also aims to detect if HR and HRV derived parameters when calculated from this wearable device can substitute reciprocal Holter measurements in our simulation environment.

Who can participate?

Male medical school trainees aged 23 to 26 years old

What does the study involve?

Participants are introduced to the simulator, briefed of the tasks to follow and given detailed information about the data recording equipment and the video-evaluation of their performance. HTye are given the Empatica E4 wristband (E4WB) in their non-dominant hand to wear. Additionally, all participants wear an Holter ECG rhythm monitoring and electrodes positioned in certain positions. A baseline recording phase of 10 minutes is initiated with the subjects

engaged in leisurely reading (BL phase). Immediately after the simulation exercise starts, the participants are trained on a basic skills module for 9 minutes and are videotaped for later visual analysis of any major errors (failures) detection followed by a 6 minutes recovery period.

What are the possible benefits and risks of participating? There are no direct benefits or risks with participating.

Where is the study run from? Athens University Medical school

When is the study starting and how long is it expected to run for? March 2015 to March 2018

Who is funding the study? National and Kapodistrian University of Athens (Greece)

Who is the main contact? Mr Konstantinos Georgiou (Scientific)

Contact information

Type(s)

Scientific

Contact name

Mr Konstantinos Georgiou

ORCID ID

https://orcid.org/0000-0003-3615-2500

Contact details

Medical Physics Laboratory Simulation Center Athens Medical School National and Kapodistrian University of Athens 75, Mikras Asias street Athens Greece 11527

Additional identifiers

Protocol serial number 7276

Study information

Scientific Title

Feasibility of a new wearable device to estimate acute stress in novices during high-fidelity surgical simulation

Study objectives

The first objective of this study is to assess the feasibility of a new watch-sized device to noninvasively measure stress parameters in novices during a simulation task in a high fidelity simulator as well as to explore the best stress detector among the recorded parameters.
The second objective is to detect if HR and HRV derived parameters when calculated from this wearable device can substitute reciprocal Holter measurements in our simulation environment.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Athens Medical School Ethics Committee, 19/04/201, ref:1516023954

Study design

Observational cohort study

Primary study design

Observational

Study type(s)

Other

Health condition(s) or problem(s) studied

Healthy participants

Interventions

Prior to the simulation task, all participants complete a questionnaire regarding age, height, weight, and prior simulation experience. An orientation phase is then implemented (approximately 15 minutes) during which the participants are introduced to the simulator, briefed of the tasks to follow and given detailed information about the data recording equipment and the video-evaluation of their performance. Thereafter they wear the wristband (E4WB) in their non dominant hand. Additionally, all participants wear an ambulatory Holter ECG rhythm monitoring and electrodes were positioned in predetermined thorax positions. Throughout the whole experiment, the subjects wear both the E4WB and the Holter rhythm monitor and an ambient temperature was kept in the simulation room in order to avoid any sweating artifacts.

A baseline recording phase of 10 minutes is initiated with the subjects engaged in leisurely reading (BL phase). Immediately after the simulation exercise started (T phase) where the subjects were trained on a basic skills module (Lap mentor, 3D Systems) for 9 minutes and were videotaped for later visual analysis of any major errors (failures) detection (F) followed by a 6 minutes recovery period (R phase). Each phase was tagged by triggering concomitantly the markers on both devices.

Heart Rate (HR), inter-beat interval duration (IBI), electrodermal activity (EDA), 3-axis hand motion activity (Acc), and skin temperature (ST) data were recorded from the E4WB. Furthermore, HR and heart rate variability (HRV) data were obtained from the Holter device.

Intervention Type

Primary outcome(s)

Stress is detected using the EDA at baseline and after the stimulation.

Key secondary outcome(s))

- 1. Heart Rate (HR) is recorded using the E4WB at baseline and after the stimulation
- 2. Inter-beat interval duration (IBI) is recorded using the E4WB at baseline and after the stimulation
- 3. Electrodermal activity (EDA) is recorded using the E4WB at baseline and after the stimulation
- 4. 3-axis hand motion activity (Acc) is recorded using the E4WB at baseline and after the stimulation
- 5. Skin temperature (ST) data is recorded from the E4WB at baseline and after the stimulation
- 6. HR and heart rate variability (HRV) data is recorded from the Holter device at baseline and after the stimulation

Completion date

01/03/2018

Eligibility

Key inclusion criteria

- 1. Male novice trainees aged 23 to 26 years old
- 2. Body mass index (BMI) from 18.5 to 24.9
- 3. Medical students or PHY1
- 4. No simulation experience prior

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Male

Key exclusion criteria

if experienced surgeons in simulation

Date of first enrolment

12/12/2016

Date of final enrolment

20/12/2017

Locations

Countries of recruitment

Greece

Study participating centre Athens University Medical school

MPLSC11527 Athens Greece 11527

Sponsor information

Organisation

National and Kapodistrian University of Athens, Greece

ROR

https://ror.org/04gnjpq42

Funder(s)

Funder type

University/education

Funder Name

National and Kapodistrian University of Athens, Greece

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Konstantinos Georgiou MD at kongeeorgiou@med.uoa.gr

IPD sharing plan summary

Available on request

Study outputs

Output type Details Date created Date added Peer reviewed? Patient-facing?

Participant information sheet

Participant information sheet 11/11/2025 11/11/2025 No