

# Exploring how oxygen therapy and smart limb exercises can help treat diabetes-related leg artery disease

<b>Submission date</b>	<b>Recruitment status</b>	<input type="checkbox"/> Prospectively registered
28/11/2024	No longer recruiting	<input type="checkbox"/> Protocol
<b>Registration date</b>	<b>Overall study status</b>	<input type="checkbox"/> Statistical analysis plan
02/12/2024	Completed	<input checked="" type="checkbox"/> Results
<b>Last Edited</b>	<b>Condition category</b>	<input type="checkbox"/> Individual participant data
19/01/2026	Nutritional, Metabolic, Endocrine	

## Plain English summary of protocol

### Background and study aims

Diabetes mellitus (DM) is a long-term condition caused by abnormal insulin levels, leading to high blood sugar. Lower extremity arterial disease (LEAD) is a condition where arteries in the legs are blocked, causing reduced limb function or disability. People with both DM and LEAD face high risks of amputation and death, which severely impacts their quality of life and finances. Current treatments are often ineffective and expensive. This study aims to explore a new treatment combining oxygen therapy (ASOT) and limb exercises to improve blood flow and reduce symptoms, potentially lowering amputation and death rates.

### Who can participate?

Adults aged 18 to 80 years who have been diagnosed with both DM and LEAD, are in stages I to IV of LEAD, are conscious, able to cooperate with treatment, and have had the disease for less than a year.

### What does the study involve?

Participants will first attend a screening visit to confirm eligibility. If eligible, they will undergo assessments to measure the impact of their condition. Participants will then be divided into two groups: a control group receiving standard treatment plus exercise therapy, and a study group receiving the same treatment plus ASOT. Both groups will follow their treatment regimen once daily, 5 days a week, for 10 days. The study will compare pain levels, daily living abilities, blood glucose, cholesterol, inflammation markers, and blood flow before and after treatment.

### What are the possible benefits and risks of participating?

The combined therapy of ASOT and limb exercises may reduce pain, improve self-care abilities, slow disease progression, relieve symptoms, and enhance quality of life. The therapy has few risks but should be used cautiously.

### Where is the study run from?

The Second Hospital of Hebei Medical University (China)

When is the study starting and how long is it expected to run for?

November 2021 to May 2024

Who is funding the study?

The Second Hospital of Hebei Medical University (China)

Who is the main contact?

Hongling Li, [lihongling\\_ys@163.com](mailto:lihongling_ys@163.com)

## Contact information

### Type(s)

Public, Scientific, Principal investigator

### Contact name

Dr Hongling Li

### ORCID ID

<https://orcid.org/0000-0002-5336-8701>

### Contact details

The Second Department of Rehabilitation, The Second Hospital of Hebei Medical University.

No. 215 Heping West Road, Xinhua District

Shijiazhuang City

China

050000

+86 311-66002789

[lihongling\\_ys@163.com](mailto:lihongling_ys@163.com)

## Additional identifiers

### Clinical Trials Information System (CTIS)

Nil known

### ClinicalTrials.gov (NCT)

Nil known

### Protocol serial number

Nil known

## Study information

### Scientific Title

A study to explore the effect of atmospheric saturated oxygen combined with intelligent upper and lower limb rehabilitation training on diabetic lower extremity arterial disease

### Study objectives

Atmospheric saturated oxygen therapy (ASOT) combined with upper and lower limb rehabilitation exercise therapy can improve the self-care ability of patients with diabetic lower limb arterial disease, delay disease progression, relieve symptoms, and improve quality of life.

### **Ethics approval required**

Ethics approval required

### **Ethics approval(s)**

approved 27/05/2024, Ethics committee of the Second Hospital of Hebei Medical University (No. 215 Heping West Road, Xinhua District, Shijiazhuang City, 050000, China; +86 311-66002789; pub@hb2h.com), ref: 2024-R255

### **Study design**

Single-center interventional double-blind randomized controlled trial

### **Primary study design**

Interventional

### **Study type(s)**

Quality of life

### **Health condition(s) or problem(s) studied**

Lower limb rehabilitation training for patients with diabetes and lower extremity arterial disease (LEAD)

### **Interventions**

Both groups receive standard treatment: blood glucose control (liraglutide injection, Novo Nordisk (China)), lipidlowering therapy (atorvastatin, Qilu Pharmaceutical), depressor (verapamil hydrochloride sustainedrelease capsules, Shanghai Tengry), antiplatelet therapy (aspirin, Hunan Zhongnan Pharmaceutical), anticoagulation therapy (warfarin, Qilu Pharmaceutical), circulation booster (urokinase, Guangdong Techpool BioPharma), antiinfection therapy (piperacillin /tazobactam, North China Pharmaceutical), neurotrophic support, and physical therapy. Patients with gangrene are given debridement and dressing changes to control local infection. Nursing care and education are provided in a routine fashion.

The control group receive exercise therapy in addition to the standard treatment. This involves the use of a German Motomed upper and lower limb rehabilitation trainer. Patients are seated or lying on the equipment with their backs firmly against the backrest, stabilized with a belt. For upper limb training, patients have their feet fixed on the pedals and remained stationary while their hands are fixed on the handles to perform circular movements for 30 min per session. For lower limb training, patients have their hands fixed on the handles and remained stationary while their feet are fixed on the pedals to perform circular movements for 30 min per session. When patients have difficulty performing active exercise, the responsible nurse will provide professional assistance. During the exercise, vital signs are monitored to ensure patient safety. This regimen is performed once daily, 5 days a week, for 10 days per course.

The study group receive ASOT in addition to the abovementioned regimen administered to the control group. Patients lie flat on the hospital bed with one end of a multifunctional saturated oxygen breathing apparatus connected to the ward's oxygen supply and the other connected to an oxygen tube and mask. The mask is placed correctly over the patient's nose and mouth and

secured with an appropriately tightened strap. Pure oxygen is inhaled using normobaric oxygen supply, with the partial pressure of oxygen set at 760 mmHg. Each session lasts 60 min, once daily, 5 days a week, for 10 days per course.

## **Intervention Type**

Behavioural

## **Primary outcome(s)**

1. Pain is measured using a 10 cm sliding ruler (VAS score) at baseline and after treatment
2. Barthel Index is measured using the Barthel Index rating scale at baseline and after treatment
3. Blood glucose levels are measured using a fingerstick test in the fasting state in the early morning and 2 hours after breakfast at baseline and after treatment
4. Levels of TC, TG, HDL-C, LDL-C, IL-6, CRP, and HCY are measured using a fully automated biochemical analyzer from fasting venous blood samples at baseline and after treatment
5. ABI index is measured using a Doppler stethoscope at baseline and after treatment
6. Partial pressure of oxygen within tissue microcirculation (TcPO<sub>2</sub>) is measured using a TcPO<sub>2</sub> test at baseline and after treatment
7. DPA flow velocity on the affected side is measured using color Doppler ultrasound at baseline and after treatment

## **Key secondary outcome(s)**

The general characteristics of the patients were measured using a questionnaire at baseline

## **Completion date**

01/05/2024

# **Eligibility**

## **Key inclusion criteria**

1. Meeting the diagnostic criteria for DM and LEAD
2. Confirmed to have Fontaine stage I to IV LEAD (stage I: asymptomatic; stage IIa: mild intermittent claudication; stage IIb: moderate to severe intermittent claudication; stage III: ischemic rest pain; stage IV: ischemic ulcer or gangrene)
3. Age between 18 and 80 years
4. Clear consciousness and ability to cooperate with treatment
5. Disease duration within one year
6. Provision of signed informed consent

## **Participant type(s)**

Patient

## **Healthy volunteers allowed**

No

## **Age group**

Mixed

## **Lower age limit**

18 years

**Upper age limit**

80 years

**Sex**

All

**Total final enrolment**

60

**Key exclusion criteria**

1. Severe cardiac, pulmonary, hepatic, or renal insufficiency
2. Recent history of fractures and chronic orthopedic diseases
3. Severe carbon dioxide retention
4. Known malignant diseases and tumor history
5. Coexisting peripheral vascular disease
6. Lack of cooperation or no provision of signed informed consent from the patient or his/her family members

**Date of first enrolment**

01/01/2022

**Date of final enrolment**

31/12/2023

## Locations

**Countries of recruitment**

China

**Study participating centre**

**The Second Hospital of Hebei Medical University**

No. 215 Heping West Road, Xinhua District

Shijiazhuang City

China

050000

## Sponsor information

**Organisation**

The Second Hospital of Hebei Medical University

## Funder(s)

**Funder type**

Other

**Funder Name**

Investigator initiated and funded

## Results and Publications

**Individual participant data (IPD) sharing plan**

The datasets generated and/or analysed during the current study will be included in the subsequent results publication

**IPD sharing plan summary**

Published as a supplement to the results publication

**Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#"><u>Results article</u></a>		14/01/2026	19/01/2026	Yes	No