

Posterior pericardiotomy for prevention of atrial fibrillation and pericardial effusion after aortic valve replacement

Submission date 25/12/2013	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 08/01/2014	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 12/07/2017	Condition category Surgery	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

In patients who underwent a heart surgery called coronary artery bypass graft (CABG), posterior pericardiotomy (a surgical procedure carried out on the pericardium of the heart) has been shown to be protective against postoperative atrial fibrillation (abnormal heart rate after heart surgery). This study aims to find out whether it could help patients who underwent valve surgery.

Who can participate?

Patients admitted for aortic valve replacement surgery can participate in this study.

What does the study involve?

Patients are divided randomly into two groups. One group receives posterior pericardiotomy during the valve replacement surgery and the other group, called the control group, receives the standard surgical technique.

What are the possible benefits and risks of participating?

This technique may reduce the occurrence of postoperative atrial fibrillation and accumulation of fluid around the heart in patients who underwent aortic valve replacement surgery.

Where is the study run from?

This study is run from Ochapowski Regional Hospital, Russia.

When is the study starting and how long is it expected to run for?

The study started in October 2013 and is expected to run for 1.5 years.

Who is funding the study?

The study is funded by the Kuban State Medical University, Russia.

Who is the main contact?

Dr Vasily Kaleda
vasily.kaleda@gmail.com

Contact information

Type(s)

Scientific

Contact name

Dr Vasily Kaleda

Contact details

140, Rossiyskaya str.

Krasnodar

Russian Federation

350086

-

vasily.kaleda@gmail.com

Additional identifiers

Protocol serial number

N/A

Study information

Scientific Title

The efficacy of posterior pericardiotomy in prevention of atrial fibrillation and pericardial effusion after aortic valve replacement

Study objectives

The hypothesis is that posterior pericardiotomy can reduce the incidence of atrial fibrillation and pericardial effusion after aortic valve replacement.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Kuban State Medical University Ethics Comimittee, 01/10/2013, Protocol #22

Study design

Single-centre non-blinded randomised controlled trial

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Cardiac surgery postoperative atrial fibrillation and pericardial effusion

Interventions

Patients will be randomised to two groups: intervention and control

1. Intervention group: will undergo posterior pericardiotomy in addition to conventional aortic valve replacement
2. Control group: will undergo only conventional aortic valve replacement

Intervention Type

Procedure/Surgery

Phase

Not Applicable

Primary outcome(s)

1. Mortality
2. The incidence of stroke
3. The incidence of cardiac tamponade

Concerned to in-hospital period.

Key secondary outcome(s)

1. The incidence of atrial fibrillation, measured with 48-h ECG monitoring and then with daily ECG
2. The incidence of pericardial effusion measured by echo twice after removing chest tubes and before discharge
3. The incidence of left pleural effusion measured by echo twice after removing chest tubes and before discharge
4. Length of stay in hospital

Completion date

17/04/2015

Eligibility

Key inclusion criteria

1. Age 18-69 years
2. Informed consent obtained
3. Isolated primary aortic valve replacement

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

69 years

Sex

All

Key exclusion criteria

1. Known history of atrial fibrillation before surgery
2. Amiodarone intake before surgery
3. Known history of thyrotoxicosis
4. Pericardial effusion with a maximum pericardial separation of more than 0.5 cm by transthoracic echocardiography (TTE) before surgery
5. Severe chronic obstructive pulmonary disease (COPD)
6. Left ventricular ejection fraction (LVEF) <30%
7. Left atrium (LA) diameter >50 mm
8. Pericardial adhesions
9. Any active inflammatory disease at the time of surgery (including active infection endocarditis)

Date of first enrolment

17/10/2013

Date of final enrolment

17/04/2015

Locations

Countries of recruitment

Russian Federation

Study participating centre

140, Rossiyskaya str.

Krasnodar

Russian Federation

350086

Sponsor information

Organisation

Kuban State Medical University (Russia)

ROR

<https://ror.org/04wa91k02>

Funder(s)

Funder type

University/education

Funder Name

Kuban State Medical University (Russia)

Results and Publications

Individual participant data (IPD) sharing plan

The current data sharing plans for the current study are unknown and will be made available at a later date.

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	14/02/2017		Yes	No