

# Standard (D1) versus extended (D2) lymph node dissection during gastrectomy for gastric cancer in Italy: a randomised comparison of morbidity, mortality and survival

<b>Submission date</b> 30/03/2009	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 07/04/2009	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 23/04/2021	<b>Condition category</b> Cancer	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

Not provided at time of registration

## Contact information

### Type(s)

Scientific

### Contact name

Prof Maurizio Degiuli

### Contact details

Department of Surgery  
SCDU Chirurgia Generale 1  
Ospedale S Giovanni Battista - Le molinette  
corso Bramante 88/90  
Turin  
Italy  
10126  
+39 (0)11 633 6121  
mdegiuli@hotmail.com

## Additional identifiers

EudraCT/CTIS number

IRAS number

**ClinicalTrials.gov number**

**Secondary identifying numbers**

N/A

## **Study information**

### **Scientific Title**

Italian Gastric Cancer Study Group (IGCSG) randomised clinical trial of morbidity, mortality and survival after D1 and D2 gastrectomy for gastric cancer in western patients

### **Acronym**

IGCSG-R01

### **Study objectives**

Nowadays D2 gastrectomy is the gold standard in most countries from the Far East. In western countries extended procedures have never gained much popularity for two main reasons. First, the evidence of survival benefit after D2 in Japan had been based solely on observational and retrospective studies. Second, although low morbidity and mortality rates have been reported also in western centres in retrospective studies, the fear of increasing post-operative complications and death rates has been confirmed by two large scale randomised controlled trials (RCTs) recently performed in Europe, which strongly concluded that D2 dissection was not recommended for western patients and made the British National Health service guidance discourage the use of D2 technique in routine clinical practice.

However, more recently the evidence of survival benefit after extended dissection in the Far East has finally come also from a RCT. Moreover, different critical appraisals of the European RCTs have concluded that the increase of morbidity and mortality observed in these trials were mostly related to spleno-pancreatectomy routinely performed in D2 total gastrectomies and to the little experience/volume of hospitals and surgeons involved, that this significant increase may have offset the long-term survival benefit of the procedure and that D2 dissection may offer cure to subset of patients with N2 disease; therefore, D2 gastrectomy may be of benefit if morbidity and mortality can be avoided.

Based on these previous reports, in 1994 the Italian Gastric Cancer Study Group (IGCSG) designed a multicentre phase II trial to test the feasibility and the safeness of the extended procedure performed in western patients with pancreas preserving splenectomy and strict quality control applied over all the centres. We found post-operative morbidity and mortality rates of 20.9% and 3.1%, comparable with those reported after the western standard D1-gastrectomy. Once documented the feasibility and safeness of D2 procedure in Italy, we have set up this new multicentre RCT to compare outcome and survival effects of D1 versus D2 (IGCSG-R01). This is the fourth RCT performed all over the world comparing these two procedures.

### **Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

When our RCT was designed in 1998, there was no need for ethical approval according to regional and national regulations.

## **Study design**

Interventional multicentre two-armed randomised controlled surgical trial

## **Primary study design**

Interventional

## **Secondary study design**

Randomised controlled trial

## **Study setting(s)**

Hospital

## **Study type(s)**

Treatment

## **Participant information sheet**

## **Health condition(s) or problem(s) studied**

Gastric cancer

## **Interventions**

The study was performed according to the rules of the JRGCC as regards the extent of stomach removal and the technique of lymph-node dissection (D1 and D2 procedures) and to the Japanese Classification of Gastric Carcinoma - second English Edition by the Japanese Gastric Cancer Association, particularly as concerning the definitions of classifications and grouping of regional lymph nodes, the extent of lymph node metastasis (N) and the curative potential of gastric resection (Resection A, B or C). In this new classification the regional lymph nodes are classified into three groups (compartments or levels or tiers 1 - 3), depending upon the location of the primary tumour. The operative details of the two procedures respected the general rules for gastric cancer study, as described by the Japanese Research Society for Gastric Cancer in 1981. D1 resection entailed removal of the nodes usually defined as perigastric nodes (n1 nodes) 'en bloc' with the specimen, according to the JGCA. In D2 surgery lymph nodes of n1 tier were removed together with n2 tier nodes. In the D2 arm, during total gastrectomy, the pancreas was removed only when it was suspected to be involved by the tumour. When required (clinical T greater than 1 on the greater curvature of the proximal and middle thirds of the stomach), splenectomy was performed with the pancreas preservation technique as described by Maruyama. We have considered two different types of protocol deviation: "contamination" for D1 and "non-compliance" for D2. "Contamination" was the deviation with the pathological proof of more than 2 LN stations that were not supposed to be removed while "non-compliance" was the deviation represented by the absence of LN from more than two LN stations that were required.

Arm 1, D1 gastrectomy: the time for this procedure ranged from 3 hours to 4 hours depending on the extent of gastrectomy (3 hours for distal gastrectomy and 4 hours for total gastrectomy)

Arm 2, D2 gastrectomy: the time for this procedure ranged from 4 hours to 5 hours depending on the extent of gastrectomy (4 hours for distal gastrectomy and 5 hours for total gastrectomy)

Analysis of main outcome (survival) will be performed when at least 95% of recruited patients have accrued at least 5 years of follow up.

## **Intervention Type**

Other

**Phase**

Not Applicable

**Primary outcome measure**

Survival: follow-up for mortality analysed with the Kaplan-Mayer risk assessment.

**Secondary outcome measures**

1. Complications (general and surgical) and reoperations assessed post-operatively
2. Hospital mortality, defined as death occurring within 30 days from the procedure or during patient's hospital stay

Follow-up was done at regular intervals (4 months) and follow-up data were sent every 6 months to the Reference Centre where all data were collected.

**Overall study start date**

01/06/1998

**Completion date**

13/11/2006

## **Eligibility**

**Key inclusion criteria**

1. Histology proven and potentially curable gastric adenocarcinoma, not requiring emergency surgery
2. Aged younger than 80 years, either sex
3. In adequate physical condition without serious co-morbid cardio-respiratory diseases which could preclude a safe D2 procedure
4. Have not undergone previous gastric surgery
5. Does not harbor a coexisting cancer

**Participant type(s)**

Patient

**Age group**

Adult

**Sex**

Both

**Target number of participants**

320

**Key exclusion criteria**

1. Absence of macroscopic involvement of liver and peritoneum (HO, PO)
2. Absence of macroscopic involvement of adjacent organs (T less than 4)
3. Absence of macroscopic massive involvement of N2 nodes (enlarged nodes at coeliac area)
4. Absence of malignant cells in para-aortic nodes (16B1) at biopsy and frozen section

5. Absence of malignant cells in peritoneal washing fluid, during intra-operative fresh examination
6. Absence of macroscopic residual tumour (RO)
7. No involvement of the oesophagus, cardia or duodenum

**Date of first enrolment**

01/06/1998

**Date of final enrolment**

13/11/2006

## Locations

**Countries of recruitment**

Italy

**Study participating centre**

Department of Surgery

Turin

Italy

10126

## Sponsor information

**Organisation**

Health Research Aimed at the Regione Piemonte (Ricerca Sanitaria Finalizzata della Regione Piemonte) (Italy)

**Sponsor details**

Corso Stati Uniti 1

Turin

Italy

10128

-

sanita.pubblica@regione.piemonte.it

**Sponsor type**

Government

**Website**

<http://www.ricerca-sanitaria-finalizzata.it/>

## Funder(s)

**Funder type**

Government

**Funder Name**

Ministry of Health (Italy) - Health Research Program 2003 (ref: DEG/001 - delibera Prot. n. 1837 /27.001)

## Results and Publications

**Publication and dissemination plan**

Not provided at time of registration

**Intention to publish date****Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration

**Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Other publications</a>	interim analysis	01/04/2004		Yes	No
<a href="#">Results article</a>	results	01/05/2010		Yes	No
<a href="#">Results article</a>	results	01/01/2014		Yes	No
<a href="#">Results article</a>	fifteen-year follow up results	01/06/2021	23/04/2021	Yes	No