

Reducing antibiotic use in primary care: a pharmacist-led educational and audit & feedback strategy

Submission date 25/11/2024	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 11/12/2024	Overall study status Ongoing	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 21/07/2025	Condition category Other	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

This study aims to improve how general practitioners (GPs) prescribe antibiotics by providing them with educational support and feedback. The goal is to ensure antibiotics are used correctly to combat resistance and improve patient care.

Who can participate?

All general practitioners (GPs) working in primary health care centers in three Spanish Health Care Districts: Mallorca (Balearic Islands), Paterna (Valencia), and Tarragona-Reus (Catalonia) will participate. A total of 112 GPs are required for the study.

What does the study involve?

GPs will receive personalized feedback on their antibiotic prescribing habits, including information on selection, dosage, duration, and the conditions treated. Those who request additional help or are in the highest prescribing quartile will get one-on-one support from a study pharmacist. This includes setting goals, creating action plans, and discussing specific cases. Participants will also have access to an online educational course.

What are the possible benefits and risks of participating?

The main benefit is improved antibiotic prescribing practices, which can lead to better patient outcomes and reduced antibiotic resistance. There are no significant risks for the GPs participating in this study.

Where is the study run from?

The study is conducted in three Spanish Health Care Districts: Mallorca (Balearic Islands), Paterna (Valencia), and Tarragona-Reus (Catalonia).

When is the study starting and how long is it expected to run for?

January 2023 to December 2026

Who is funding the study?

The study is funded by the Carlos III Institute of Health (PI22/01742) (Spain)

Who is the main contact?

1. Dr Laura Gallardo, laura.gallardo@ibsalut.es

2. Aina Soler, aina.soler@ibsalut.es

Contact information

Type(s)

Public, Principal investigator

Contact name

Dr Laura Gallardo

ORCID ID

<https://orcid.org/0000-0002-5769-706X>

Contact details

c/Escola Graduada, 3

Palma

Spain

07002

+34 971 17 58 97

laura.gallardo@ibsalut.es

Type(s)

Scientific

Contact name

Ms Aina Soler

Contact details

c/Escola Graduada, 3

Palma

Spain

07002

+34 971 17 58 97

aina.soler@ibsalut.es

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

Nil known

Study information

Scientific Title

A pharmacist-led educational intervention and audit & feedback approach to reduce antibiotic prescriptions in primary care

Acronym

AFA study

Study objectives

Current study hypothesis as of 06/01/2025:

An Audit & Feedback intervention targeting general practitioners (GP), which includes individualized graphical data on antibiotic prescriptions (covering antibiotic selection, dosages, treatment duration, and associated conditions) combined with a pharmacist-led educational intervention consisting of an individualized, face-to-face session with a primary care pharmacist for GPs with significant potential for improvement in their prescribing practices (above the 75th percentile), along with an online educational component, reduces antibiotic prescriptions by at least 1 point in the ratio of total antibiotic prescriptions per 100 visits, for individuals aged 14 years and older.

Previous study hypothesis:

An Audit & Feedback intervention targeting general practitioners (GP), which includes individualized graphical data on antibiotic prescriptions (covering antibiotic selection, dosages, treatment duration, and associated conditions) combined with a pharmacist-led educational intervention consisting of an individualized, face-to-face session with a primary care pharmacist for GPs with significant potential for improvement in their prescribing practices (above the 75th percentile), along with an online educational component, reduces antibiotic prescriptions by at least 4 daily doses per 1,000 inhabitants per day (DHD) in individuals over 14 years of age.

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 14/07/2023, Balearic Islands Ethics Committee (c/Calçat, 2A, 2n, Palma, 07011, Spain; +34 971177378; ceic_ib@caib.es), ref: IB 5219/23 PI

Study design

Multicenter parallel randomized clinical trial

Primary study design

Interventional

Study type(s)

Efficacy

Health condition(s) or problem(s) studied

Reduction of antibiotic prescription in primary care

Interventions

The method of randomization is simple random sampling.

General practitioners (GPs) in the intervention arm will receive individualized monthly information about their antibiotic prescription (incorporating graphical data, including antibiotic selection, dosages, treatment duration, and associated conditions), combined with an individualized, face-to-face intervention by a primary care pharmacist for GPs with significant potential for improvement in their prescribing practices (above the 75th percentile), along with an online educational component.

GPs in the control group will receive an intervention to reduce potentially inappropriate medication use (benzodiazepines, proton pump inhibitors and antipsychotics in patients with dementia) in patients over 65 years old.

The follow-up of the intervention is 12 months.

Intervention Type

Behavioural

Primary outcome(s)

Current primary outcome measure as of 06/01/2025:

The efficacy of the interventions on the prescription of antibiotics in the primary care area, measured using the number of antibiotic prescriptions extracted from the e-prescription databases of each health district at 12-month follow-up. The primary outcome will be measured as the total number of antibiotic prescriptions per 100 visits.

Previous primary outcome measure:

The efficacy of the interventions on the prescription of antibiotics in the primary care area, measured using the number of antibiotic prescriptions extracted from the e-prescription databases of each health district at 12-month follow-up

Key secondary outcome(s)

Current secondary outcome measures as of 07/05/2025:

1. Percentage of patients who start antibiotic treatment measured using the number of antibiotic prescriptions extracted from the e-prescription databases of each health district at 12-month follow-up.
2. Antibiotic prescription according to therapeutic group of antibiotics measured using number of antibiotic prescriptions according to therapeutic group (beta-lactams, glycopeptides, macrolides, aminoglycosides, tetracyclines, quinolones and fluoroquinolones, sulfonamides, lincosamides, oxazolidinones, polymyxins, rifamycins, nitroimidazoles, nitrofurans, fosfomycin) extracted from the e-prescription databases of each health district at 12-month follow-up.
3. Antibiotic prescription according to prescription pathology measured using the number of antibiotic prescriptions and their associated pathology (lower and upper respiratory infections, lower and upper urinary tract infections, sexually transmitted infections, skin and soft tissue infections, oral and odontogenic infections) at 12-month follow-up.
4. Antibiotic prescription according to gender measured using the number of antibiotic prescriptions extracted from the e-prescription databases of each health district, considering the gender of the GP and gender of the patient at 12-month follow-up.
5. The reach, engagement, adoption, fidelity and maintenance of the intervention. Reach will be defined as the percentage of participating GPs relative to the total number of eligible GPs. Engagement will measure the degree of active involvement in the intervention, including use of the feedback reports, participation in pharmacist-led sessions, and integration of recommendations into clinical practice. Adoption will be measured at the organizational level as the percentage of healthcare management units that have agreed to implement the

intervention. Fidelity to the intervention will be assessed through audits of adherence to the planned feedback delivery process. This includes consistency in the timing and content of reports, implementation of pharmacist consultations, and completion of online training modules. Maintenance of the intervention will be assessed at 6 and 12 months post-intervention, focusing on the degree to which the intervention becomes institutionalized within the healthcare system.

Previous secondary outcome measures:

1. Percentage of patients who start antibiotic treatment measured using the number of antibiotic prescriptions extracted from the e-prescription databases of each health district at 12-month follow-up.
2. Antibiotic prescription according to therapeutic group of antibiotics measured using number of antibiotic prescriptions according to therapeutic group (beta-lactams, glycopeptides, macrolides, aminoglycosides, tetracyclines, quinolones and fluoroquinolones, sulfonamides, lincosamides, oxazolidinones, polymyxins, rifamycins, nitroimidazoles, nitrofurans, fosfomycin) extracted from the e-prescription databases of each health district at 12-month follow-up.
3. Antibiotic prescription according to prescription pathology measured using the number of antibiotic prescriptions and their associated pathology (lower and upper respiratory infections, lower and upper urinary tract infections, sexually transmitted infections, skin and soft tissue infections, oral and odontogenic infections) at 12-month follow-up.
4. Antibiotic prescription according to gender measured using the number of antibiotic prescriptions extracted from the e-prescription databases of each health district, considering the gender of the GP and gender of the patient at 12-month follow-up.
5. The reach of the intervention as well as the fidelity of the intervention execution. The reach of the intervention will be measured using the percentage of patients assigned to GPs, with any antibiotic prescription and eligible to receive the intervention at 12 months follow-up. The fidelity will be measured using a comprehensive record with a detailed description of the execution process and any adaptations made in each intervention, as well as the strategies they comprise at the 12-month follow-up.

Completion date

12/12/2026

Eligibility

Key inclusion criteria

All GPs with assigned patients in their primary care health center will be included

Participant type(s)

Health professional

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

Does not meet the inclusion criteria

Date of first enrolment

01/05/2025

Date of final enrolment

31/12/2025

Locations

Countries of recruitment

Spain

Study participating centre

Balearic Islands Health Service (IB-Salut)

Carrer de l'Escola Graduada, 3, Centre

Palma

Spain

07002

Study participating centre

Paterna Health District (Conselleria de Salut Universal; Arnau de Vilanova-Llíria district)

Paterna

Spain

46980

Study participating centre

Tarragona-Reus Health district (Institut Català de la Salut; Tarragona-Reus district)

Tarragona

Spain

43480

Sponsor information

Organisation

Primary Care Management of Mallorca (Gerencia de Atención Primaria de Mallorca)

Funder(s)

Funder type

Government

Funder Name

Instituto de Salud Carlos III

Alternative Name(s)

SaludISCI, InstitutodeSaludCarlosIII, Instituto de Salud Carlos III | Madrid, Spain, Carlos III Institute of Health, Institute of Health Carlos III, Carlos III Health Institute, La misión del Instituto de Salud Carlos III (ISCI), ISCI

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

Spain

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be stored in a publicly available repository

IPD sharing plan summary

Stored in publicly available repository

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol article		19/07/2025	21/07/2025	Yes	No