

Effectiveness of surgical procedures for acute cranial expansion in traumatic brain injury

Submission date 02/01/2025	Recruitment status Recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 01/04/2025	Overall study status Ongoing	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 01/04/2025	Condition category Circulatory System	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Traumatic brain injury (TBI) patients often exhibit an increase in their intracranial volume due to blood collection or brain tissue edema. When the volume of any intracranial compartment exceeds a critical threshold, the compensatory mechanisms become overwhelmed, compromising intracranial compliance and blood supply, which leads to intracranial compartment syndrome (ICCS) and further exacerbates brain damage through secondary injury. When less invasive measures to counteract ICCS prove insufficient, cranial decompression is recommended, with decompressive craniectomy (DC) currently being the preferred technique. Although its effectiveness has been demonstrated, DC is also associated with a high incidence of complications. Expansive craniotomy (EC) has been proposed as an alternative that leverages the benefits of cranial decompression provided by DC while reducing the associated adverse effects. This study will compare the functional outcomes and complications linked to DC and EC.

Who can participate?

Adults aged 18 to 70 years old with TBI diagnosed with ICCS who require cranial decompression

What does the study involve?

In DC a cranial vault bone graft is removed and left out to be stored in a bone bank or an abdominal pouch created during the surgery. In EC the cranial vault bone graft is fixed in an elevated position 1 - 1.5 cm above the cranial rim by 3-5 metallic plates placed during surgery. Participants' progress will be observed during their hospital stay and assessed for 1 year through structured telephone follow-ups. The main outcomes include functional recovery and the rate of complications such as operative site infections, and/or reinterventions. By comparing these two methods, the study seeks to determine whether the expansion craniotomy improves outcomes for TBI patients while maintaining or enhancing the safety and reliability of the cranial decompression procedure.

What are the possible benefits and risks of participating?

Participants may benefit from a safer and more effective method of reducing ICP, especially if they are assigned to the expansion craniotomy group. The study provides access to expert surgical care and close monitoring throughout the recovery process. Contributions to this research could lead to improved treatment protocols for future TBI patients worldwide.

Both procedures carry inherent surgical risks, including infection and bleeding, among others, not related to the study per se. Because the indications for any of the two interventions will be the same for the original procedure at the discretion of the treating neurosurgeon. Participants will be required to attend follow-up assessments, which may involve some time and effort. All procedures will be performed by experienced neurosurgeons following established safety protocols. Participants will be closely monitored, and any complications will be promptly addressed. This study represents an important step toward advancing surgical techniques for managing severe TBI and improving the quality of care for patients worldwide.

Where is the study run from?

Meditech Foundation in Cali (Colombia)

When is the study starting, and how long is it expected to run for?

December 2024 to February 2027

Who is funding the study?

NTPlast S.r.l.

Who is the main contact?

Dr Andrés M. Rubiano MD, PhD(c), FACS, IFAANS, direccion@meditechhubcol.org, andresrubiano@aol.com

Study website

<https://space-tbi-study.com/>

Contact information

Type(s)

Scientific

Contact name

Dr Santiago Cardona

ORCID ID

<https://orcid.org/0000-0002-1446-0636>

Contact details

Calle 7A #44-103

Cali

Colombia

760036

(+57) 324 593 1186

cardona.santiago.meditechf@outlook.com

Type(s)

Principal Investigator

Contact name

Dr Andres M Rubiano

ORCID ID

<https://orcid.org/0000-0001-8931-3254>

Contact details

Calle 7A #44-103
Cali
Colombia
760036
+57 313 251 4132
andresrubiano@aol.com

Type(s)

Public

Contact name

Ms Wendy Gonzalez

ORCID ID

<http://orcid.org/0009-0000-7106-3568>

Contact details

Calle 7A # 44-103
Cali
Colombia
760036
+57 310 537 4949
gonzalez.wendy.meditechf@outlook.com

Type(s)

Scientific

Contact name

Dr Santiago Cardona

Contact details

Calle 7A #44-103
Cali
Colombia
760036
(+57) 324 593 1186
proyectos@meditechhubcol.org

Type(s)

Principal Investigator

Contact name

Dr Andres M Rubiano

Contact details

Calle 7a #44-103
Cali
Colombia

760036
(+57) 313 251 4132
direccion@meditechhubcol.org

Additional identifiers

EudraCT/CTIS number

Nil known

IRAS number

ClinicalTrials.gov number

Nil known

Secondary identifying numbers

Nil known

Study information

Scientific Title

A comparative-effectiveness study on expansion craniotomy versus decompressive craniectomy for surgical management of traumatic brain injury in patients with intracranial compartment syndrome

Acronym

E-SPACE-TBI

Study objectives

This study aims to determine the outcomes of decompressive craniectomy versus expansion craniotomy in the surgical management of patients with TBI and intracranial compartment syndrome

Ethics approval required

Ethics approval required

Ethics approval(s)

Approved 31/01/2025, Meditech Ethics and Research Committee (Calle 7A # 44-103, Cali, 760036, Colombia; +57 602 372 06 72; comitedeeticainvestigacion@meditechhubcol.org), ref: CEIM-2024-12-102

Study design

Prospective multicentric observational comparative-effectiveness study

Primary study design

Observational

Secondary study design

Comparative-effectiveness study

Study setting(s)

Hospital, Medical and other records, Telephone

Study type(s)

Efficacy

Participant information sheet

Not available in web format, please use contact details to request a participant information sheet

Health condition(s) or problem(s) studied

Intracranial compartment syndrome in traumatic brain injury patients

Interventions

The study will include patients with traumatic brain injury and intracranial compartment syndrome managed either by hemispheric expansion craniotomy or hemispheric decompressive craniectomy.

Enrollment:

Participants are screened upon admission to the emergency room after experiencing a traumatic brain injury (TBI). Eligibility is determined using clinical and imaging criteria, including abnormal brain CT scans and evidence of intracranial compartment syndrome (ICCS), which is defined as a primary intracranial injury generating a loss of intracranial compliance and generating risk for brain tissue hypoxia and/or hypoperfusion. Informed consent is obtained from participants or their legal representatives as many of these patients can arrive with a low level of consciousness due to the TBI.

Surgical interventions: participants are allocated to one of two groups:

This study will conduct surgical procedures for cranial decompression to treat intracranial compartment syndrome. Different techniques can achieve this. Currently, decompressive craniectomy (DC) is the most frequently used technique. It is favored by the Brain Trauma Foundation's current guidelines and will serve as a comparator for expansion craniotomy (EC). The allocation of patients to the intervention they will undergo will be based on the clinical criteria of the treating neurosurgeon as long as the patients meet the inclusion criteria and do not present any exclusion criteria. Neurosurgeons will be required to describe the reason(s) why the surgery was indicated and why the surgical approach was selected.

The EC will be performed through a retro-auricular "C-shape" incision and executing a 15x15 cm craniotomy with a "C-shape" durotomy. Dural closure will be at the discretion of the surgeon and the availability of resources, including duroplasty with aponeurotic galea, a suturable dural patch, a non-suturable dural patch, or simply a superficial dural cover with hemostatic materials like Surgicel® or Gelfoam®. The expansion craniotomy will be finished with a cranioplasty using 3-5 "Rialto" plates for bone graft closure.

There are multiple techniques for DC. The elected technique will be at the discretion of the treating neurosurgeon, but only fronto-parieto-temporal DC will be considered. The removed bone flap may be stored in an abdominal pouch in the right or left upper abdominal quadrants or by freezing in a bone or blood bank freezer.

Patients who receive an EC may subsequently be taken to DC if the expected improvement is not achieved with the intervention, according to the postsurgical images and invasive and non-invasive neuromonitoring, based on the criteria of the treating neurosurgeon. If this happens, the treating neurosurgeon must specify why they opted for the procedure.

On the other hand, the intervention cannot be changed in patients taken to DC once the surgery is completed since the bone would have already been removed and stored.

Follow-up and monitoring:

Observation During Hospitalization: Patients with this type of procedure, always go for postoperative management to intensive care units (ICUs), independent of the study. Follow-up is performed by ICU personnel and includes systemic multi-monitoring and evaluations for postoperative complications such as infections. They regularly require head imaging control with CT not specifically associated with the study (is an observational CEI study). The data collectors of the study will be collecting physiological variables daily from the clinical records of the ICU. Data monitoring will be performed by the central research team in permanent contact with the local investigators.

Long-Term Follow-Up: Data will be collected from the medical record until the patient is discharged or transferred and by telephone interview at 3, 6, and 12 months after the admission. The telephone follow-up will include applying the Glasgow Outcome Scale—Extended version (GOSE) and evaluating functional recovery.

Duration:

Total Observation: Participants will be observed continuously during their hospital stay.

Data Collection and Sources:

Data will be collected from the medical records.

Demographic and Clinical Data: Includes age, sex, baseline health, and mechanism of trauma.

Surgical Outcomes: we will include complications reported in the medical record or observed inside the ICU.

Functional Outcomes: Evaluated from the medical records and GOSE scores.

Data Sources: Data will be collected from medical records, direct observations, and after discharge by telephone interviews conducted by trained personnel.

Intervention Type

Procedure/Surgery

Primary outcome measure

Functional outcomes measured using the Glasgow Outcome Scale Extended at discharge, 3 months, 6 months, and 12 months after surgery

Secondary outcome measures

1. Control of the Intracranial Compartment Syndrome measured using ICP waveform via invasive and noninvasive devices and/or noninvasive neuromonitoring techniques (digital pupillometry, optic nerve sheath diameter ultrasonographic measurement, and transcranial Doppler) measured daily during ICU admission.
2. Frequency and type of neurosurgical interventions required measured using data collected from medical records when possible and telephone follow-up as an alternative until the end of the follow-up at 12 months after the initial surgery
3. Medical and surgical treatment intensity measured according to the categories of stratified management of intracranial pressure proposed in the study methodology that are based on international guidelines for the management of intracranial hypertension at discharge from the ICU
4. Surgical site complications measured using data collected from medical records when possible and telephone follow-up as an alternative until the end of the follow-up at 12 months after the initial surgery
5. ICU and in-hospital length of stay collected from medical records at hospital discharge

Overall study start date

27/12/2024

Completion date

01/02/2027

Eligibility

Key inclusion criteria

1. TBI patients arriving at the emergency room in the first 24 hours following trauma.
2. Abnormal CT scan, with a primary injury including any epidural, intracerebral or subdural collection with a midline shift >3 mm and any basal cistern compression with at least 2 abnormal findings in the initial evaluation at the emergency room including optic nerve ultrasound > 6 mm on the eye at the same side of the CT's primary injury or and/or an abnormal pupillometry with a reduced MCV in the pupil of the same side of the CT's primary injury, or/and a TCD with PI > 1.3 and/or MCA-DV < 20 cm/seg on the same side of the CT's primary injury or/and a P2>P1 waveform pattern in the same side of the CT's primary injury.
3. Age 18 to 70 years old.
4. Patients with or without polytrauma with survival expectancy >24 hours.
5. Cranial decompression or cranial expansion surgical procedures less than 24 hours after the trauma.

Participant type(s)

Patient

Age group

Mixed

Lower age limit

18 Years

Upper age limit

70 Years

Sex

Both

Target number of participants

292

Key exclusion criteria

1. TBI patients arriving at the emergency room after 24 hours following trauma.
2. Normal CT scan at the emergency room.
3. Abnormal CT scan at the emergency room with any primary injury and midline shift less than 3mm or without basal cistern compression and with normal values in at least two different modalities of assessing ICCS (pupillometry, optic nerve sheath ultrasound, transcranial Doppler and/or non-invasive ICP waveform analyzer.
4. Age less than 18 or more than 70 years old.
5. Polytrauma or massive brain injury with survival expectancy < 24 hours.
6. Cranial decompression or cranial expansion surgical procedures performed > 24 hours after the trauma.

Date of first enrolment

01/02/2025

Date of final enrolment

01/02/2026

Locations

Countries of recruitment

Bolivia

Brazil

Cambodia

Cameroon

Chile

Colombia

Dominican Republic

Ecuador

Egypt

Guatemala

Italy

Nigeria

Paraguay

Peru

Philippines

Rwanda

Serbia

Tanzania

Thailand

Venezuela

Study participating centre

Hospital Universitario del Valle Evaristo García
Calle 5 #36-00
Cali
Colombia
760042

Study participating centre
Hospital de Kennedy
Av. 1 de Mayo #40B-54
Bogota
Colombia
110871

Study participating centre
Clínica Santa Gracia
Cl. 14 N #9-15
Popayan
Colombia
190003

Study participating centre
Clinica Abel Gonzalez
Av. Independencia # 101
Santo Domingo
Dominican Republic
10109

Study participating centre
Hospital de Emergencia Jose Casimiro Ulloa
Av. Roosevelt 6355
Lima
Peru
150122

Study participating centre
Hospital Nacional Daniel Alcides Carrion
Av. Guardia Chalaca 2176
Callao
Peru
07001

Study participating centre

Hospital Belen de Trujillo

Jiron Bolivar 350

Trujillo

Peru

13001

Study participating centre

Hospital Miguel Pérez Carreño

Av. San Martín, a la altura de Maternidad

Caracas

Venezuela

1020

Study participating centre

Hospital Central de Maracay

Av. Las Delicias, Maracay

Maracay

Venezuela

2101

Study participating centre

Hospital Enrique Tejera

Av. Lisandro Alvarado

Valencia

Venezuela

2001

Study participating centre

Hospital das Clínicas da FMUSP

Av. Dr. Enéas de Carvalho Aguiar, 255

São Paulo

Brazil

05403-900

Study participating centre

Fundação Hospitalar Getúlio Vargas

Avenida Apuriná, número 4, en el barrio Praça 14 de Janeiro

Manaus

Brazil
69020

Study participating centre

Hospital Obrero No. 1

Calle Lucas Jaimes 76

La Paz

Bolivia

0000

Study participating centre

Hospital Regional San Juan de Dios

Calle Santa Cruz

Tarija

Bolivia

0000

Study participating centre

Hospital Clínico Mutual de Seguridad

Av. Libertador Bernardo O'Higgins

Santiago de Chile

Chile

4848

Study participating centre

Hospital Puerto Montt

Los Aromos 65, Puerto Montt

Los Lagos

Chile

5480000

Study participating centre

Hospital General San Juan de Dios

1ra Avenida "A" 10-50

Ciudad de Guatemala

Guatemala

01010

Study participating centre

Hospital Roosevelt
Calzada Roosevelt
Ciudad de Guatemala
Guatemala
01011

Study participating centre
Hospital del Trauma Prof. Dr. Manuel Giagni
Av Gral Máximo Santos
Asunción
Paraguay
1500

Study participating centre
Hospital de Especialidades Eugenio Espejo
Av. Gran Colombia
Quito
Ecuador
170403

Study participating centre
Hospital General Monte Sinai
Av. Gran Colombia
Quito
Ecuador
170515

Study participating centre
Hospital General Docente Ambato
Av. Pasteur y Unidad Nacional
Ambato
Ecuador
180104

Study participating centre
Hospital General Rodríguez Zambrano
Barrio Sta. Martha Calle 12 V-A San Mateo
Manta
Ecuador
130213

Study participating centre
University College Hospital
Queen Elizabeth Road, Ibadan, Oyo State
Ibadan
Nigeria
200212

Study participating centre
Rwanda Military Hospital
Street KK739ST, Kanombe, Distrito de Kicukiro
Kigali
Rwanda
250

Study participating centre
Garoua Regional Teaching Hospital
Boulevard Docteur Jamot, Garoua, Región Norte
Garoua
Cameroon
237

Study participating centre
El Qasr El Ainy Hospital
27 Nafezet Sheem El Shafaey St, Kasr Al Ainy
Cairo
Egypt
11562

Study participating centre
Muhimbili Orthopaedic Hospital
Kalenga Street, West Upanga,
Dar es Salaam
Tanzania
141111

Study participating centre
Bicol Regional Hospital and Medical Center
Rizal St. 4501
Daraga

Philippines
4501

Study participating centre

Techo Santepheap National Hospital

JRC6+JR Win Win Blvd, Phnom
Penh
Cambodia
12000

Study participating centre

Khon Kaen Hospital

54 Sri Chant Rd, Nai Mueang, Mueang Khon Kaen District
Khon Kaen
Thailand
40000

Study participating centre

Neurosurgical Clinic University Clinical Centre of Serbia

Dr Koste Todorovića 4
Belgrade
Serbia
11000

Study participating centre

Ozienda Ospedaliero – Universitaria Policlinico Umberto I

Viale del Policlinico, 155, 00161
Rome
Italy
00161

Study participating centre

Centro Diagnóstico de Especialistas

Cl. 13 #1175
Riohacha
Colombia
440009

Sponsor information

Organisation

Meditech Foundation

Sponsor details

Calle 7-A # 44-103, Cali, Colombia

Cali

Colombia

760036

+57 (602) 372 0672

asistenciafundacionmeditech@gmail.com

Sponsor type

Research organisation

Website

<https://www.meditechfoundationglobal.org/es>

Funder(s)**Funder type**

Industry

Funder Name

NTPlast

Results and Publications**Publication and dissemination plan**

Planned publication in a peer-reviewed journal.

Intention to publish date

01/02/2028

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available upon request from MR Dr Santiago Cardona (cardona.santiago.meditecf@outlook.com). Individual participant data (IPD) will be stored in a private repository and made available upon request. Once requested, the unprocessed data collected on the RedCap platform will be shared as soon as possible. The shared data will be anonymous, as the only identifier will be the ID assigned to patients participating in the study. No additional consents are required for sharing IPD, as this possibility is included in the informed consent form signed by patients or their next of kin during their involvement in the study.

IPD sharing plan summary

Stored in non-publicly available repository, Available on request