The effects on asthma of dietary supplements that induce the growth or activity of beneficial microorganisms.

Submission date	Recruitment status	[X] Prospectively registered
27/03/2019	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
04/04/2019	Completed	Results
Last Edited	Condition category	[] Individual participant data
05/04/2019	Respiratory	Record updated in last year

Plain English summary of protocol

Background and study aims

Asthma is an inflammatory disorder that causes the narrowing of airways, increased production of mucus and therefore difficulty breathing. Asthma sufferers may experience shortness of breath, wheezing, tightness of the chest and a dry irritating cough. Asthma is a common condition that affects approximately 300 million people worldwide, in the UK 5.4 million people are currently recieving treatment. The NHS spend up to £1 billion a year for the care and treatment of those who have asthma.

This study will investigate how prebiotic supplementation affects asthma, by observing changes in quality of life, use of current medication and inflammation. Prebiotics are a carbohydrate that feed the 'good bacteria' in the gut, otherwise known as Probiotics. These bacteria have an important role in look after our immune system and a reduced number/activity of these 'good bacteria' may lead to development of asthma or increased severity of asthma. This study aims to increase our understanding of how the gut bacteria may affect asthma and the results may lead to a better understanding of the factors that might influence asthma and how these can be managed by targeting the diet.

Who can participate?
Adults diagnosed with asthma

What does the study involve?

Participants are randomly allocated into one of two groups. Group one are given a prebiotc supplement, known as B-GOS. Group two are given a placebo (dummy pill). Both groups will take the supplement, B-GOS, or placebo for 3 weeks. After this time, participants will continue as normal for two weeks, followed by another 3 weeks with either the placebo or B-GOS. Each participant will attend the laboratory for a lung function test, blood sample and to fill out questionnare before the start of each treatment; a total of 4 times over 8 weeks.

What are the possible benefits and risks of participating? Participants may benefit from gaining a wider understanding into managing and controlling their asthma. Potential risks include possible discomfort when taking blood samples.

Where is the study located? Nottingham Trent University, Clifton Campus (UK)

When does the study start and end? June 2019 to June 2020

Who is the main contact.
Mr. Jacob Jayaratnasingam
Jacob.Jayaratnasingam2014@my.ntu.ac.uk

Contact information

Type(s)

Scientific

Contact name

Mr Jacob Jayaratnasingam

ORCID ID

https://orcid.org/0000-0002-6497-5531

Contact details

Nottingham Trent University Clifton Campus Clifton Lane Clifton Nottingham United Kingdom NG11 8NS +44 (0)115 941 8418 jacob.jayaratnasingam2014@my.ntu.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Integrated Research Application System (IRAS)

253339

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

IRAS Application: 253339

Study information

Scientific Title

The effects of Prebiotic Supplementation on Quality of Life, Control of Asthma, and Markers of Systemic Inflammation in Adults with Asthma. A Double-Blind, Placebo-Controlled, Crossover Trial.

Acronym

PSA

Study objectives

Prebiotic supplementation will lead to significant changes in pro-inflammatory/anti-inflammatory markers in the blood.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approval pending, East Midlands - Nottingham 1 Research Ethics Committee (The Old Chapel, Royal Standard Place, Nottingham, NG1 6FS; 0207 104 8104; NRESCommittee.EastMidlands-Nottingham1@nhs.net), ref: 19/EM/0088

Study design

Single centre double-blind placebo-controlled intervention crossover trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Asthma, steps 1-4 according to the Birtish Thoracic Society Guidelines (BTS)

Interventions

Block randomisation will be used to randomise subjects into equal groups. At baseline, both groups will give a blood sample, undergo lung function tests, complete a 4-day food diary and asthma related questionnaires. The same measurements will be completed following prebiotic or placebo supplementation after 3-weeks. The measurements will then be completed again following a 2-week washout period. Groups will then crossover research arms and continue a 3 week supplementation period with either the prebiotic or placebo, before the final measurement. Supplements will be randomised by an independent party into 'X' and 'Y', ensuring the study is double blind.

Blood samples will be analysed using flow cytometry and mass spectrometry to observe specific and general changes in protein expression.

Intervention Type

Supplement

Primary outcome(s)

Changes in FOXP3 expression by regulatory T cells in adults with asthma, following a prebiotic intervention crossover trial. Measured using flow cytometry at 4 timepoints as aforementioned, baseline, prebiotic, washout and placebo.

Key secondary outcome(s))

- 1. Changes in asthma control, and frequency of reliever medication in adults with asthma following prebiotic supplementation. Measured at 4 time points, baseline, prebiotic, washout and placebo; using the Asthma Control Questionnaire.
- 2. Changes in perception of asthma control (e.g. reduced frequency/severity of symptoms) following prebiotic supplementation. Measured at 4 time points, baseline, prebiotic, washout and placebo; using the Asthma Control Test.
- 3. Changes in quality of life following prebiotic supplementation. Measured at 4 time points, baseline, prebiotic, washout and placebo; using the Asthma Quality of Life Questionnaire

Completion date

24/06/2020

Eligibility

Key inclusion criteria

- 1.18-50 years of age at the date of your first visit.
- 2. Body mass index (BMI) of 18.5-25 kg·m2 (we can work this out for you using your height and body weight).
- 3. Non-smoker.
- 4. Asthma is defined as Steps 1 to 4 based on British Thoracic Society guidelines.
- 5. On a stable asthma treatment for 3 months
- 6. Current medication prescription from your GP if diagnosed with asthma (e.g. maintenance and reliever inhalers).
- 7. Be able and willing to follow all trial requirements.
- 8. Disclose any nutritional supplements you take to the researcher, to determine whether these may be considered as 'exclusion criteria'.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

50 years

Sex

All

Key exclusion criteria

- 1. Suffer from asthma but do not have a current medication prescription from your GP (e.g. maintenance and reliever inhalers).
- 2. Regularly consume Omega-3 supplements, and/or eat high levels of Omega-3 (e.g. more than 1-2 portions of oily fish such as salmon or mackerel a week).
- 3. Take aspirin or other non-steroidal anti-inflammatory drugs such as ibuprofen once a day on 5 of the 7 days of the week.
- 4. Pregnant or planning a pregnancy during the time of the study (on each visit you will be asked to complete a pregnancy test if you are within childbearing age).
- 5. Consumed prebiotics and/or probiotics (supplements), drugs that affect gastrointestinal mobility or laxatives in the 4 weeks before signing the consent form.
- 6. Daily dose of anti-histamine, which you could not temporarily avoid for 72 hours before each testing session without exacerbation of symptoms.
- 7. Vegetarian or vegan diet.
- 8. Previously diagnosed with chronic obstructive pulmonary disease (COPD), emphysema, chronic bronchitis, or similar respiratory (breathing-related) illness.
- 9. Admitted to hospital during the past 12 months for your asthma.
- 10. History of heart failure, pulmonary hypertension, embolism, or other pulmonary heart disease.
- 11. History of recurrent chest infections.
- 12. Acute infection in the last four weeks, and/or major operation in the past four months.
- 13. History of gastrointestinal drug reaction.
- 14. Taken antibiotics in the past 3 months.
- 15. History or current evidence of gastrointestinal disease (e.g. chronic constipation, diarrhoea, irritable bowel syndrome, Chrohn's disease).
- 16. Recently taken part in other research projects. Please notify the chief investigator.
- 17. Lactose intolerant.
- 18. Regularly take antioxidant supplements, such as beta-carotene, vitamin A, vitamin C, vitamin E, lutein and selenium. Standard multivitamin and mineral supplements are acceptable; however, If a single antioxidant supplement (e.g. Vitamin C), is more than the recommended daily DRV's this must be checked with the chief investigator.

Date of first enrolment 24/06/2019

Date of final enrolment 24/06/2020

Locations

Countries of recruitment

United Kingdom

England

Study participating centre
Nottingham Trent University Clifton Campus
Nottingham Trent University Clifton Campus
Clifton Lane

Clifton Nottingham Nottingham United Kingdom NG11 8NS

Sponsor information

Organisation

Nottingham Trent University

ROR

https://ror.org/04xyxjd90

Funder(s)

Funder type

University/education

Funder Name

Nottingham Trent University

Alternative Name(s)

NTU

Funding Body Type

Private sector organisation

Funding Body Subtype

Universities (academic only)

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

The current data sharing plans for this study are unknown and will be available at a later date

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output typeDetailsDate createdDate addedPeer reviewed?Patient-facing?HRA research summary28/06/2023NoNoParticipant information sheet11/11/202511/11/2025NoYes