A group interpersonal psychotherapy intervention for caregivers of children with nodding syndrome to improve the mental health of both

| Submission date | Recruitment status No longer recruiting | Prospectively registered | | |
|-------------------------------------|--|--|--|--|
| 27/05/2016 | | ☐ Protocol | | |
| Registration date 08/06/2016 | Overall study status Completed | Statistical analysis plan | | |
| | | [X] Results | | |
| Last Edited 10/07/2023 | Condition category Mental and Behavioural Disorders | [] Individual participant data | | |

Plain English summary of protocol

Background and study aims

Nodding syndrome (NS) is a recent, little-known disease that occurs only in children and only in small regions in South Sudan, Tanzania and northern Uganda. It was first described in 1962. The disease causes affected children to have so-called nodding seizures, often triggered by eating or when they feel cold. The seizures are short lived and stop once the trigger is removed (i.e. they stop eating or become warm again). Over time, the condition leads to a permanent stunting of growth and cognitive development and leads to mental handicap. It is not known what causes the disease, but it may be associated with being infested with a parasitic worm called Onchocerca volvulus. There is no known cure for the disease and so the available treatments concentrate on alleviating symptoms; this includes the taking if anticonvulsant drugs, for example phenobarbitol. Looking after children with NS can cause significant strain on the family looking after them and it can cause the mental health of both sufferers and their caregivers (for example, parents). The study aims to design and investigate the effectiveness of a family based, group interpersonal psychotherapy treatment (IPT-F) for caregivers of patients with Nodding Syndrome (NS) on the mental health of both the caregivers and patients compared to usual care (provided for in the national NS response plan).

Who can participate?

Adults (aged at least 18) caring for a child with NS.

What does the study involve?

Participants are randomly allocated to one of two groups. Those in group 1 are allocated to the control group and are given usual care. Those in group 2 are allocated to the intervention group. Participants in this group take part in interviews and group discussions with other caregivers. The information gained from these is then used to help design the intervention. The IPT-F intervention will be tested in Pader district, northern Uganda and involves one and a half hour group sessions with caregivers held weekly over a period of 12 weeks. It is delivered by trained village health team members who are supervised by trained health workers from the local health

facility. All participants are then followed up during the course of a year to see what affect the intervention has had on their mental wellbeing.

What are the possible benefits and risks of participating? Not provided at time of registration

Where is the study run from?
Makerere University School of Public Health, Makerere University (Uganda)

When is the study starting and how long is it expected to run for? April 2013 to September 2014

Who is funding the study?
Grand Challenges Canada (Canada)

Who is the main contact? Dr Byamah Mutamba byamamutamba@yahoo.com

Contact information

Type(s)

Scientific

Contact name

Dr Byamah Mutamba

ORCID ID

https://orcid.org/0000-0002-9430-443X

Contact details

Butabika Hospital Uganda Kampala Uganda P.O.Box 7017 0772609671 byamamutamba@yahoo.com

Additional identifiers

Protocol serial number S4 0232-01

Study information

Scientific Title

Effectiveness of a family based group interpersonal psychotherapy intervention (IPT-F) for caregivers of children with nodding syndrome to improve the mental health of both in comparison to usual care

Acronym

IPT-F

Study objectives

In comparison to the control, the culturally sensitive IPT-F intervention will alleviate psychological distress symptoms and improve the mental health of caregivers leading to improved child-parent relationships and mental health of the children with Nodding syndrome (NS).

Ethics approval required

Old ethics approval format

Ethics approval(s)

Makerere University College Of Health Sciences, School Of Public Health. Higher Degrees Research and Ethics Committee, 14/06/2013, ref: IRB00011353

Study design

Single centre quasi experimental (intervention cohort) study design

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Common mental disorders

Interventions

There are two arms in the trial; intervention and control. To obtain the participants in the intervention arm, caregivers are consecutively selected based on updated lists of caregivers from the local health facility NS (Nodding syndrome) register at Atanga Health centre and from Village Health Team (VHT) members resident in selected study parishes of Atanga sub-county. The participants for the control arm will be randomly chosen from an updated list of caregivers resident in selected study parishes in Awere sub-county.

The group interpersonal psychotherapy intervention (IPT-F) will involve weekly group sessions each lasting 1 and a half hours delivered by trained village health team (VHT) members for a period of 12 weeks. The VHT members will be supervised by trained health workers from the local health facility.

Caregivers in the control parishes will access the usual care as provided for in the national response plan for NS. The national response plan for NS includes health education about NS for affected communities, management of patients with NS, caregiver education and counselling at health facilities and designated treatment centres, outreach clinics and home visits to families affected by the syndrome.

Patients in both study arms will continue to receive the usual care during and after the intervention period.

The activities conducted during each of the 12 IPT-F weekly sessions depends on which session it is in the 12 week schedule.

During the first phase (sessions 1 and 2), the facilitator:

- 1. Helps group members develop rapport among themselves, feel comfortable in the group so that they can talk about how they feel and what is happening in their lives
- 2. Explains that everyone in the group has depression and can help one another
- 3. Explains that because of depression they may not feel that they can do as much in their lives as they had in the past, but that as they feel better they'll get back to their usual ways
- 4. Explains what will happen during the group sessions
- 5. Reviews everyone's depression symptoms using the translated version of primary health questionnaire-version 9.
- 6. Reminds each one of their particular problem areas and the connection to depression symptoms.
- 7. Prompts group members to talk about the interpersonal problem that is triggering the depression, and finds out how the person has been dealing with the problem during the previous week

During the middle phase (sessions 3-10) the facilitator:

- 8. Continues to make group members feel comfortable within the group and aims at ensuring that all members have an opportunity to talk during the session
- 9. Links each person's depression symptoms to current life problem
- 10. Helps members listen to each other and tries to get each one of them to offer ideas that could be useful to another group member in dealing with a problem
- 11. Helps members act in a caring way towards one other
- 12. Continues to encourage members to be hopeful about realising changes in their life that will make them feel better

During the termination phase (sessions 11-12), the facilitator encourages each member:

- 13. To talk about any life changes and their feelings about these changes
- 14. To talk about their depression: what symptoms have gone and what symptoms remain
- 15. To talk about what worries them about the future
- 16. To talk about what has not changed in their life and how they feel about that.and
- 17. To talk about how they feel about the cessation of the group meetings

Intervention Type

Behavioural

Primary outcome(s)

Proportion of caregivers with improved mental health status (a decrease in proportion of caregivers with depression) after one year follow up, assessed using mini plus neuropsychiatric interview (depression module)

Key secondary outcome(s))

- 1. The proportion of caregivers with:
- 1.1. Generalised anxiety disorder (GAD) measured using mini plus neuropsychiatric interview (GAD module.
- 1.2. Post traumatic stress disorder (PTSD) measured using mini plus neuropsychiatric interview (PTSD module)
- 1.3. A high risk of suicide measured using mini plus neuropsychiatric interview(suicide module)
- 1.4. High levels of psychological distress measured using the self-report questionnaire
- 1.5. Low levels of functioning measured using the assessment of functioning questionnaire

- 1.6. Low levels of social support measured using the social support scale
- 1.7. High levels of stigma measured using the devaluation of consumer's scale
- 2. Proportion of affected children with:
- 2.1. An abnormal total difficulties score measured using the strengths and difficulties questionnaire
- 2.2. High parent attachment scores measured using the inventory of parent and peer attachment questionnaire
- 2.3. Depression measured using the depression self-rating scale
- 2.4. Separation anxiety disorder measured using the mini kid neuropsychiatric interview
- 2.5. Generalised anxiety disorder measured using the mini kid neuropsychiatric interview
- 2.6. Attention deficit hyperactivity disorder measured using the mini kid neuropsychiatric interview
- 2.7. Psychological trauma measured using the revised child impact of events scale

Completion date

30/09/2014

Eligibility

Key inclusion criteria

- 1. A caregiver will be included in the study if she/he is an adult ≥18 years of age
- 2. Is living in the selected study parishes
- 3. Is a primary caregiver for a patient (child/adolescent) with Nodding Syndrome
- 4. Gives informed written consent
- 5. Gives assent for and presents with one of their NS-affected children for assessment

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

142

Key exclusion criteria

- 1. Is severely ill and unable to participate in the study
- 2. Does not give informed written consent to participate in the study

Date of first enrolment

02/09/2013

Date of final enrolment

Locations

Countries of recruitment

Uganda

Study participating centre
Makerere University School of Public Health
Makerere University
Kampala
Uganda
P.O.Box 7072

Sponsor information

Organisation

Makerere University

ROR

https://ror.org/03dmz0111

Organisation

Uganda National Council for Science and Technology

Funder(s)

Funder type

Government

Funder Name

Grand Challenges Canada

Alternative Name(s)

Grands Défis Canada, gchallenges, Grand Challenges Canada / Grands Défis Canada, grandchallengescanada, GCC

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

Canada

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

| Output type | Details | Date created | Date added | Peer reviewed? | Patient-facing? |
|-------------------------------|-------------------------------|--------------|------------|----------------|-----------------|
| Results article | | 28/06/2018 | | Yes | No |
| Results article | | 15/02/2018 | 10/07/2023 | Yes | No |
| Participant information sheet | Participant information sheet | 11/11/2025 | 11/11/2025 | No | Yes |