

# Assessment of lung water in patients with low blood pressure after infection

<b>Submission date</b> 16/01/2015	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 04/02/2015	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 09/10/2018	<b>Condition category</b> Circulatory System	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Extravascular lung water refers to fluid within the lung. Increase in extravascular lung water is associated with death in very ill patients. The purpose of this study is to compare three methods of lung ultrasound with transpulmonary thermodilution technique.

### Who can participate?

Individuals with septic shock, 18 years old or over

### What does the study involve?

All patients will be assessed with three methods of lung ultrasound (total B-line scores, BLUE points and scanning eight regions) and the results will be compared with the transpulmonary thermodilution technique.

### What are the possible benefits and risks of participating?

A benefit is improvement in the monitoring of the patients. The risks are bleeding, haematoma and pneumothorax.

### Where is the study run from?

Phramongkutklao Hospital (Thailand)

### When is study starting and how long is it expected to run for?

From March 2013 to July 2014

### Who is funding the study?

Phramongkutklao Hospital Foundation Under Her Royal Highness Princess Maha Chakri Sirindhorn Patronage (Thailand)

### Who is the main contact?

Dr. Pattarin Pirompanich

## Contact information

**Type(s)**

Public

**Contact name**

Dr Pattarin Pirompanich

**ORCID ID**

<https://orcid.org/0000-0003-0181-5394>

**Contact details**

Division of Pulmonary and Critical Care Medicine  
Department of Internal Medicine  
Faculty of Medicine  
Thammasat University  
Klong 1  
Klong Luang  
Pathumthani  
Thailand  
12120

**Additional identifiers****Protocol serial number**

N/A

**Study information****Scientific Title**

Comparison of three methods of lung ultrasound with the transpulmonary thermodilution technique for assessment of extravascular lung water in patients with septic shock

**Study objectives**

Extravascular lung water index greater than 10 mL/kg in patients with cardiogenic pulmonary oedema correlates with pulmonary capillary wedge pressure of greater than 20 mm Hg. Measurement of extravascular lung water needs sophisticated tools and use of the invasive transpulmonary thermodilution technique. By contrast, multiple B-lines by transthoracic portable ultrasound have been recently proposed to correlate with increased extravascular lung water in patients with cardiogenic pulmonary oedema.

Lung ultrasound has a good correlation with transpulmonary thermodilution technique for the assessment of extravascular lung water in patients with septic shock.

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

Institutional review board of the Royal Thai Army Medical Department, 22/04/2009  
Reference number: R032h/56

**Study design**

Cross sectional study

**Primary study design**

Observational

**Study type(s)**

Diagnostic

**Health condition(s) or problem(s) studied**

Septic shock

**Interventions**

1. Total B-line scores

2. BLUE points

3. Scanning eight regions

These three methods of lung ultrasound were used on each patient to assess extravascular lung water and the results were compared with those from obtained with the transpulmonary thermodilution technique.

**Intervention Type**

Device

**Primary outcome(s)**

Best method of lung ultrasound to diagnose pulmonary oedema in patients with septic shock: cutoff point of extravascular water index measured by transpulmonary thermodilution greater than

10 mL/kg predicted bodyweight, which is correlated with a pulmonary capillary wedge pressure of greater than 20 mmHg to indicate pulmonary oedema

**Key secondary outcome(s))**

Cutoff point of total B-line scores to predict extravascular lung water index of 10 mL/kg or more

**Completion date**

30/07/2014

## **Eligibility**

**Key inclusion criteria**

1. Age 18 years old or over

2. Septic shock requiring central venous catheterisation and arterial catheterisation for continuous pressure monitoring

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

18 years

**Sex**

All

**Key exclusion criteria**

1. Valvular heart disease
2. Pregnancy
3. Acute or history of pulmonary embolism
4. Lung resection surgery
5. Acute respiratory distress syndrome
6. Interstitial lung diseases
7. Multilobar pneumonia
8. Pleural diseases

**Date of first enrolment**

01/09/2013

**Date of final enrolment**

30/06/2014

**Locations****Countries of recruitment**

Thailand

**Study participating centre**

Phramongkutklao Hospital

315 Phayathai Road

Ratchathewi

Bangkok

Thailand

10400

**Sponsor information****Organisation**

Phramongkutklao Hospital Foundation Under Her Royal Highness Princess Maha Chakri Sirindhorn Patronage

**Funder(s)**

**Funder type**

Charity

**Funder Name**

Phramongkutklao Hospital Foundation Under Her Royal Highness Princess Maha Chakri  
Sirindhorn Patronage

## Results and Publications

**Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Stored in repository

**Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	01/09/2018		Yes	No