Evaluating the effectiveness and costeffectiveness of Tobacco Cessation services in Swedish primary healthcare targeting socioeconomically disadvantaged areas in Stockholm - the "Motivation 2 Quit" (M2Q) study

Submission date 07/01/2016	Recruitment status No longer recruiting	[X] Prospectively registered		
		[X] Protocol		
Registration date 11/01/2016	Overall study status Completed	Statistical analysis plan		
		[X] Results		
Last Edited	Condition category	Individual participant data		
02/12/2022	Mental and Behavioural Disorders			

Plain English summary of protocol

Background and study aims

The aim of this study is to test whether a new method to help people quit their tobacco use is better than the methods currently offered by primary health care providers (i.e. family doctors) in Sweden. The method specifically targets primary healthcare centres in Stockholm where tobacco use is more common, where the local population are likely to be poorly paid, not be as well-educated and suffer from poorer health when compared to the general population. The study also aims to determine whether this new method is worth its cost.

Who can participate?

Adult Swedish or Arabic speaking patients with Swedish social security numbers and permanent resident permits, without cognitive impairment, who use tobacco on a daily basis, visit participating primary healthcare centres and who are not already undergoing treatment for tobacco cessation.

What does the study involve?

The primary healthcare centres participating in this study are assigned to either the intervention or control group. Participants who attend centres in the control group are given their usual treatment. This includes counselling and drug (nicotine replacement) therapy. Participants who attend centres in the intervention group are treated according to the "tobacco cessation on prescription" programme. The main difference between the treatment for this group and the control group is that the counselling and drug therapy services offered are individually prescribed. They are also given information on other ways to help them quit their tobacco use (for example, doing exercise and finding other ways to cope with withdrawal symptoms) and self-help resources (such as smartphone apps, web-based counseling and websites that have more information and support). All participants are followed up after 6 and 12 months to see whether they have quit their tobacco use.

What are the possible benefits and risks of participating?

Participants will be offered support to quit their tobacco use. Quitting tobacco use can be a stressful experience since it can cause short term withdrawal symptoms and psychological stress. However, counselling on coping strategies and therapies that can help ease withdrawal symptoms will be given to avoid discomfort for participants. Participants that successfully quit their tobacco use should experience an improvement in both short and long-term health and quality of life.

Where is the study run from? 14-20 primary healthcare centres located in socioeconomically disadvantaged areas in Stockholm County (Sweden)

When is the study starting and how long is it expected to run for? January 2015 to August 2019

Who is funding the study?

- 1. Stockholm County Council (Sweden)
- 2. Skandia Life Insurance (Sweden)
- 3. The Public Health Agency of Sweden (Folkhälsomyndigheten) (Sweden)

Who is the main contact?

1. Dr Tanja Tomson (scientific) tanja.tomson@ki.se

2. Miss Anne Leppänen (public) anne.leppanen@ki.se

Contact information

Type(s)

Scientific

Contact name

Dr Tanja Tomson

ORCID ID

https://orcid.org/0000-0002-4577-4304

Contact details

Karolinska Institute (Karolinska Institutet) Tomtebodavägen 18A Stockholm Sweden 17177 +46 (0)852 480 173 tanja.tomson@ki.se

Type(s)

Public

Contact name

Miss Anne Leppänen

ORCID ID

https://orcid.org/0000-0003-4273-4072

Contact details

Karolinska Institute (Karolinska Institutet) Tomtebodavägen 18A Stockholm Sweden 17177 +46 (0)852 483 612 anne.leppanen@ki.se

Additional identifiers

Protocol serial number

N/A

Study information

Scientific Title

Evaluating the effectiveness and cost-effectiveness of Tobacco Cessation on Prescription in Swedish primary healthcare targeting socioeconomically disadvantaged areas in Stockholm - the "Motivation 2 Quit" (M2Q) study, a cluster-randomised controlled trial

Acronym

M2Q

Study objectives

Compared to standard treatment tobacco cessation on prescription is an effective and costeffective treatment in achieving 7-day abstinence from tobacco use at 6 months after intervention among tobacco users visiting primary healthcare centres located in socioeconomically disadvantaged areas in Stockholm.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Regional Ethical Review Board in Stockholm, April 2015, ref: 2015/207-31, 2015-1226-32, 2016 /2080-32

Study design

An interventional pragmatic cluster-randomised controlled trial.

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Smoking

Interventions

Participants are randomly allocated to either tobacco cessation on prescription or standard treatment according to which primary healthcare centre (PHC) they attend.

- 1. Tobacco cessation on prescription is defined as:
- 1.1. Tobacco cessation counselling (minimum 10 minutes) provided by a qualified healthcare professional in combination with
- 1.2. A prescription for individualised tobacco cessation treatment, including options for:
- 1.2.1. Further counselling (referral to healthcare provider with more competence or the Swedish Quit Smoking Line)
- 1.2.2. Pharmacotherapy (nicotine replacement therapy, varenicline, bupropion)
- 1.2.3. Other measures for tobacco cessation (physical activity and other strategies to cope with withdrawal symptoms)
- 1.2.4. Follow-up (by telephone or revisit)
- 1.2.5. Support for self-management (reference to mobile applications, web-based counselling and websites for more information and support). The approach will be individualised in the sense that providers will discuss the available treatment options, contraindications, preferences and other relevant circumstances with the patient and then decide together on which treatment alternative(s) suit the individual best.
- 3. Follow-up of the prescription by the prescriber on at least one occasion is also included in the intervention.

Standard treatment is defined as:

Current treatment practices for tobacco cessation at the participating PHC in the control group and include different types of counselling and pharmacotherapy. The major difference between the trial conditions is how the counselling is administered (with or without a prescription form).

Participating PHC centres will be randomised with a 1:1 ratio to either intervention- or control conditions. Cluster-randomisation will be employed at the PHC centre level, meaning that all study participants recruited from a particular PHC centre will receive the same treatment (tobacco cessation treatment either with or without tobacco cessation on prescription).

Intervention Type

Mixed

Primary outcome(s)

Point prevalence of 7-day abstinence (total abstinence from tobacco use during the 7 days preceding follow-up) at 6 months after the intervention

Key secondary outcome(s))

- 1. Point prevalence of 7-day abstinence at 12 months after the intervention
- 2. Daily tobacco consumption (number of cigarettes), at 6 and 12 months after the intervention
- 3. Number of quit attempts and health-related quality of life (on a scale from 0-1 where 0 represents death and 1 represents perfect health) at 6 and 12 months after the intervention
- 4. Cost effectiveness, measured as the incremental cost per quality-adjusted life year

All outcomes will be based on self-reports from patient questionnaires.

Completion date

31/12/2019

Eligibility

Key inclusion criteria

- 1. Swedish or Arabic speaking daily tobacco users with Swedish social security number and permanent resident permit
- 2. Over 18 years of age
- 3. Visiting participating primary health care centres in the study

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

250

Key exclusion criteria

- 1. Ongoing treatment for tobacco cessation
- 2. Cognitive impairment affecting ability to participate in the study on a voluntary basis

Date of first enrolment

01/02/2016

Date of final enrolment

31/08/2018

Locations

Countries of recruitment

Sweden

Study participating centre Valsta Medical Centre (Valsta Vårdcentral)

Hammargatan 6

Märsta

Study participating centre Norrtälje Northern Medical Centre (Norrtälje Norra Vårdcentral)

Lasarettsgatan 6 Norrtälje Sweden 761 45

Study participating centre Hallstaviks Medical Centre (Hallstaviks Vårdcentral)

Carl Wahrens väg 24 Hallstavik Sweden 763 34

Study participating centre

Capio Primary Care Centre Hagsätra (Capio Vårdcentral Hagsätra)

Hagsätra torg 7B Bandhagen Sweden 124 73

Study participating centre Capic Primary Care Centre Wasa (Capi

Capio Primary Care Centre Wasa (Capio Vårdcentral Wasa) Prästgårdsvägen 4 Södertälie

Sweden 151 61

Study participating centre

Cevita Care - Rimbo-Edsbro family doctors (Husläkarna Rimbo-Edsbro)

Stockholmsvägen Rimbo Sweden 762 31

Study participating centre

Capio Primary Care Centre Skogås (Capio Vårdcentral Skogås)

Melodivägen 6 Skogås Sweden 142 40

Study participating centre

Capio Primary Care Centre Vårberg (Capio Vårdcentral Vårberg)

Vårbergstorget 5 Skärholmen Sweden 127 43

Study participating centre Handens Medical Centre (Handens Vårdcentral)

Dalarövägen 6 Handen Sweden 136 46

Study participating centre Family Doctors Husby (Familjeläkarna Husby)

Edvard Griegsgången 13 Kista Sweden 164 32

Study participating centre Hallonbergens Medical Centre (Hallonbergens Vårdcentral)

Hallonbergsplan 7 Sundbyberg Sweden 174 52

Study participating centre Fisksätra Medical Centre (Fisksätra Vårdcentral)

Fisksätra torg 20 Saltsjöbaden Sweden 133 41

Study participating centre The Health Medical Centre Tensta (Hälsans Vårdcentral Tensta)

Tenstagången 18 Spånga Sweden 163 64

Study participating centre Liljeholmens Medical Centre (Liljeholmens Vårdcentral)

Liljeholmstorget 7 Stockholm Sweden 117 63

Study participating centre Rinkeby Medical Centre (Rinkeby Vårdcentral)

Skårbygränd 3 Spånga Sweden 163 72

Study participating centre Vårbergs Medical Centre (Vårbergs Vårdcentral)

Vårbergsplan 31 Skärholmen Sweden 127 43

Study participating centre Skärholmens Medical Centre (Skärholmens Vårdcentral)

Storholmsgatan 19 Skärholmen Sweden 127 48

Study participating centre Solna City Medical Centre (Solna Centrum Vårdcentral)

Hotellgatan 3 Solna

Sponsor information

Organisation

Karolinska Institute (Karolinska Institutet) (Sweden)

ROR

https://ror.org/056d84691

Funder(s)

Funder type

Government

Funder Name

Stockholms Läns Landsting

Alternative Name(s)

Stockholm County Council

Funding Body Type

Government organisation

Funding Body Subtype

Local government

Location

Sweden

Funder Name

Skandia Life Insurance (Livförsäkringsbolaget Skandia)

Funder Name

The Public Health Agency of Sweden (Folkhälsomyndigheten)

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are not expected to be made available because the trialists do not have consent to share this information.

IPD sharing plan summary

Not expected to be made available

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		01/12/2022	02/12/2022	Yes	No
<u>Protocol article</u>	protocol	16/09/2016		Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes