Trial of online support for primary care patients tapering off antidepressant medication

Submission date	Recruitment status Recruiting	Prospectively registered		
04/04/2022		☐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
05/05/2022	Ongoing	Results		
Last Edited	Condition category	Individual participant data		
15/07/2025	Mental and Behavioural Disorders	[X] Record updated in last year		

Plain English summary of protocol

Background and study aims

The use of antidepressants (ADs) is increasing globally, including within Australia, which has one of the highest rates of AD prescribing. Despite clear benefits for many people, there is reason to believe that the ongoing use of these medications is often not properly monitored or stopped (deprescribed) when a person returns to better Mental health. This trial sets out to test how well an online support tool (WiserAD) can help patients and their general practitioner to manage the careful and appropriate reducing and stopping of antidepressants, in primary care patients.

Who can participate?

Patients aged 18-75 years, who have been stable on their ADs for <=12 months, with no or mild depressive symptoms and no history of recurrent depression.

What does the study involve?

All participants will receive a phone call from the study team one week after randomisation. Participants allocated to the intervention arm will be provided with a personal login code for the study web portal and encouraged to complete the initial components of the intervention (Assess and Advise) which seek to determine participants' current support and management strategies and help them to understand their specific antidepressants. They will then complete the third component which contains three sub-sections to assist in creating a personal plan to help them cease their ADs: i) Prepare – management strategies for withdrawal symptoms and opportunities to discuss the plan with their GP or trusted mental health worker; ii) Schedule -Selecting a start time to begin tapering; iii) Share – Print out of the personalised action plan to keep and share with supportive family and/or friends. Participants will also be required to complete a daily check-in through the portal which will check current symptoms and highlight any negative changes in emotional wellbeing, they will also receive texts reminders to complete these tasks. Participants allocated to the usual care arm will receive usual care plus attention control which comprises a link to the beyondblue website and directed to the AD factsheet. Participants will not receive any instructions regarding their personal AD use (i.e. they will not be advised to cease or continue). Both arms will receive reminders to complete their research follow up measures online (or by phone if they prefer) at 3, 6, 12, 18 and 24 months.

What are the possible benefits and risk of participating?

WiserAD will provide AD users with the opportunity to reflect on their emotional health and well-being and current need for antidepressant medication. This will also help us to find out if the WiserAD support tool is effective. We do not anticipate any risks, side-effects, or discomforts. However, if patients are asked to taper (reduce down) their medication as part of the trial, we will monitor how well they are coping with this through a daily update from the WiserAD tool and the dosage may be adjusted accordingly, in consultation with a GP, who will remain their responsible clinician.

Where is the study run from? The University of Melbourne, Department of General Practice, Australia.

When is the study starting and how long is it expected to run for? March 2018 to December 2026

Who is funding the study? National Health and Medical Research Council (NHMRC; ID no.: 1157337)

Who is the main contact?
Dr Cath Kaylor-Hughes
wiserad-enquiries@unimelb.edu.au

Contact information

Type(s)

Public

Contact name

Ms Amy Coe

ORCID ID

https://orcid.org/0000-0003-3723-7645

Contact details

Level 2, 780 Elizabeth St Melbourne Australia 3004 +61 3 90356335 amy.coe@unimelb.edu.au

Type(s)

Scientific

Contact name

Dr Cath Kaylor-Hughes

ORCID ID

https://orcid.org/0000-0003-3353-4108

Contact details

Level 2, 780 Elizabeth St Melbourne Australia 3004 +61 383444921 cath.kaylorhughes@unimelb.edu.au

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

NCT05355025

Protocol serial number

ACTRN126220000567729, NHMRC ID no. 1157337

Study information

Scientific Title

WiserAD: A randomised trial of a structured online intervention to promote and support antidepressant deprescribing in primary care

Acronym

The WiserAD Study

Study objectives

WiserAD (a novel, structured approach to deprescribing antidepressants) will be more effective than usual practice in enabling GPs to help primary care patients to cease (or decrease) their antidepressant medication whilst maintaining patient mental health and wellbeing.

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 21/07/2021, University of Melbourne Human Research Ethics Committee (HREC) (Level 5, Alan Gilbert Building, 161 Barry Street, Carlton, 3010, Australia; +61 03 8344 1376; HumanEthics-Enquiries@unimelb.edu.au), ref: 20558

Approved 12/08/2021, University of Melbourne Human Research Ethics Committee (HREC, Office of Research Ethics and Integrity, University of Melbourne, VIC 3010, Australia; +61 8344 1376; HumanEthics-Enquiries@unimelb.edu.au), ref: 20558

Study design

Single-blind parallel-arm randomized controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Decreasing antidepressant (AD) medication for depression

Interventions

All participants will receive a brief <10 minute check-in phone call from the study team one week after randomisation to make sure that they have access to the intervention or attention control.

Participants will be provided with a link and personal login code for the study web portal which has been purposely built for the trial. Participants will receive the code via automated email and will be encouraged to complete the initial components of the intervention (Assess and Advise) which seek to determine participants' current support and management strategies and help them to understand their specific antidepressants (ADs) (approx 10 minutes). They will then complete the third component which contains three sub-sections to assist in creating a personal plan to help them cease their ADs:

- i) Prepare management strategies for withdrawal symptoms and opportunities to discuss the plan with their GP or trusted mental health worker;
- ii) Schedule Selecting a start time to begin tapering;
- iii) Share Print out of the personalised action plan to keep and share with supportive family and /or friends (approx 5 minutes total to completed).

At this point, participants will also be asked to share their tapering schedule with their GP for modification if necessary. Participants will asked to complete their appointment with their GP within two weeks. Once the tapering schedule has been approved, participants will log back into the web portal and will be required to complete a daily check-in through the portal which will check current symptoms and highlight any negative changes in emotional wellbeing (approx 10 minutes), they will also receive texts reminders to complete these tasks. Apart from the reminder texts and email communications, all information will be provided to participants via the WiserAD web portal.

Participants will have access to the WiserAD online support tool for the duration of the study (minimum of 3-months post randomisation and until last data collected from the last participant, therefore up to 18-months), however they will be encouraged to begin tapering their ADs within two weeks of receiving approval of their tapering schedule. Use of the WiserAD online support tool will be monitored using purpose built analytics within the WiserAD web portal (unblinded study team access only).

Randomisation:

The treatment to which a participant is assigned will be determined by a computer generated pseudo-random code using random permuted blocks of varying size, created by the web developers and held on a secure server. Computer generated pseudo-random code using random permuted blocks of varying size, created by the web developers and held on a secure server. Participants will be allocated with equal probability to each treatment arm and stratified by site (GP clinic).

Intervention Type

Behavioural

Primary outcome(s)

Proportion of patients successfully ceasing ADs at 6 months where successful cessation is defined as no AD use (as measured by the Resource Use Questionnaire (RUQ)) and the absence of clinically significant depressive symptoms (as measured by the Patient Health Questionnaire 9-item (PHQ-9)). Measured at 3-, 6-, 12-, 18-months post-baseline. Primary outcome is at 6-months post baseline.

Key secondary outcome(s))

- 1. Depressive symptoms (PHQ-9) Measured at baseline, 3-, 6-, 12-, 18months.
- 2. Anxiety symptoms (GAD-7) Measured at baseline 3-, 6-, 12-, 18-months.
- 3. Patient Activation (PAM) Measured at baseline, 3-, 6-months.
- 4. Quality of Life (AQoL-4D) Measured at baseline, 3-, 6-, 12-, 18-months.
- 5. Health Service Use (Resource Use Questionnaire) Measured at baseline, 3-, 6-, 12-, 18-months.
- 6. Beliefs About Medication Questionnaire (BMQ) Measured at baseline, 3-months.
- 7. Signs and Symptoms (A study specific 3-item text response measure for patient reported withdrawal symptoms) Measured at 3-, 6-months.
- 8. User Engagement Scale-Short Form (UES-SF) Measured at 3-, 6-months.
- 9. Accountability Measurement Tool (AMT) Measured at 3-, 6-months.
- 10. Medical Benefit Scheme (MBS; health service use) and the Pharmaceutical Benefit Scheme (PBS; pharmaceutical use) data collected at
- 11. the completion of the study (for the whole duration of participation in the study (18-months)).
- 12. Proportion of patients successfully ceasing ADs where successful cessation is defined as no AD use (as measured by the Resource Use Questionnaire (RUQ)) and the absence of clinically significant depressive symptoms (as measured by the Patient Health Questionnaire 9-item (PHQ-9))Measured at 3-, 12-, 18-months post baseline.

Completion date

31/12/2026

Eligibility

Key inclusion criteria

- 1. 18-75 years
- 2. Stable on AD for >=12m: (no depressive episodes)
- 3. No history of recurrent depression
- 4. Sufficient English language proficiency to provide informed consent
- 5. No or mild depressive symptoms as measured on the Personal Health Questionnaire (PHQ9)
- 6. Low risk of Suicide or Self-harm
- 7. Agree to consider reviewing their AD use
- 8. Agree to be randomized into the study
- 9. Willing to provide informed consent

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

75 years

Sex

All

Key exclusion criteria

- 1. Those currently experiencing a major life event in the next 3 months
- 2. Currently using ADs for any other health condition (other than depression)
- 3. Currently using non-SSRI or SNRI ADs, antipsychotics, or other mood stabiliser medication
- 4. Have no daily access to the internet

Date of first enrolment

25/04/2022

Date of final enrolment

31/12/2025

Locations

Countries of recruitment

Australia

Study participating centre Parkville Precinct Medical

1F Royal Parade Parkville Australia 3050

Sponsor information

Organisation

University of Melbourne

ROR

https://ror.org/01ej9dk98

Funder(s)

Funder type

Government

Funder Name

National Health and Medical Research Council

Alternative Name(s)

National Health and Medical Research Council, Australian Government, NHMRC National Health and Medical Research Council, NHMRC

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

Australia

Results and Publications

Individual participant data (IPD) sharing plan

IPD will not be available publicly due to confidential information being collected as part of the trial. Data collected as part of the trial will be for research purposes only.

IPD sharing plan summary

Not expected to be made available

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Other publications		13/02/2024	15/02/2024	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes