Interval exercise versus continuous exercise in patients with moderate to severe chronic obstructive pulmonary disease

| Submission date | Recruitment status No longer recruiting | Prospectively registered | |
|-------------------------------|---|--------------------------------|--|
| 08/06/2004 | | [X] Protocol | |
| Registration date | Overall study status | Statistical analysis plan | |
| 08/06/2004 | Completed | [X] Results | |
| Last Edited 20/10/2016 | Condition category Respiratory | [] Individual participant data | |
| 20/10/2010 | Respiratory | | |

Plain English summary of protocol

Background and study aims

Chronic obstructive pulmonary disease is a disease of the lungs that is usually caused by smoking. In COPD, lung damage gradually worsens over time. People with COPD are short of breath and sometimes cough and wheeze. Symptoms slowly get worse over time, but most patients also have intermittent bouts of worsening of symptoms. Lung infections, cold weather, and exertion may bring on these bouts. In addition to medications, the treatment guidelines recommend high-intensity, continuous exercise to improve lung function. However, this type of exercise can be difficult for people with COPD to tolerate. Some believe that short intervals of exercise may also benefit patients with COPD and may be easier for them to accomplish. The aim of this study is to find out whether interval exercise produces the same benefits for people with COPD and is easier for them to tolerate than continuous exercise.

Who can participate?
Adult patients with COPD

What does the study involve?

Participants are randomly allocated to attend 12 to 15 supervised sessions of either continuous or intermittent high-intensity exercise over 3 weeks followed by unsupervised exercise at home. Participants used exercise bikes during the supervised exercise sessions. Participants allocated to continuous exercise warmed up for 2 minutes, pedaled at high intensity for 20 minutes (or until they needed to stop because of fatigue or other symptoms), and cooled down for 2 minutes. Participants in the intermittent exercise group warmed up for 2 minutes, pedaled for 20 minutes, alternating between 20 seconds at high intensity and 40 seconds at low intensity (or until they needed to stop because of fatigue or other symptoms), and cooled down for 2 minutes. Participants completed a standard questionnaire to assess their lung function at the start of the study and 5 weeks into the study. The researchers also collected information on the number of times participants had to take unplanned breaks during the supervised exercise sessions.

What are the possible benefits and risks of participating? Not provided at time of registration

Where is the study run from? AstraZeneca (Switzerland)

When is the study starting and how long is it expected to run for? May 2004 to September 2005

Who is funding the study?

- 1. AstraZeneca (Switzerland)
- 2. The Helmut Horten Foundation

Who is the main contact? Dr Milo Puhan milo.puhan@evimed.ch

Contact information

Type(s)

Scientific

Contact name

Dr Milo Puhan

Contact details

Horten Centre
University of Zurich
Postfach Nord
University Hospital
Zurich
Switzerland
8091
+41 1 255 87 09
milo.puhan@evimed.ch

Additional identifiers

Protocol serial number

N/A

Study information

Scientific Title

Interval exercise versus continuous exercise in patients with moderate to severe chronic obstructive pulmonary disease

Study objectives

The aim of this study is to assess if interval exercise compared to high intensity continuous exercise is not of inferior effectiveness in terms of health-related quality of life (HRQL) and

exercise capacity improvements but associated with better exercise tolerance in patients with moderate to severe COPD at the beginning of a respiratory rehabilitation.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics Committee of the Kantonsspital Aarau, Aargau, Switzerland

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Chronic obstructive pulmonary disease (COPD)

Interventions

Group 1: 12-15 sessions of high intensity continuous exercise

Group 2: 12-15 sessions of interval exercise

Between group comparisons in terms of changes of clinical outcomes (health-related quality of life, functional exercise capacity and subjective patient experience of exercise) and physiological outcomes (exercise tests with gas exchange and ventilatory variables) during respiratory rehabilitation.

Intervention Type

Behavioural

Primary outcome(s)

Health-related quality of life (HRQL), measured by the Chronic Respiratory Questionnaire (CRQ) two weeks after the end of rehabilitation

Key secondary outcome(s))

Secondary endpoints include additional clinical outcomes such as functional exercise capacity, other HRQL measures, patients' experience of physical exercise as well as physiological measures of the effects of physical exercise such as cardiopulmonary exercise testing.

Completion date

30/09/2005

Eligibility

Key inclusion criteria

COPD with Global Initiative for Chronic Obstructive Lung Disease (GOLD) stage III and IV admitted to an inpatient respiratory rehabilitation

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

Not provided at time of registration

Date of first enrolment

01/05/2004

Date of final enrolment

30/09/2005

Locations

Countries of recruitment

Switzerland

Study participating centre University of Zurich

Zurich Switzerland 8091

Sponsor information

Organisation

AstraZeneca Switzerland

ROR

https://ror.org/034rhks82

Funder(s)

Funder type

Industry

Funder Name

AstraZeneca

Alternative Name(s)

AstraZeneca PLC, Pearl Therapeutics, AZ

Funding Body Type

Government organisation

Funding Body Subtype

For-profit companies (industry)

Location

United Kingdom

Funder Name

Helmut Horten Foundation

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Study outputs

| Output type | Details | Date created Date added | Peer reviewed? | Patient-facing? |
|-------------------------------|-------------------------------|-------------------------|----------------|-----------------|
| Results article | results | 05/12/2006 | Yes | No |
| <u>Protocol article</u> | protocol | 13/08/2004 | Yes | No |
| Participant information sheet | Participant information sheet | 11/11/2025 11/11/2025 | No | Yes |