

Effects of the amount of SARS-CoV-2 in the maternal airways on outcomes before, during, and after childbirth

Submission date 31/10/2021	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 08/11/2021	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 20/12/2023	Condition category Pregnancy and Childbirth	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection has interested a relevant number of pregnant women worldwide with negative consequences both on the maternal and neonatal side. For instance, COVID-19 is more severe in pregnant women, preterm delivery is more common and neonates have a higher chance to need intensive care admission. Finally, SARS-CoV-2 mother-to-child transmission is possible and neonatal COVID-19 can be observed although rarely. Despite the viral load has been linked to the clinical severity of COVID-19 in non-pregnant patients, the relationship between SARS-CoV-2 viral load and perinatal outcomes in pregnant women affected by COVID-19 in late pregnancy is unknown. We sought to investigate whether or not nasopharyngeal SARS-CoV-2 viral load has any effect on perinatal outcomes, when COVID-19 is diagnosed in the third trimester of pregnancy.

Who can participate?

All women affected by COVID-19 during the third trimester of pregnancy (and not meeting exclusion criteria) were eligible.

What does the study involve?

Patient records from the start of the pandemic to mid-2021 will be analysed.

What are the possible benefits and risks of participating?

The study is retrospective, does not modify the routine care in any way and does not carry any risk for the patients.

Where is the study run from?

Hôpital "A. Bécclère" (France)

The study is conducted in four academic tertiary referral perinatal center in major EU and UK cities. These centers have been chosen through contacts between peers for the availability of electronic databases of pregnant women affected by COVID-19 and their expertise on perinatal COVID-19 research.

When is the study starting and how long is it expected to run for?

March 2020 to August 2021

Who is funding the study?

The study has no sponsor of any type and no honorarium is previewed for the participation to the study. Authors are performing the study for free during their worktime and they do not have any conflict of interest to disclose in relation to the project.

Who is the main contact?

Prof. Daniele De Luca, dm.deluca@icloud.com

Contact information

Type(s)

Scientific

Contact name

Prof Daniele De Luca

ORCID ID

<https://orcid.org/0000-0002-3846-4834>

Contact details

Service de Pédiatrie et Réanimation Néonatale,
Hôpital "A. Béclère", GHU Paris Saclay - APHP

157 rue de la Porte de Trivaux

Clamart

France

92140

+33 (0)145374837

DANIELE.DELUCA@APHP.FR

Type(s)

Public

Contact name

Prof Daniele De Luca

Contact details

Service de Pédiatrie et Réanimation Néonatale,
Hôpital "A. Béclère", GHU Paris Saclay - APHP

157 rue de la Porte de Trivaux

Clamart

France

92140

+33 (0)145374837

dm.deluca@icloud.com

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

Nil known

Study information

Scientific Title

Effect of SARS-CoV-2 estimated nasopharyngeal viral load on perinatal outcomes in pregnant women affected by COVID-19 during the third trimester (VALOROUS study)

Acronym

VALOROUS

Study objectives

To verify if maternal viral load in the airways is associated with perinatal outcomes in women affected by COVID-19 in the third trimester of pregnancy.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 11/11/2020, Ethical committee of the Department of Women and Newborn Health (Paris Saclay University Hospitals, APHP, Paris, France; +33 0492034409; cppsudmediterraneeV@chu-nice.fr), ref: CPP Sud Méditerranée n.2020-A00924-35

Study design

Multicenter international observational retrospective cohort study

Primary study design

Observational

Study type(s)

Other

Health condition(s) or problem(s) studied

Effect of SARS-CoV-2 estimated nasopharyngeal viral load on perinatal outcomes in pregnant women affected by COVID-19 during the third trimester

Interventions

Nasopharyngeal swabs will be obtained following US Center for Disease Control and Prevention guidelines. Extraction and amplification will be performed with commercial assays validated for SARS-CoV-2 diagnosis by WHO or local healthcare authorities. Manufacturer's recommendations will always be followed. The SARS-CoV-2 load will be estimated for any viral gene, according to each laboratory protocol. RT-PCR technique will be performed according to European Center for Disease Prevention and Control.

The following data will be collected: basic maternal and neonatal demographics, birth weight Z-score, time between COVID-19 diagnosis and delivery, 5' Apgar score, cord pH, estimated viral load for any vital gene, COVID-19 severity. These variables were chosen in order to keep a pragmatic design and make from the different centers easy to merge.

Data will be collected retrospectively to the start of the pandemic and prospectively to the end of the study.

Intervention Type

Other

Primary outcome(s)

1. Estimated viral load measured at birth using the number of cycles at the RT-PCR on maternal nasopharyngeal swab according to best microbiological practice
2. Gestational age at the birth measured using patient records
3. Birth weight and its Z-score measured using patient records
4. 5' Apgar score measured at birth by adequately trained midwives or neonatologists
5. Cord pH measured at birth using potentiometric point of care method on cord arterial puncture

Key secondary outcome(s)

There are no secondary outcome measures

Completion date

31/08/2021

Eligibility

Key inclusion criteria

1. Third trimester of pregnancy
2. Diagnosis of COVID performed according to World Health Organization criteria
3. Viral load estimated by real-time polymerase chain reaction cycle threshold (Ct) for any viral gene

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Female

Total final enrolment

393

Key exclusion criteria

1. Fetal congenital malformations
2. Major genetic or chromosomal abnormalities
3. Ongoing pregnancies at the time of the analysis
4. Missing viral load or outcome data

Date of first enrolment

01/03/2020

Date of final enrolment

31/08/2021

Locations

Countries of recruitment

United Kingdom

England

Belgium

France

Italy

Study participating centre

Department of Women and Newborn Health, Medical Center A. Bécclère, Paris Saclay University Hospitals, APHP

157, Rue de la porte de Trivaux

Clamart

France

92140

Study participating centre

Fetal Medicine Unit, Saint George's Hospital

Blackshaw Road

London

United Kingdom

SW17 0QT

Study participating centre

Department of Translational Medical Sciences, University of Naples Federico II

5, Via Sergio Pansini

Naples

Italy

80131

Study participating centre

Department of Obstetrics and Gynecology, Brugmann University Hospital, Université Libre de Bruxelles

4, Place A. Van Gehuchten

Brussels

Belgium

1020

Sponsor information

Organisation

Hôpital Antoine-Béclère

ROR

<https://ror.org/04sb8a726>

Funder(s)

Funder type

Other

Funder Name

Investigator initiated and funded

Results and Publications

Individual participant data (IPD) sharing plan

Anonymous raw data will be available on reasonable request.

Data can be asked to investigators in each of the participating centers (for the data contributed by each center). Data will be totally anonymous. A restricted duration of availability and other restrictions may apply according to local regulations enforced in each center.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		05/03/2023	20/12/2023	Yes	No
Protocol file			02/11/2021	No	No