# Reducing distress from somatic symptoms

Submission date	Recruitment status  No longer recruiting	<ul><li>Prospectively registered</li></ul>		
24/02/2015		☐ Protocol		
Registration date	Overall study status Completed	Statistical analysis plan		
13/05/2015		☐ Results		
<b>Last Edited</b> 12/08/2020	<b>Condition category</b> Mental and Behavioural Disorders	Individual participant data		
		[ ] Record updated in last year		

### Plain English summary of protocol

Background and study aims

Somatic symptom disorders are mental disorders characterized by physical symptoms that suggest physical illness but which cannot be explained fully by a general medical condition. A new brief psychological therapy may help people worry less. The aim of this study is to test how people find this therapy and whether it works. The researchers want to see whether it reduces worry, especially worries about health and illness. If it is shown to work then it will be possible to use the therapy in NHS services.

Who can participate?

Adults meeting criteria for somatic symptom disorders

### What does the study involve?

Participants receive a 'worry intervention' in 6 sessions over 8 weeks, either at home or at the GP surgery. The worry reduction strategies include psycho-education about worry, reviewing of beliefs about worry, increasing awareness of the initiation of worry and identification of individual triggers, use of worry periods, substituting problem-solving in place of worry and relaxation exercises. There are homework exercises between sessions. At the end of the study, participants are invited to take part in another interview to talk about their experience of the therapy sessions and suggest improvements. Participants are paid £15 for each assessment session, and travel expenses are paid.

What are the possible benefits and risks of participating?

It is hoped that participants benefit from the opportunity to engage in the worry intervention. The main possible burden is time taken to complete the questions.

Where is the study run from? South East England (UK)

When is the study starting and how long is it expected to run for? June 2014 to August 2015

Who is funding the study? University of Southampton (UK)

Who is the main contact? Prof. David Kingdon D.Kingdon@soton.ac.uk

### Contact information

### Type(s)

Scientific

### Contact name

Prof David Kingdon

#### Contact details

University Department of Psychiatry Academic Centre, College Keep 4-12 Terminus Terrace Southampton United Kingdom SO14 3DT +44 (0)2380718520 D.Kingdon@soton.ac.uk

### Additional identifiers

### Protocol serial number

12798

## Study information

#### Scientific Title

Reducing worry in patients with somatic symptom disorder using brief cognitive behavioural therapy

#### Acronym

**REDRESS** 

### Study objectives

To determine the acceptability and explore the clinical effectiveness of a worry-based intervention for severe and distressing health concerns.

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

- 1. University of Southampton Ethics and Research Governance Committee (ERGO12798) National Research Ethics Service, 17/02/2014, ref: 14/SC/0040
- 2. Southern Health NHS Foundation Trust, 27/05/2014, ref: SHT120
- 3. Sussex Partnership NHS Foundation Trust, 01/05/2014, ref: 5038-2014

### Study design

Observational case series

### Primary study design

Observational

### Study type(s)

Treatment

### Health condition(s) or problem(s) studied

Somatic symptom disorder

#### **Interventions**

Patients with Somatic Symptom Disorder will be offered 6 sessions of individual manualised psychological intervention focussed on worry. A control group is not included in this design.

The worry intervention will be provided in 6 sessions over 8 weeks. Intervention sessions will take place at home or at the participant's GP surgery, to minimise travel and burden for the participant. The worry reduction strategies included are indicated in the anxiety literature to be effective at reducing worry and do not challenge the illness interpretation. The main techniques are psycho-education about worry, reviewing of beliefs about worry, increasing awareness of the initiation of worry and identification of individual triggers, use of worry periods, substituting problem-solving in place of worry and relaxation exercises. Homework exercises are set between sessions. Written information is provided for patients (see appendix B). With patients' consent, sessions will be taped for assessment of adherence and for competence.

At the end of sessions, a consultation letter will be developed with the patient and shared with the patients' referring clinician and their general practitioner.

### Intervention Type

Other

### Primary outcome(s)

Health Anxiety Inventory (Salkovskis, Rimes, Warwick & Clark, 2002). Assessments will take place at baseline and 8 weeks.

### Key secondary outcome(s))

- 1. Intolerance of Uncertainty Questionnaire (Freeston, Rheaume, Letarte, Dugas & Ladouceur (1994)
- 2. Penn State Worry Questionnaire (Meyer, Miller, Metzger & Borkovec, 1990) Hospital Anxiety and Depression Scale (Zigmond & Snaith, 1983)
- 3. EQ-5D (Rabin & de Charro, 2001)
- 4. Short Warwick-Edinburgh Mental Well-being Scale (SWEMWBS, Tennant, Hiller, Fishwick, Platt, Weich, Parkinson et al, 2007)
- 5. A question around the acceptability of a referral for psychological therapy, developed for this study
- 6. Number of attendences at A&E.
- 7. Outpatients and General Practice will also be recorded
- 8. New physical diagnoses will also be noted.

The scales will provide comparative measures of symptoms, functioning and well being with some estimate of resource utilisation.

Assessments will take place at baseline and 8 weeks.

### Completion date

31/08/2015

## **Eligibility**

### Key inclusion criteria

- 1. Over 18 years old
- 2. Meet criteria for Somatic Symptom Disorder

### Participant type(s)

**Patient** 

### Healthy volunteers allowed

No

### Age group

Adult

### Lower age limit

18 years

#### Sex

All

### Key exclusion criteria

- 1. Under 18 years old
- 2. Not meeting criteria for Somatic Symptom Disorder
- 3. Having an insufficient understanding of English to read and complete the study questionnaires
- 4. Having a primary diagnosis of drug or alcohol misuse
- 5. Taking part in current Cognitive Behavourial Therapy

### Date of first enrolment

17/06/2014

#### Date of final enrolment

07/01/2015

### Locations

#### Countries of recruitment

United Kingdom

England

Study participating centre
Southern Health NHS Foundation Trust
United Kingdom
SO40 2RZ

Study participating centre
Sussex Partnership NHS Foundation Trust
United Kingdom
BN13 3EP

## Sponsor information

### Organisation

University of Southampton

#### **ROR**

https://ror.org/01ryk1543

## Funder(s)

### Funder type

University/education

### **Funder Name**

University of Southampton

### Alternative Name(s)

University of Southampton UK

### **Funding Body Type**

Government organisation

### **Funding Body Subtype**

Universities (academic only)

#### Location

**United Kingdom** 

## **Results and Publications**

## Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not expected to be made available

### **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			28/06/2023	No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes