# Is two-dimensional ultrasound necessary six weeks after insertion of the 52-mg levonorgestrel-releasing intrauterine system (LNG-IUS) contraceptive?

Submission date 29/07/2021	Recruitment status No longer recruiting	Prospectively registered		
		[X] Protocol		
Registration date 07/09/2021	Overall study status Completed	Statistical analysis plan		
		[X] Results		
<b>Last Edited</b> 07/11/2023	Condition category Urological and Genital Diseases	[] Individual participant data		

## Plain English summary of protocol

Background and study aims

The correct position of the contraceptive levonorgestrel intrauterine system (LNG-IUS) can be confirmed by two-dimensional ultrasound (2DUS). However, best the timing of a routine 2DUS to confirm the correct position is subject to discussion. The main aim of this study was to assess how often the LNG-IUS is found to be incorrectly positioned on 2DUS during routine investigation at six weeks after insertion in the gynecological outpatient department. Additionally, the study will investigate the relation between malposition of the LNG-IUS and clinical symptoms like unacceptable bleeding patterns and pelvic pain.

## Who can participate?

Women seeking a 52mg LNG-IUS (Mirena®), over 18 years of age, for contraception or treatment of heavy menstrual bleeding.

## What does the study involve?

2DUS was performed to check the position of the IUD immediately after insertion and six weeks later. Women were excluded if they did not attend the appointment for 2DUS six weeks after insertion. Women were asked to complete questionnaires six weeks after insertion and were asked about bleeding pattern and pelvic pain.

What are the possible benefits and risks of participating? No additional benefits or risks to participants are anticipated.

Where is the study run from? Máxima Medical Centre (Netherlands)

When is the study starting and how long is it expected to run for? May 2015 to December 2018

Who is funding the study? Investigator initiated and funded

Who is the main contact?
Patty van der Heijden, pattyvanderheijden@gmail.com

# Contact information

## Type(s)

Scientific

#### Contact name

Miss Patty van der Heijden

#### **ORCID ID**

https://orcid.org/0000-0002-5290-7646

## Contact details

Elkerliek Hospital Wesselmanlaan 25 Helmond Netherlands 5707HA +31 615699305 pvanderheijden@elkerliek.nl

# Additional identifiers

## Clinical Trials Information System (CTIS)

Nil known

# ClinicalTrials.gov (NCT)

Nil known

## Protocol serial number

15.025

# Study information

#### Scientific Title

Is two-dimensional ultrasound (2DUS) necessary six weeks after insertion of 52-mg LNG-IUS?

# **Study objectives**

In this prospective cohort study we assessed the additional value of 2DUS six weeks after insertion. We described the frequency of malposition and the relationship between malposition of the LNG-IUS and clinical symptoms like bleeding pattern and/or pelvic pain.

# Ethics approval required

Old ethics approval format

## Ethics approval(s)

Approved 23/05/2015, Dagelijks Bestuur METC (Máxima Medisch Centrum, Locatie Veldhoven, Postbus 7777, 5500 MB Veldhoven; 3140-8888525; m.rutten@mmc.nl), ref: 15.025

## Study design

Single-centre observational study

## Primary study design

Observational

# Study type(s)

Diagnostic

## Health condition(s) or problem(s) studied

Diagnostics of malposition and complaints in women with 52-mg LNG-IUS

#### **Interventions**

Immediately after insertion of the LNG-IUS, the position of the LNG-IUS was assessed with the 2DUS by the resident, gynaecologist or physician assistant who performed the insertion. Six weeks after insertion a 2DUS was performed to evaluate the position of the LNG-IUS. Participants were asked to complete questionnaires 6 weeks after insertion. In the questionnaires women were asked about their bleeding pattern (presence of bleeding and/or spotting: 'no bleeding', 'regular menstruation', 'sometimes a day of spotting (maximum of once a week)', 'heavy menstrual bleeding', 'several days a week bleeding days', 'several days a week spotting days', 'continuously spotting', 'complete irregular cycle'). Women were asked to rate pelvic pain on a five-point Likert scale.

## Intervention Type

Drua

#### Phase

Not Applicable

# Drug/device/biological/vaccine name(s)

levonorgestrel

## Primary outcome(s)

Incidence of malposition on 2DUS during routine investigation six weeks after insertion

# Key secondary outcome(s))

Clinical symptoms at six weeks:

- 1. Unacceptable bleeding pattern measured using a questionnaire
- 2. Pelvic pain measured using a 5-point Likert scale

## Completion date

31/12/2018

# **Eligibility**

# Key inclusion criteria

- 1. Women who visited our clinic for the insertion of a 52mg LNG-IUS (Mirena®) for contraception or treatment of heavy menstrual bleeding
- 2. Over 18 years of age

## Participant type(s)

Patient

## Healthy volunteers allowed

No

## Age group

Adult

## Lower age limit

18 years

#### Sex

Female

## Total final enrolment

500

## Key exclusion criteria

- 1. Presence of intracavitary polyps or fibroids on 2DUS prior to insertion
- 2. Unprotected intercourse one month prior to insertion if insertion took place outside menstruation
- 3. Women were excluded for analyses if no ultrasound was performed six weeks after insertion

### Date of first enrolment

01/03/2015

## Date of final enrolment

31/12/2016

# Locations

## Countries of recruitment

Netherlands

## Study participating centre Máxima Medical Centre

de Run 4600 Veldhoven Netherlands 5504DB

# **Sponsor information**

## Organisation

Máxima Medisch Centrum

## **ROR**

https://ror.org/02x6rcb77

# Funder(s)

# Funder type

Other

## Funder Name

investigator initiated and funded

# **Results and Publications**

# Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request. (pattyvanderheijden@gmail.com)

## IPD sharing plan summary

Available on request

# **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		04/10/2022	07/11/2023	Yes	No
Participant information sheet	In Dutch		10/08/2021	No	Yes
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Protocol file	In Dutch		10/08/2021	No	No