

# Waiting list controlled trial of surgical treatment for adults with chronic constipation

|  |   |   |
|--|---|---|
| <b>Submission date</b><br>30/09/2015   | <b>Recruitment status</b><br>No longer recruiting | <input type="checkbox"/> Prospectively registered<br><input checked="" type="checkbox"/> Protocol |
| <b>Registration date</b><br>30/09/2015 | <b>Overall study status</b><br>Completed          | <input type="checkbox"/> Statistical analysis plan<br><input checked="" type="checkbox"/> Results |
| <b>Last Edited</b><br>27/10/2022       | <b>Condition category</b><br>Digestive System     | <input type="checkbox"/> Individual participant data  |

## Plain English summary of protocol

### Background and study aims

Constipation is a common condition which affects most people in their lifetime. Chronic constipation is where the sufferer has been affected for at least six months, and non-surgical treatments such as medications (laxatives), exercise and diet changes are ineffective. A laparoscopic ventral mesh rectopexy (LVMR) is a keyhole surgery procedure which is used to treat patients suffering from rectal prolapse (when the bowel slides out of the anus or slides in on itself). This procedure is designed to straighten and reattach the rectum (part of the bowel nearest to the anus) back into its normal position. The aim of this study is to find out whether LVMR could be an effective treatment for patients with chronic constipation who are unresponsive to standard treatment options.

### Who can participate?

Adults with chronic constipation who are unresponsive to non-surgical treatment.

### What does the study involve?

Eligible participants are randomly allocated into three groups, with different waiting times before their LVMR surgery. Those in the first group receive the surgery when they enter the study, those in the second group receive the surgery 12 weeks after entering the study, and those in the third group receive the surgery 24 weeks after entering the study. The effectiveness of the surgery on improving constipation symptoms and quality of life is measured using a number of questionnaires every 12 weeks (until 48 weeks), with the option of long term follow up to 72 weeks.

### What are the possible benefits and risks of participating?

Participants may benefit from an improved quality of life as a result of taking part and will be carefully selected and monitored during the course of the study. The surgeons performing the surgery will also be carefully selected and monitored to ensure a high level of care. The risks of trial participation are very low over and above standard surgical risks and risks of anaesthesia. Radiological investigations (scans) routinely performed to select appropriate patients for surgery require a small ionising radiation dose, although this is not considered to be harmful. A number of questionnaires contain personal questions about bowel problems and the effect of these on quality of life and behaviour, which may be difficult for some participants.

Where is the study run from?

The Royal London & Whipps Cross Hospitals and nine other hospitals in England (UK)

When is the study starting and how long is it expected to run for?

July 2015 to March 2020

Who is funding the study?

National Institute for Health Research (UK)

Who is the main contact?

Ms Natasha Stevens

## Contact information

### Type(s)

Scientific

### Contact name

Mrs Shiva Taheri

### Contact details

Barts and The London School of Medicine and Dentistry

Centre for Neuroscience Trauma

Blizard Institute

2 Newark Street

London

United Kingdom

E1 2AT

## Additional identifiers

### Protocol serial number

19016

## Study information

### Scientific Title

Stepped wedge randomised trial of laparoscopic ventral mesh rectopexy in adults with chronic constipation

### Acronym

CapaCiTY03

### Study objectives

This trial aims to address the evidence gap for laparoscopic ventral mesh rectopexy (LVMR) and will assess the efficacy of LVMR for the treatment of adults with chronic constipation.

### Ethics approval required

Old ethics approval format

**Ethics approval(s)**

First Medical Research Ethics Committee, 06/07/2015, ref: 15-LO-0609

**Study design**

Randomised; Interventional and Observational; Design type: Treatment, Cohort study

**Primary study design**

Interventional

**Study type(s)**

Treatment

**Health condition(s) or problem(s) studied**

Chronic constipation

**Interventions**

Patients are randomly allocated into three groups with different waiting times for the Laparoscopic Ventral Mesh Rectopexy (LapVMR) surgery. All groups have a run in period of 4 weeks to arrange surgery, with an average waiting time of 3 months.

Group 1: The LVMR surgery is performed at T = 0 weeks

Group 2: The LVMR surgery is performed at T = 12 weeks

Group 3: The LVMR surgery is performed at T = 24 weeks

Within the intervention, participants are admitted to hospital for at least one day post-operative care. The stepped wedged design allows data before surgery to be compared to data after surgery, effectively, patients act as their own controls.

Post-operatively, patients are followed up after 1 day, after 30 days and once every 12 weeks to maximum 72 weeks.

**Intervention Type**

Other

**Primary outcome(s)**

Patient assessment of quality of life is assessed using the Patient Assessment of Constipation Quality of Life (PAC-QOL) questionnaire at 24 weeks post-surgery compared to pre-surgery controls.

**Key secondary outcome(s)**

1. Quality of Life is measured using at 24, 48 and 72 weeks using the following questionnaires:
  - 1.1. Patient Assessment of Constipation Symptom (PAC-SYM) total and individual domain scores
  - 1.2. Patient Assessment of Constipation Quality of Life (PAC-QOL) total and individual domain score
  - 1.3. Patient Health Questionnaire-9 (PHQ-9)
  - 1.4. Generalized anxiety disorder questionnaire (GAD7)
  - 1.5. Global patient satisfaction/improvement score (VAS) and whether they would recommend each treatment experienced to other patients
  - 1.6. Chronic Constipation Brief Response Questionnaire (CC-BRQ), and brief illness perception questionnaire (BIPQ)
  - 1.7. St Marks Incontinence score (for concurrent symptoms)
  - 1.8. Baseline brief sexual function questionnaire (PISQ-12 for women and MSHQ-EjD Short Form

for men)

- 1.9. A two week patient bowel diary (for 2 weeks prior to each assessment) to record bowel frequency and whether each evacuation was 'spontaneous (no use of laxatives) and/or complete
2. Health Economics measured at 24, 48 and 72 weeks using a patient health journal and generic quality of life using the EQ-5D-5L descriptive system and EQ-VAS
3. Specific adverse events, morbidity and mortality data measured at 24, 48 and 72 weeks using:
  - 3.1. Perioperative findings e.g. scarring, adhesions, tissue laxity, pelvic depth, ovarian or uterine pathology
  - 3.2. Procedural data: duration of surgery, blood loss, approach (lap vs. conversion to open), type of mesh and sutures (make, diameter, number);
  - 3.3. Length of post-operative stay
  - 3.4. 30 day re-admission rates;
  - 3.5. Complications (30-day morbidity and mortality)
  - 3.6. Specific outcomes e.g. dyspareunia and sexual function, pelvic pain, urinary dysfunction, new onset-faecal incontinence, early mesh complications (displacement, erosion, infection).
  - 3.7. Early clinical recurrence of structural defect e.g. prolapse or rectocele based on rectal examination +/- adjunctive investigations (as clinically indicated)
4. Qualitative Interviews to assess patient and health professional experiences. Method of measurement include semi structured, digitally recorded, anonymised and transcribed interviews. Interviews will be conducted with a subset of 20 patients and 5 surgeons, before and after surgery (timing will vary depending on group allocation)

#### **Completion date**

01/03/2020

## **Eligibility**

#### **Key inclusion criteria**

1. Aged between 18 and 70 years
2. Patient selfreports problematic constipation
3. Symptom onset > 6 months prior to recruitment
4. Symptoms meet American College of Gastroenterology definition of constipation
5. Constipation failed treatment to a minimum basic standard (NHS Map of Medicine 2012 (lifestyle AND dietary measures AND =2 laxatives or prokinetics) tried (no time requirement)
6. Ability to understand written and spoken English (due to questionnaire validity)
7. Ability and willingness to give informed consent
8. Failure of nonsurgical interventions (minimum of nurse led behavioural therapy)
9. Internal rectal prolapse as determined by clinical examination and INVEST

#### **Participant type(s)**

Patient

#### **Healthy volunteers allowed**

No

#### **Age group**

Adult

#### **Lower age limit**

18 years

**Sex**

All

**Total final enrolment**

28

**Key exclusion criteria**

1. Significant organic colonic disease (red flag symptoms e.g. rectal bleeding prior investigated); IBD; megacolon or megarectum (if diagnosed beforehand); severe diverticulosis/stricture/birth defects deemed to contribute to symptoms (incidental diverticulosis if known not an exclusion)
2. Major colorectal resectional surgery
3. Current overt pelvic organ prolapse (bladder, uterus) or disease requiring obvious surgical intervention other than lap VMR
4. Previous rectopexy
5. Sacral nerve stimulation (SNS) in situ
6. Rectal impaction (as defined by digital and abdominal examination: these form part of the NHS Map of Medicine basic standard)
7. Significant neurological disease deemed to be causative e.g. Parkinson's, spinal injury, multiple sclerosis, diabetic neuropathy (not uncomplicated diabetes alone)
8. Significant connective tissue disease: scleroderma, systemic sclerosis and SLE (not hypermobility alone)
9. Significant medical comorbidities and activity of daily living impairment (based on Bartell index in apparently frail patients Barthel index  $\leq 11$ )
10. Major active psychiatric diagnosis (e.g. schizophrenia, major depressive illness and mania)
11. Chronic regular opioid use (at least once daily use) where this is deemed to be the cause of constipation based on temporal association of symptoms with onset of therapy
12. Pregnancy or intention to become pregnant during study period
13. Known severe intraabdominal adhesions

**Date of first enrolment**

01/07/2015

**Date of final enrolment**

30/06/2019

**Locations****Countries of recruitment**

United Kingdom

England

**Study participating centre**

**The Royal London & Whipps Cross Hospitals**

Barts Health NHS Trust

Whitechapel Road

London

United Kingdom

E1 1BB

**Study participating centre**

**St Thomas' Hospital**

Westminster Bridge Road  
Guy's and Thomas' NHS Foundation Trust  
London  
United Kingdom  
SE1 7EH

**Study participating centre**

**Wythenshawe Hospital**

University Hospital of South Manchester NHS Foundation Trust  
Southmoor Road  
Wythenshawe  
Manchester  
United Kingdom  
M23 9LT

**Study participating centre**

**Northern General Hospital**

Sheffield Teaching Hospital NHS Foundation Trust  
Herries Road  
Sheffield  
United Kingdom  
S5 7AU

**Study participating centre**

**Poole Hospital**

Poole Hospital NHS Foundation Trust  
Longfleet Road  
Poole  
United Kingdom  
BH15 2JB

**Study participating centre**

**Southmead Hospital**

North Bristol NHS Foundation Trust  
Southmead Road  
Westbury-on-Trym

Bristol  
United Kingdom  
BS10 5NB

**Study participating centre**  
**Royal Shrewsbury Hospital**  
The Shrewsbury and Telford Hospital NHS Trust  
Mytton Oak Road  
Shrewsbury  
United Kingdom  
SY3 8XQ

**Study participating centre**  
**Broomfield Hospital**  
Mid Essex Hospital Services NHS Trust  
Court Road  
Chelmsford  
United Kingdom  
CM1 7ET

**Study participating centre**  
**St. James's University Hospital**  
The Leeds Teaching Hospitals NHS Trust  
Leeds  
United Kingdom  
LS9 7TF

**Study participating centre**  
**Southend University Hospital NHS Foundation Trust**  
Prittlewell Chase  
Southend-on-Sea  
Westcliff-On-sea  
United Kingdom  
SS0 0RY

## **Sponsor information**

**Organisation**  
Queen Mary University of London

ROR

<https://ror.org/026zzn846>

## Funder(s)

### Funder type

Government

### Funder Name

National Institute for Health Research

### Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

### Funding Body Type

Government organisation

### Funding Body Subtype

National government

### Location

United Kingdom

## Results and Publications

### Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be stored in a repository held by the Barts and the London Pragmatic Clinical Trials Unit, Queen Mary University of London.

### IPD sharing plan summary

Stored in repository

### Study outputs

| Output type                                   | Details                       | Date created | Date added | Peer reviewed? | Patient-facing? |
|---|-------------------------------|--------------|------------|----------------|-----------------|
| <a href="#">Results article</a>               |                               | 19/05/2022   | 23/05/2022 | Yes            | No              |
| <a href="#">Results article</a>               |                               | 01/11/2021   | 27/10/2022 | Yes            | No              |
| <a href="#">Protocol article</a>              | protocol                      | 05/02/2018   |            | Yes            | No              |
| <a href="#">Participant information sheet</a> | version V3                    |              | 03/05/2016 | No             | Yes             |
| <a href="#">Participant information sheet</a> | Participant information sheet | 11/11/2025   | 11/11/2025 | No             | Yes             |
| <a href="#">Study website</a>                 | Study website                 | 11/11/2025   | 11/11/2025 | No             | Yes             |



