

What is the diagnostic accuracy of midwife-delivered point-of-care ultrasound for detecting breech presentation at term?

Submission date 03/09/2024	Recruitment status Recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 30/04/2025	Overall study status Ongoing	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 10/02/2026	Condition category Pregnancy and Childbirth	<input type="checkbox"/> Individual participant data
		<input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Most babies are cephalic (head down) at the end of pregnancy, however, 3-4% of babies are breech (bottom down). At a late pregnancy check, midwives feel a woman's tummy (abdomen) using their hands (palpation) to check for the baby's position but up to 40% of breech babies are missed. Being specific about the baby's position is particularly important so that plans for safe delivery can be made. Breech babies are at higher risk of injury during normal birth than cephalic babies. Undiagnosed breech births can be associated with poor outcomes for the baby and mother, so determining which way the baby is positioned is important to provide women with the information they need in order to make an informed choice about their care.

The 'gold standard' for determining a baby's position is by ultrasound scan. This is performed by a trained person called a sonographer or a specialist doctor using a hospital-based ultrasound machine. In recent years, smaller handheld ultrasound machines have been introduced into some clinical areas to help diagnose conditions where a conventional ultrasound is not present. An advantage of these machines is that they are portable. These machines connect to a mobile phone or e-tablet.

This study aims to find out whether midwives are able to tell what position the baby is in before it is born using a small handheld ultrasound machine and to see if this is in agreement with a conventional scan used in hospitals. The researchers also want to know what maternity service users think about the use of these devices (explored through recorded interviews with selected women and midwives), and whether these devices will reduce the risk of undiagnosed breech presentation and its potential complications, and in turn whether this will save money for the NHS. The team running the study include experts in trial design and statistics, obstetricians, midwives, sonographers, neonatologists, qualitative researchers and health economists from a number of institutions with direct patient involvement.

Who can participate?

Pregnant women who are between 35+0-36+6 weeks pregnant with one baby, of different backgrounds from collaborating maternity units which are geographically diverse and include the North of England, the Midlands, South Coast, London and the East of England

What does the study involve?

Participants will be invited to participate in the study by a midwife at their 36-week appointment. The researchers will compare handheld ultrasound to the 'gold standard' conventional ultrasound. The conventional ultrasound will need to be performed within one day of the handheld ultrasound.

What are the possible benefits and risks of participating?

Ultrasound, both using conventional machines and handheld devices, is very safe and will not cause harm to the pregnant woman or their baby. Depending on when and where the midwife appointment and confirmation scan take place, this may involve an additional visit to the hospital. There is a small chance that the baby could change position after the scan. If this happens, options will be discussed with the study participant as per the hospital's guidelines.

Where is the study run from?

1. Imperial College London (UK)
2. The Centre for Trials Research, Cardiff University (UK)
3. City St George's (UK)
4. University College London (UK)

When is the study starting and how long is it expected to run for?

May 2023 to March 2027

Who is funding the study?

NIHR Health Technology Assessment (HTA) (UK)

Who is the main contact?

Dr Eleri Owen-Jones, sono-breech@cardiff.ac.uk

Contact information

Type(s)

Scientific, Principal investigator

Contact name

Prof Christoph Lees

Contact details

Centre for Fetal Care
Queen Charlotte's & Chelsea Hospital
Du Cane Road
London
United Kingdom
W12 0HS
+44 (0)207 594 2104
christoph.lees@nhs.net

Type(s)

Public

Contact name

Dr Eleri Owen-Jones

ORCID ID

<https://orcid.org/0000-0003-0850-4724>

Contact details

Centre for Trials Research
Cardiff University
Heath Park
Cardiff
United Kingdom
CF14 4YS
+44 (0)29 2068 7620
owen-jonesce@cardiff.ac.uk

Type(s)

Scientific, Principal investigator

Contact name

Dr Amarnath Bhide

Contact details

Ground Floor
Lanesborough Wing
St George's Hospital
Blackshaw Road
Tooting
London
United Kingdom
SW17 0QT
+44 (0)2087250007
amar.bhide@nhs.net

Additional identifiers**Clinical Trials Information System (CTIS)**

Nil known

Integrated Research Application System (IRAS)

318520

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

CPMS 53525, IRAS 318520

Study information

Scientific Title

Diagnostic accuracy of handheld ultrasound at 36 weeks of gestation to determine fetal presentation

Acronym

Sono-breech

Study objectives

What is the diagnostic accuracy of midwife-delivered point of care ultrasound (PoCUS) for detecting breech presentation at term?

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 03/07/2024, West Midlands - South Birmingham Research Ethics Committee (2 Redman Place, Stratford, London, E20 1JQ, United Kingdom; +44 (0)2071048121; southbirmingham.rec@hra.nhs.uk), ref: 24/WM/0143

Study design

Multicentre prospective observational study

Primary study design

Observational

Study type(s)

Other

Health condition(s) or problem(s) studied

Pregnant women

Interventions

There are three components to the study: the main diagnostic accuracy study; a nested acceptability study using qualitative methodology; and a health economic evaluation to evaluate cost-effectiveness.

36-week appointment:

At routine 36-week antenatal appointments, eligible pregnant women will be consented to take part in the Sono-breech study. They will receive their usual 36-week examination including abdominal palpation to determine fetal presentation and check fetal viability. Following this, the Sono-breech trained midwife will perform a handheld PoCUS scan to check presentation and fetal viability. This should take no longer than 30 minutes.

Following the handheld scan the Sono-breech midwife will arrange for the woman to attend an ultrasound scan performed by an Ultrasound Practitioner on a conventional ultrasound machine within one day. The Ultrasound Practitioner will check fetal presentation, and this will be recorded. This should take no longer than 10 minutes. The conventional ultrasound confirmation must NOT be done on a handheld PoCUS device. The pathway for ultrasound scans will differ from site to site, and liaison with the maternity unit and ultrasound department during study set-up will ensure availability for same/next day ultrasound assessment using a static conventional ultrasound machine.

Questionnaire:

Around 6 weeks after the birth, two online questionnaires will be sent to the study participants via email. The questionnaires will ask some questions about the pregnancy and the baby.

Interview:

Between 6-12 weeks after the birth, some study participants will be invited to take place in a short interview, where a member of the research team will ask questions about how the study participant felt about the study and the use of the handheld ultrasound devices. The researchers plan to interview some women who choose not to take part in the study to explore why they preferred not to have the scan.

Intervention Type

Other

Primary outcome(s)

Diagnostic accuracy (sensitivity and specificity) of midwife-conducted handheld PoCUS at 36 weeks for the detection of breech presentation compared in the same women to 'gold standard' conventional ultrasound

Key secondary outcome(s)

1. Acceptability to midwives and pregnant women of handheld PoCUS in the detection of breech presentation at term, measured using focus groups (midwives) during the study period; and interviews (pregnant women) up to 12 weeks after birth
2. Resource use including mode of birth and neonatal intensive care unit (NICU) – reasons and outcome of admissions, and length of stay obtained from patient records for up to 28 days
3. Evaluation of training requirements for midwives using handheld PoCUS for fetal presentation, measured using focus groups during the study period
4. Proportion of breech presentations that remain undiagnosed in labour and resulting change in management, measured using a Pregnancy Outcome CRF recorded up to 12 weeks after birth
5. Birth experience measured using the Birth Experience Assessment Measure completed up to 6 weeks after birth
6. Infant quality of life measured using the Infant Quality of Life Instrument (IQI) completed up to 6 weeks after birth

Completion date

31/03/2027

Eligibility

Key inclusion criteria

1. Singleton live pregnancy
2. 35+0 - 36+6 weeks of gestational age
3. Ability to give valid informed consent
4. Commitment to attend a second scan within 1 day
5. Midwifery sample eligibility – being part of the Sono-breech study and undertaking PoCUS scanning as part of this study

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Female

Total final enrolment

0

Key exclusion criteria

1. Multiple pregnancy
2. Unable to attend a second scan within 1 day

Date of first enrolment

05/08/2024

Date of final enrolment

30/11/2026

Locations**Countries of recruitment**

United Kingdom

England

Study participating centre**Queen Charlotte's and Chelsea Hospital**

Imperial College Healthcare NHS Trust

London

England

W12 0HS

Study participating centre**St Mary's Hospital**

Imperial College Healthcare NHS Trust

London

England

W2 1NY

Study participating centre

Ipswich Hospital

Heath Road
Ipswich
England
IP4 5PD

Study participating centre

West Suffolk Hospital

West Suffolk NHS Foundation Trust
Lowestoft
England
IP33 2QZ

Study participating centre

East Surrey Hospital

Canada Avenue
Redhill
England
RH1 5RH

Study participating centre

Northwick Park Hospital

Watford Road
Harrow
England
HA1 3UJ

Study participating centre

James Cook University Hospital

Marton Road
Middlesbrough
England
TS4 3BW

Study participating centre

United Lincolnshire Teaching Hospitals NHS Trust

Lincoln County Hospital
Greetwell Road
Lincoln
England
LN2 5QY

Study participating centre

St Georges Hospital

Blackshaw Road
Tooting
London
England
SW17 0QT

Study participating centre

City Hospital NHS Trust

City Hospital N H S Trust
Dudley Road
Birmingham
England
B18 7QH

Study participating centre

Birmingham Women's and Children's NHS Foundation Trust

Steelhouse Lane
Birmingham
England
B4 6NH

Study participating centre

Manchester University NHS Foundation Trust

Cobbett House
Oxford Road
Manchester
England
M13 9WL

Study participating centre

University Hospitals of Leicester NHS Trust

Leicester Royal Infirmary
Infirmary Square
Leicester
England
LE1 5WW

Sponsor information

Organisation

Imperial College London

ROR

<https://ror.org/041kmwe10>

Funder(s)

Funder type

Government

Funder Name

Health Technology Assessment Programme

Alternative Name(s)

NIHR Health Technology Assessment Programme, Health Technology Assessment (HTA), HTA

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Study website	Study website	11/11/2025	11/11/2025	No	Yes