Comparing the cost-effectiveness of two commonly used types of dye for determining whether cancer has spread into the lymphatic system

| Submission date 13/04/2022 | Recruitment status No longer recruiting | [X] Prospectively registered[X] Protocol |
|------------------------------|---|---|
| Registration date 08/12/2022 | Overall study status Completed | Statistical analysis planResults |
| Last Edited 14/12/2023 | Condition category Cancer | Individual participant dataRecord updated in last year |

Plain English summary of protocol

Background and study aims

Women with a small breast cancer require surgery to remove the cancer from their breast and also to remove a few lymph nodes from their armpit (sentinel node biopsy) at the same time to see whether the cancer has spread. These women have an injection of a dye before their surgery which demonstrates the lymph nodes the surgeon needs to remove. This study aims to compare the cost-effectiveness of two commonly used types of dye for sentinel node biopsy - Magtrace and technetium.

Who can participate? Women aged 18 years or above with breast cancer

What does the study involve?

Participants will be randomly allocated to have either Magtrace or technetium. A research assistant will shadow these women in their visits to the hospital before surgery and work out the hospital costs associated with each treatment.

What are the possible benefits and risks of participating?

The potential benefits of using Magtrace® will be in simplifying the patient pathway before surgery. Magtrace® will be given on the day of surgery, whereas technetium would be given on the day of surgery or the day before. This means around the time of surgery there will be less stress for the patient, they will be ready to have their surgery at any time of the day (rather than having to wait until they have a technetium injection). Technetium requires aliquoting out from the Christie and transporting across to the nuclear medicine department on a daily basis, and following radioactive licences, must be given by two trained professionals in a very controlled manner. Magtrace® can be given by any trained member of staff without the need for radioactivity licences. It will negate the need for the complex process of transferring the nuclear medicine across the city and the need for staff to make up the injections every morning and give them to patients. Magtrace® will therefore simplify the pathway and remove multiple steps

freeing up staff for other jobs. Magtrace® will make theatre scheduling in the morning easier as any patient will be available for theatre, potentially making less waiting for theatre staff and more efficient operating. It is unlikely that extra cases will be done as a result of the change but it may avoid overruns and delays in theatre.

All processes and technologies have previously been proven to be effective. Technetium sentinel node injection is often combined with a blue dye. The blue dye has a small risk of anaphylaxis (a severe, potentially life-threatening allergic reaction) of about 1 in 10,000 and can stain the skin of the breast blue for several months. Magtrace® does not have a risk of anaphylaxis but can stain the skin of the breast brown for several months post-operatively.

Where is the study run from? Manchester University NHS Foundation Trust (UK)

When is the study starting and how long is it expected to run for? October 2020 to May 2025

Who is funding the study? Endomag (UK)

Who is the main contact? Mr James Harvey, james.harvey@mft.nhs.uk

Contact information

Type(s) Principal Investigator

Contact name Mr James Harvey

ORCID ID http://orcid.org/0000-0002-1359-2186

Contact details Nightingale Centre Manchester University NHS Foundation Trust Southmoor Road Manchester United Kingdom SK23 7EJ +44 (0)1612914436 james.harvey@mft.nhs.uk

Additional identifiers

EudraCT/CTIS number Nil known

IRAS number 293053 **ClinicalTrials.gov number** Nil known

Secondary identifying numbers

IRAS 293053, CPMS 53234

Study information

Scientific Title

Magtech: cost-effectiveness and benefits of Magtrace® versus Technetium in sentinel node biopsy for breast cancer

Acronym

Magtech

Study objectives

Single-site case-control study investigating the healthcare costs of Magtrace®, and a discretechoice qualitative experiment of the perceived value of Magtrace® vs the current standard of care

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 23/09/2022, North West - Greater Manchester East Research Ethics Committee (3rd Floor, Barlow House, 4 Minshull Street, Manchester, M1 3DZ, UK; +44 (0)2071048009; gmeast. rec@hra.nhs.uk), ref: 22/NW/0192

Study design

Single-site case-control study and a discrete-choice qualitative experiment

Primary study design

Interventional

Secondary study design Randomised controlled trial

Study setting(s) Hospital

Study type(s) Treatment

Participant information sheet

Not available in web format, please use the contact details to request a participant information sheet

Health condition(s) or problem(s) studied

Early breast cancer

Interventions

Using a stopwatch, research staff have the ability to consent and then actively monitor (shadow around the hospital) a group of ~20 patients in each treatment group (Magtrace® vs routine care using Technetium nuclear medicine). It is critical prior to recruitment that key covariates and factors likely to influence not only outcomes but also health economic factors of care are identified. These factors will require consideration before randomisation to Magtrace or routine care, with randomisation recommended to take place using sealed envelopes in blocks of ten. If imbalances in key covariates occur following randomisation, adjustment of health economic outcomes will be performed to mitigate these imbalances using gold-standard two-stage bootstrapping for cost-effectiveness analyses.

Intervention Type

Device

Phase Phase III/IV

Drug/device/biological/vaccine name(s)

Magtrace®, technetium

Primary outcome measure

Cost-effectiveness of Magtrace® compared to Technetium in the identification of sentinel nodes in breast cancer, measured using data records over duration of the study.

Secondary outcome measures

 Patient time spent in hospital and in transit in total over the preoperative and perioperative journey, measured using electronic patient records over the duration of the patient visit.
 Patient anxiety on the day of surgery measured using the State-Trait Anxiety Inventory (STAI-Y) given on the ward on the morning of surgery

3. Number and length of patient hospital visits comparing preoperative visits (clinic visits, preoperative assessment, tracer injection visits, ward visits and operative visits), measured on the day of surgery

4. Safety of Magtrace® and efficacy: adverse events relatable to tracers, number of nodes removed, number of nodes containing tracer, % of patients with detectable tracer in the axillary sentinel nodes, can surgeons differentiate Magtrace® and Magseed signal®, recorded for as long as the participant is on the trial

5. Cost per episode and for total care, measured using activity-based costing (TDABC). TDABC is a bottom-up approach to healthcare pathway mapping and costing, which records pathways observed during routine clinical practice, identifies all points and durations of interaction with healthcare providers therein; and assigns time-dependent costs to each constituent.

6. Surgical start time, days of week of operating, and delays to getting patients to the theatre. Delays will be recorded by asking the theatre team directly pre-operatively whether they had to wait for the patient to arrive and what the reason was, and also whether the list had to be rescheduled/moved to accommodate a delay, measured using electronic patient records over the duration of the patient visit.

7. Current pathways of patient care: the research assistants provide a 'learned' assessment of what they believe represents a 'typical' pathway at the Nightingale centre. A multidisciplinary team will then assess the validity of this pathway and voice any concerns/objections/potential additions. The duration of each patient visit is monitored throughout the trial.

8. Key aspects of care pathways important to patients and healthcare professionals in the management of breast cancer; in order to ensure that all relevant points of interaction with

healthcare staff, and utilisation of health services, are captured, the pathway will be amended as necessary. Duration of the patient visit.

9. Preferences/overall satisfaction with the care package for Magtrace vs Technetium from the perspective of patients and healthcare providers. Once data is collected for 20 patients undergoing Magtrace, and 20 patients undergoing routine care, it will be possible to multiply NHS resource utilisation by representative NHS unit costs (staff salaries per minute, radiography, inpatient, nuclear medicine) and determine an array of health economic outcomes. The result will be a health-economic comparison of Magtrace® over Technetium nuclear medicine. Sensitivity analyses will be performed on this dataset to measure the robustness of the findings to changes in the values of the parameters measured during the TDABC stopwatch exercise. Rather than relying on point estimates, this will enable the derivation of credible intervals, and determine how specific factors such as time in the hospital, number of visits, requirements for isotopes etc.

Overall study start date

07/10/2020

Completion date

01/05/2025

Eligibility

Key inclusion criteria

1. Participant is willing and able to give informed consent for participation in the study

2. Female aged 18 years or above

3. Diagnosed with breast cancer (invasive) requiring Magseed® localisation and sentinel node biopsy

4. Willing to allow her General Practitioner and consultant, if appropriate, to be notified of participation in the study

5. Undergoing breast-conserving surgery with sentinel node biopsy

6. Surgeons may only operate on the Magtrace arm of the study if they have completed a minimum of five training cases with Magtrace

Participant type(s)

Patient

Age group Adult

Lower age limit 18 Years

Sex Female

Target number of participants 50

Key exclusion criteria

- 1. Patients with a pacemaker or implanted device in the chest wall
- 2. Patients who are pregnant or lactating
- 3. Patients who have received Magtrace® (iron oxide) injection in the previous 6 months
- 4. Patients with previous ipsilateral axillary surgery
- 5. Patients whose breast and axillary surgery are not due to be performed synchronously
- 6. Patients following neoadjuvant chemotherapy

7. Patients who require MRI follow-up of the ipsilateral breast in the year following surgery (as Magtrace® may interfere with MRI)

8. Patients requiring an interpreter

9. Patients involved in current research or have recently been involved in any research prior to recruitment

Date of first enrolment 01/05/2024

Date of final enrolment

01/05/2025

Locations

Countries of recruitment England

United Kingdom

Study participating centre

Wythenshawe Hospital

Southmoor Road Wythenshawe Manchester United Kingdom M23 9LT

Sponsor information

Organisation

Manchester University NHS Foundation Trust

Sponsor details

Southmoor Road Manchester England United Kingdom M23 9LT +44 (0)1612914436 charlotte.bryant@mft.nhs.uk

Sponsor type Hospital/treatment centre

Website https://mft.nhs.uk/

ROR https://ror.org/00he80998

Funder(s)

Funder type Industry

Funder Name Endomag

Results and Publications

Publication and dissemination plan

Planned publication in a peer-reviewed journal

Intention to publish date

31/12/2025

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are not expected to be made available

IPD sharing plan summary

Not expected to be made available

Study outputs

| Output type | Details version 1 | Date created | Date added | Peer reviewed? | Patient-facing? |
|----------------------|-----------------------------|--------------|------------|----------------|-----------------|
| <u>Protocol file</u> | | 03/08/2022 | 24/10/2022 | No | No |
| HRA research summary | | | 26/07/2023 | No | No |