Does antithrombotic prophylaxis increase the incidence of post-operative bleeding and duration of post-operative hospital stay after thyroidectomy? A retrospective study

Submission date	Recruitment status No longer recruiting	Prospectively registered		
23/01/2018		Protocol		
Registration date 05/02/2018	Overall study status Completed	Statistical analysis plan		
		[X] Results		
Last Edited	Condition category	[] Individual participant data		
13/05/2019	Suraerv			

Plain English summary of protocol

Background and study aims

Providing antithrombotic medication (drugs that reduce the change of blood clots) during surgery could help prevent blood clots and pulmonary embolism (blocking of a bloog vessel in the lung), however this increases the risk of bleeding after surgery. This topic has not been adequately investigated in thyroid surgery. In thyroid surgery, even a small bleeding may result in airway compression and death by asphyxia (a lack of oxygen). The aim of this study is to establish whether providing antithrombotic drugs during surgery can negatively affect the early outcome of patients undergoing total thyroidectomy, with particular reference to the incidence of bleeding and duration of post-operative hospital stay. Therefore this study aims to clarify if antithrombotic drugs administered in order to prevent thromboembolic events are safe and effective in thyroid surgery.

Who can participate?

Adults aged 18 and older who are having a thyroid surgery.

What does the study involve?

Participants underwent thyroidectomy and antithrombotic drugs according with standard surgical criteria and international guidelines for antithrombotic prophylaxis. All patients are subject to a close monitoring in the first 12 hours after surgery and, after discharge, are reevaluated within 1 week in order to assess the wound and, if appropriate, remove suture. The data from the surgeries are reviewed to assess the impact of the antithrombotic drugs.

What are the possible benefits and risks of participating? There are no direct benefits or risks with participating.

Where is the study run from? Policlinico Universitario "Duilio Casula" (Italy)

When is the study starting and how long is it expected to run for? November 2017 to January 2018

Who is funding the study? University of Cagliari (Italy)

Who is the main contact? Dr Enrico Erdas (Scientific)

Contact information

Type(s)

Scientific

Contact name

Dr Enrico Erdas

Contact details

Policlinico Universitario "Duilio Casula" SS554 Cagliari Italy 09124

Additional identifiers

Protocol serial number

11/1968

Study information

Scientific Title

Does antithrombotic prophylaxis worsen early outcome of total thyroidectomy? A retrospective cohort study

Study objectives

It is expected that the incidence of post-operative bleeding and the length of hospital stay after total thyroidectomy would be higher when antithrombotic prophylaxis was administered.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Institutional Ethical Committee University Hospital of Monserrato, University of Cagliari, 20/11/2017 ref: PROT. PG/2017/8430

Study design

Retrospective cohort study

Primary study design

Observational

Study type(s)

Treatment

Health condition(s) or problem(s) studied

factors influencing outcome of thyroid surgery

Interventions

Participants have antithrombotic prophylaxis with low-molecular-weight heparin (a subcutaneous prophylactic dose (i.e., 0.3 ml) of nadroparin calcium administered the day before their operation and continued for at least 10 days. Participants undergo their thyroid surgery.

After the operation all patients are subject to a close monitoring in the first 12 hours after surgery in order to detect early signs of cervical haematoma, respiratory distress or hypocalcaemia. Drains are removed when the daily amount of fluid collection falls below 20 ml in each reservoir. Parathyroid hormone (PTH) and serum calcium levels are measured on the first and second postoperative day, while further controls are performed only if required (PTH at very low level, clinical signs of hypocalcaemia regardless of serum calcium levels). Post-operative oral calcium and vitamin D supplements are administered in all symptomatic patients with hypocalcemia. After discharge, all patients are re-evaluated within 1 week in order to assess the wound and, if appropriate, remove intradermal suture. All patients underwent telephone follow-up once a week for at least one month.

Researchers review the data about surgical outcomes to investigate the usage of the antithrombotic medication.

Intervention Type

Procedure/Surgery

Primary outcome(s)

- 1. Incidence of severe postoperative cervical hematoma is measured using the the need for surgical revision
- 2. Length of postoperative hospital stay is measured using patient records at discharge

Key secondary outcome(s))

Incidence of thromboembolic events is measured using the diagnosis of deep vein thrombosis and/or pulmonary embolism.

Completion date

15/01/2018

Eligibility

Key inclusion criteria

All patients submitted to total thyroidectomy for whom antithrombotic prophylaxis would be indicated.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Total final enrolment

526

Key exclusion criteria

- 1. Aged below 18 years
- 2. Anticoagulant or antiplatelet therapy
- 3. Recurrent goitre and operations other than total thyroidectomy (hemithyroidectomy, thyroidectomy with central or lateral neck dissection).

Date of first enrolment

01/02/2013

Date of final enrolment

31/10/2017

Locations

Countries of recruitment

Italy

Study participating centre
Policlinico Universitario "Duilio Casula"
Italy
09042

Sponsor information

Organisation

University of Cagliari

ROR

https://ror.org/003109y17

Funder(s)

Funder type

University/education

Funder Name

University of Cagliari

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Enrico Erdas at erdasenrico@libero.it.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	24/04/2019	13/05/2019	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes