The beneficial effect on sleep of a food supplement based on Scutellaria lateriflora L. (aerial parts of the plant)

Submission date	Recruitment status	[X] Prospectively registered
22/05/2024	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
03/06/2024	Completed	Results
Last Edited	Condition category	Individual participant data
14/06/2024	Mental and Behavioural Disorders	Record updated in last year

Plain English summary of protocol

Background and study aims

Insomnia is the most common sleep disorder worldwide, where people have trouble falling or staying asleep. This affects their overall well-being, and can lead to mental, cognitive, and physical health issues.

About one-third of adults experience insomnia, with 10%-15% reporting daytime problems, and 6%-10% being diagnosed with insomnia disorder. Women are 1.44 times more likely to suffer from insomnia than men.

Insomnia significantly increases the risk of heart diseases like high blood pressure, heart attacks, and chronic heart failure. Research also links insomnia to a higher risk of type 2 diabetes. Additionally, sleeping less than 6 hours a night is linked to obesity, type 2 diabetes, high blood pressure, and heart diseases. Insomnia is also connected to major depression, often preceding and predicting its onset.

A U.S. survey found that 4.5% of adults prefer natural remedies over medications for mild insomnia. This study aims to show that a dietary supplement can improve sleep in people with primary insomnia who don't have severe symptoms.

The plant genus Scutellaria, which includes over 350 species, has been traditionally used for treating various disorders like nerve issues, chronic diseases, liver problems, and anxiety. Scutellaria lateriflora, or skullcap, is the focus of this study. Traditionally used by Native Americans to improve blood flow, reduce anxiety, and treat mental health issues, skullcap is now used in North America to treat epilepsy, nerve damage from infections, insomnia, anxiety, nerve pain, and withdrawal from tranquilizers.

Scutellaria lateriflora contains many active compounds, including flavones like baicalein, baicalin, chrysin, and wogonin, which are believed to improve mood. Baicalein weakly binds to benzodiazepine receptors, explaining its mild calming effects through interaction with GABA receptors in the nervous system. Baicalein also partially blocks the GABAA receptor, while

wogonin shows anxiety-reducing effects through benzodiazepine receptors. Other compounds interact with serotonin receptors.

In vitro studies (conducted outside of a living organism) have shown that an extract of Scutellaria lateriflora inhibits enzymes that break down GABA, a neurotransmitter that calms the nervous system, contributing to its anxiety-reducing effects.

Another study tested a 100 mg dose of Scutellaria lateriflora extract on rats and found it significantly reduced anxiety and depression behaviors.

The goal of this study is to evaluate how effective a dietary supplement made from Scutellaria lateriflora is at improving sleep in people with primary insomnia.

Who can participate?

Sixty-six subjects aged 18-70 years of either sex, who present mild to moderate primary insomnia and who can understand and sign the informed consent.

What does the study involve?

A single-center, placebo-controlled, randomized, cross-over clinical study will be conducted with a two-week run-in period, in double-blind fashion.

During the two-week run-in period, during which all enrolled subjects will take the placebo, potential compliance with the study will be observed to minimize subject drop-out and thus increase the internal validity of the clinical study. Additionally, at the end of the run-in period, any physiological responses to the placebo, as well as false adverse effects (although very unlikely - see ADVERSE EVENTS section), will be identified, the eligibility of the subjects and therefore the sample population will be confirmed, and initial data regarding the quality and severity of insomnia in the recruited subjects can be obtained.

What are the possible benefits and risks of participating? No risks are foreseen. An improvement in the sleep function of the subjects treated with the food supplement is hypothesized. However, no benefit may be achieved.

Where is the study run from? Epo S.r.l. (Italy)

When is the study starting and how long is it expected to run for? July 2023 to November 2024

Who is funding the study? Epo S.r.l. (Italy)

Who is the main contact?

- 1. Prof. Maria Daglia (scientific) maria.daglia@unina.it
- 2. Dr. Alessandra Baldi (public) alessandra.baldi.alimenti@gmail.com

Contact information

Type(s)

Scientific

Contact name

Prof Maria Daglia

ORCID ID

http://orcid.org/0000-0002-4870-7713

Contact details

Via Domenico Montesano, 49 Naples Italy 800016 + 39 3398177623 maria.daglia@unina.it

Type(s)

Public

Contact name

Ms Alessandra Baldi

ORCID ID

http://orcid.org/0000-0002-2877-9445

Contact details

Viale delle Medaglie d'Oro, 305 Rome Italy 00136 +39 3483854114 alessandra.baldi.alimenti@gmail.com

Type(s)

Principal Investigator

Contact name

Dr Matteo Laringe

Contact details

Viale Maria Bakunin, 41 Naples Italy 80126 + 39 3939406629 matteo.laringe@gmail.com

Additional identifiers

EudraCT/CTIS number

Nil known

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

SCUTESLEEP23_01

Study information

Scientific Title

Efficacy study of a food supplement based on Scutellaria lateriflora L. (aerial parts of the plant) for sleep management: single-center, controlled, randomized, cross-over, double-bind clinical study with a run-in period.

Acronym

SCUTESLEEP23

Study objectives

This study aimed to evaluate the efficacy of the supplementation of the diet with a food supplement based on Scutellaria lateriflora L. (aerial parts of the plant), in the maintenance of the sleep-wake balance, through the improvement of sleep onset and maintenance phase, in subjects where insomnia is independent of other comorbid conditions. Insomnia is considered a significant risk factor for cardiometabolic diseases.

Ethics approval required

Ethics approval required

Ethics approval(s)

Approved 30/01/2024, Ethics Committee of CAMPANIA 1 (Via Mariano Semmola, 53, Naples, 80131, Italy; +39 081/17770131; comitatoetico@istitutotumori.na.it), ref: 2/23

Study design

Interventional monocentric randomized crossover two-arm double-blind placebo-controlled clinical trial

Primary study design

Interventional

Secondary study design

Randomised cross over trial

Study setting(s)

GP practice

Study type(s)

Prevention

Participant information sheet

Not available in web format, please use contact details to request a participant information sheet.

Health condition(s) or problem(s) studied

Subjects with mild to moderate primary insomnia

Interventions

The subjects recruited in the present clinical study will consume, a food supplement based on Scutellaria lateriflora L. (aerial parts of the plant) in capsules at 400 mg/day dose, and a placebo, in a different order, based on the randomization group, for two treatment periods of 56 days, interspersed with a wash-out period of 28 days.

The treatment consists of the BlueCALM® dietary supplement, notified to the Ministry of Health (notification number: 164522).

To maintain the double-blind design, both treatments will be made unrecognizable as the packaging will be identical, and the dosage forms will be the same in color, shape, weight, and taste.

The randomization sequence will be generated by a statistician using STATA 16 software (Stata Statistical Software: Release 16. College Station, TX: StataCorp LLC) and the randomization list will be kept hidden. The participants will be assigned to each of the two treatment groups (food supplement or placebo) casually and by simple randomization (1:1: allocation ratio). The randomization code will consist of a three-digit number as indicated in the respective Case Report Form (CRF).

In the clinical study, 66 participants will be enrolled and divided into two groups (33 for each group):

GROUP 1 subjects who will first take treatment A and, following the wash-out period, treatment B. TREATMENT ORDER: AB

GROUP 2 subjects who will first take treatment B and, following the wash-out period, treatment A. TREATMENT ORDER: BA

Participants will undergo seven visits (start of run-in = tr, at the start of the first treatment period = t0, after 28 days from the start of the first treatment period = t1, at the end of the first treatment period = t2, at the start of the first treatment period = t3, after 28 days from the start of the first treatment period = t4, at the end of the first treatment period = t5) in an outpatient setting. After each clinical visit, all data are filled in the CRF by physicians.

Intervention Type

Supplement

Primary outcome measure

Sleep quality: Pittsburgh Sleep Quality Index (PSQI) [Time frame: tr (run-in start), t0 (baseline), t1 (after 28 days of the first treatment period), t2 (after 56 days of the first treatment period), t3 (baseline of the second treatment period), t4 (after 28 days of the second treatment period), t4 (after 56 days of the second treatment period)]

Secondary outcome measures

Evaluation of the following parameters of the sleep-wake cycle using The Integrated Sleep Diary completed daily for the entire duration of the study, starting from the beginning of the run-in period until the last day of the second treatment period:

- 1. Total Time in Bed (TTL) = the time interval in minutes between when the patient goes to bed and when they get up from bed;
- 2. Sleep Onset Latency (SOL) = the time interval (in minutes) between when the patient goes to bed (turns off the lights or TV) and when they fall asleep;

- 3. Number of Awakenings (NR) = the number of nighttime awakenings;
- 4. Duration of nocturnal awakenings (WASO wakefulness after sleep onset) = intranight wakefulness given by the total sum of wakefulness (in minutes) during nighttime awakenings; 5. Total Sleep Time (TTS) = the time interval (in minutes) between when the patient falls asleep
- and when they wake up definitively in the morning minus intranight wakefulness;
- 6. Sleep Efficiency Index (IES) = TTS/TTL*100.
- 7. Evaluation of the possible presence of pain or discomfort that could affect sleep quality using the Visual Analogue Scale (VAS). [Time frame: tr (run-in start), t0 (baseline), t1 (after 28 days of the first treatment period), t2 (after 56 days of the first treatment period), t3 (baseline of the second treatment period), t4 (after 28 days of the second treatment period), t4 (after 56 days of the second treatment period)].

Overall study start date

01/07/2023

Completion date

30/11/2024

Eligibility

Key inclusion criteria

- 1. Subjects aged 18 70 years of both sexes
- 2. Subjects able to understand and sign the informed consent
- 3. With primary insomnia, present for at least one month
- 4. With ISI questionnaire scores <22
- 5. With GAD-7 questionnaire scores <5
- 6. With PHQ-9 questionnaire scores <5

Participant type(s)

Healthy volunteer, Patient

Age group

Adult

Lower age limit

18 Years

Upper age limit

70 Years

Sex

Both

Target number of participants

66

Total final enrolment

66

Key exclusion criteria

- 1. Aged < 18 and > 70 years
- 2. Pregnant or breastfeeding individuals
- 3. Subjects with secondary insomnia or other sleep disorders, such as obstructive sleep apnea, narcolepsy, nocturnal myoclonus, restless legs syndrome, advanced sleep phase disorder, delayed sleep phase disorder, paradoxical insomnia, parasomnias
- 4. Subjects with ISI questionnaire scores >22
- 5. Subjects with GAD-7 questionnaire scores >5
- 6. Subjects with PHQ-9 questionnaire scores >5
- 7. Subjects suffering from psychiatric disorders: major depression, generalized anxiety disorder, post-traumatic stress disorder, panic attacks, bipolar disorder, dementia, schizophrenia 8. Subjects with chronic and acute pain conditions (e.g., arthritis, fibromyalgia, back pain, headaches, respiratory diseases, asthma, diabetes, heart diseases, injuries, hyperthyroidism, gastroesophageal reflux, epilepsy, Parkinson's disease, Alzheimer's disease, kidney diseases) 9. Subjects who abuse alcohol, amphetamines, caffeine, theine
- 10. Subjects taking antidepressants, diuretics
- 11. Subjects with hypothyroidism
- 12. Subjects who have taken monoamine oxidase inhibitors in the 14 days prior to recruitment
- 13. Subjects taking antibiotics or who have taken antibiotics in the last four weeks, or in the last 6 months, based on the intensity and duration of antibiotic treatment
- 14. Subjects with cognitive disorders that may hinder the response to questionnaires
- 15. Subjects with known allergy to the ingredients of the experimental products (active placebo)
- 16. Subjects with acquired immunodeficiency syndrome from HIV

Date of first enrolment

10/06/2024

Date of final enrolment

17/06/2024

Locations

Countries of recruitment

Italy

Study participating centre COMEGEN Soc. Coop. Sociale

Viale Maria Bakunin, 41 Naples Italy 80126

Sponsor information

Epo S.r.l.

Sponsor details

Via Stadera, 19 Milano Italy 20141 +39 02895571 epo@eposrl.com

Sponsor type

Industry

Website

http://www.eposrl.com

Funder(s)

Funder type

Industry

Funder Name

Epo S.r.l.

Results and Publications

Publication and dissemination plan

Planned publication in a high-impact peer-reviewed journal.

Intention to publish date

28/02/2025

Individual participant data (IPD) sharing plan

The datasets generated and/or analysed during the current study will be published as a supplement to the subsequent results publication.

IPD sharing plan summary

Published as a supplement to the results publication