

# Niacin skin flushing response in evaluating the efficacy of modified electroconvulsive therapy in schizophrenia

<b>Submission date</b>	<b>Recruitment status</b>	<input type="checkbox"/> Prospectively registered
09/02/2025	No longer recruiting	<input type="checkbox"/> Protocol
<b>Registration date</b>	<b>Overall study status</b>	<input type="checkbox"/> Statistical analysis plan
18/02/2025	Completed	<input type="checkbox"/> Results
<b>Last Edited</b>	<b>Condition category</b>	<input type="checkbox"/> Individual participant data
10/03/2025	Mental and Behavioural Disorders	<input checked="" type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Schizophrenia is a group of chronic neurodevelopmental disorders. Drug therapy is the main treatment. Modified electroconvulsive therapy (MECT) is also considered an effective treatment method. The aim of this study was to explore the clinical characteristics of patients undergoing MECT treatment, including positive, negative, and general psychotic symptoms, and the improvement of these characteristics after treatment.

### Who can participate?

Patients with schizophrenia aged 18-65 years who are registered at the Fourth People's Hospital of Wuhu

### What does the study involve?

In this non-randomized study, a total of 70 hospitalized patients with schizophrenia underwent eligibility assessment. Participants were divided into two groups based on whether they met the criteria for treatment-resistant schizophrenia (TRS). Among the 40 patients with TRS, eight declined to undergo MECT. The remaining consenting patients received a 6-week course of MECT combined with antipsychotic medication (SZ-MECT group). Among the 30 patients with non-treatment-resistant schizophrenia (N-TRS), all patients agreed to participate in the study and underwent 6 weeks of antipsychotic medication treatment (SZ-N group).

### What are the possible benefits and risks of participating?

Participants who receive the free treatment may benefit from a reduction in their depressive symptoms. There are no known risks involved with participating.

### Where is the study run from?

The Fourth People's Hospital of Wuhu (China)

### When is the study starting and how long is it expected to run for?

December 2022 to December 2025

Who is funding the study?

1. Health Commission of Anhui Province (China)
2. Department of Science and Technology of Anhui Province (China)

Who is the main contact?

Bingbing Sui, 19956720391@163.com

## Contact information

**Type(s)**

Principal investigator

**Contact name**

Dr Zhu Wenli

**Contact details**

Wuxiashan East Road

Wuhu

China

241000

-

3081586616@qq.com

**Type(s)**

Public, Scientific

**Contact name**

Miss Sui Bingbing

**Contact details**

Wuxiashan East Road

Wuhu

China

241000

-

19956720391@163.com

## Additional identifiers

**Clinical Trials Information System (CTIS)**

Nil known

**ClinicalTrials.gov (NCT)**

Nil known

**Protocol serial number**

Nil known

## Study information

**Scientific Title**

Changes in psychiatric symptoms and niacin skin flushing response after modified electroconvulsive therapy, and the correlation between them in schizophrenia

**Acronym**

NSFR SZ

**Study objectives**

Current interventions as of 10/03/2025:

The primary aim of this study was to compare the clinical characteristics of patients with treatment-resistant schizophrenia (TRS) and non-treatment-resistant schizophrenia (N-TRS), including the severity of clinical symptoms and the niacin skin flushing response (NSFR), which is considered an auxiliary diagnostic biomarker for schizophrenia (SZ), and investigate the relationship between NSFR and the severity of symptoms. Additionally, this study also investigated whether the patients with TRS could achieve comparable clinical remission through modified electroconvulsive therapy (MECT) to that of the patients with N-TRS receiving conventional antipsychotic drugs, to provide further evidence in support of the efficacy of MECT in the treatment of SZ.

Previous interventions:

The aim of this study is to evaluate the relationship between the efficacy of modified electroconvulsive therapy and niacin skin flushing response in schizophrenia, and explore whether niacin skin flushing response is related to psychiatric symptoms.

**Ethics approval required**

Ethics approval required

**Ethics approval(s)**

approved 29/12/2023, Ethical Committee of the Fourth People's Hospital of Wuhu (The Fourth People's Hospital of Wuhu, Wuhu, 241000, China; -; whsyllwyh@163.com), ref: [2023]-KY-052

**Study design**

Single-centre interventional non-randomized controlled trial

**Primary study design**

Interventional

**Study type(s)**

Diagnostic, Treatment, Efficacy

**Health condition(s) or problem(s) studied**

Schizophrenia

**Interventions**

Current interventions as of 10/03/2025:

Participants were divided into two groups based on whether they met the criteria for treatment-resistant schizophrenia (TRS). Among the 40 patients with TRS, 8 declined to undergo MECT. The remaining consenting patients received a 6-week course of MECT combined with antipsychotic medication (SZ-MECT group, n = 32). Among the 30 patients with non-treatment-resistant schizophrenia (N-TRS), all patients agreed to participate in the study and underwent 6 weeks of antipsychotic medication treatment (SZ-N group, n = 30). The drugs include chlorpromazine,

aripiprazole, olanzapine, and risperidone (appropriate dosage and frequency of medication based on the patient's condition). The MECT frequency is 2-3 times a week, adjusted according to the patient's condition.

#### Previous interventions:

The study recruited a total of 62 patients with schizophrenia. 30 schizophrenia individuals took only anti-psychotics (SZ-N group) and 32 schizophrenia patients received MECT combined with anti-psychotics (SZ-MECT group).

The research group combined MECT with antipsychotic drug treatment. The drugs include chlorpromazine, aripiprazole, olanzapine, and risperidone (appropriate dosage and frequency of medication based on the patient's condition). The MECT frequency is 2-3 times a week, adjusted according to the patient's condition. The follow-up time was at baseline, week 2, week 4, and week 6. The total treatment time is adjusted according to the patient's symptoms.

The control group was only treated with antipsychotic drugs, including clozapine, aripiprazole, olanzapine, and risperidone (appropriate dosage and frequency of medication based on the patient's condition).

All patients underwent Positive and Negative Syndrome Scale (PANSS) and niacin skin flushing response (NSFR) scores at baseline (t1) and follow-up (t2, ~4 weeks) time points. Group differences in changes of the NSFR between two schizophrenia groups over time, as well as the correlation between psychiatric symptoms and niacin skin flushing response, were assessed.

#### Intervention Type

Mixed

#### Primary outcome(s)

Psychiatric symptoms measured using the Positive and Negative Syndrome Scale (PANSS) at baseline, weeks 2, 4, and 6.

#### Key secondary outcome(s)

Niacin skin flushing response test using 6 different concentrations (triple gradient dilution from 60 mM) of niacin dropped onto the skin at baseline, week 2, 4, and 6

#### Completion date

31/12/2025

## Eligibility

#### Key inclusion criteria

1. Patients diagnosed according to the ICD10 (International Classification of Diseases, 10th Revision) criteria for SZ
2. 18 to 65 years old
3. PANSS score was within the range of 70~120

#### Participant type(s)

Patient

#### Healthy volunteers allowed

No

**Age group**

Mixed

**Lower age limit**

18 years

**Upper age limit**

65 years

**Sex**

All

**Total final enrolment**

62

**Key exclusion criteria**

1. Severe neurological diseases or traumatic brain injury or substance dependence
2. Administration of non-steroidal or steroid anti-inflammatory drugs within 2 weeks
3. MECT administration 1 month before admission
4. MECT contraindications (including anesthesia allergy, intracranial space-occupying lesions, active pulmonary inflammation, recent myocardial infarction, recent cerebral hemorrhage, retinal detachment, untreated glaucoma, etc)
5. Pregnancy

**Date of first enrolment**

01/01/2024

**Date of final enrolment**

31/12/2024

## Locations

**Countries of recruitment**

China

**Study participating centre**

**The Fourth People's Hospital of Wuhu**

Wuxiashan East Road

Wuhu

China

241000

## Sponsor information

## Organisation

The Fourth People's Hospital of Wuhu

## Funder(s)

### Funder type

Government

### Funder Name

Health Commission of Anhui Province

### Funder Name

Anhui Provincial Department of Science and Technology

### Alternative Name(s)

Anhui Department of Science and Technology, , Department of Science and Technology of Anhui Province

### Funding Body Type

Government organisation

### Funding Body Subtype

Local government

### Location

China

## Results and Publications

### Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be available upon request from Bingbing Sui (19956720391@163.com)

### IPD sharing plan summary

Available on request

### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Participant information sheet</a>	Participant information sheet	11/11/2025	11/11/2025	No	Yes