Radiotherapy after Oesophageal Cancer Stenting (ROCS) study

| Submission date | Recruitment status No longer recruiting | [X] Prospectively registered | | |
|-------------------|---|------------------------------|--|--|
| 04/07/2012 | | [X] Protocol | | |
| Registration date | Overall study status Completed | Statistical analysis plan | | |
| 10/07/2012 | | [X] Results | | |
| Last Edited | Condition category | Individual participant data | | |
| 28/02/2023 | Cancer | | | |

Plain English summary of protocol

https://www.cancerresearchuk.org/about-cancer/find-a-clinical-trial/a-trial-looking-radiotherapy-oesophageal-cancer-difficulty-swallowing-rocs

Contact information

Type(s)

Scientific

Contact name

Dr Douglas Adamson

Contact details

Ward 32 Ninewells Hosptial Dundee United Kingdom DD1 9SY

Type(s)

Public

Contact name

Ms Martina Svobodova

Contact details

Centre for Trials Research (CTR)
College of Biomedical and Life Sciences
Cardiff University
6th Floor, Neuadd Meirionnydd
Heath Park
Cardiff

United Kingdom CF14 4YS +44(0)29 20687463 ROCS@cardiff.ac.uk

Additional identifiers

ClinicalTrials.gov (NCT)

NCT01915693

Protocol serial number

HTA 10/50/49, WCTU030

Study information

Scientific Title

Palliative radiotherapy in addition to self-expanding metal stent for improving dysphagia and survival in advanced oesophageal cancer: Radiotherapy after Oesophageal Cancer Stenting (ROCS) study

Acronym

ROCS

Study objectives

Radiotherapy in addition to self-expanding metal stent (SEMS) placement improves patient-reported dysphagia and increases time to progression in a patient population unable to undergo surgery.

More details can be found at http://www.nets.nihr.ac.uk/projects/hta/105049 Protocol can be found at http://www.nets.nihr.ac.uk/__data/assets/pdf_file/0009/81666/PRO-10-50-49.pdf

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Two-arm open randomised phase III trial with a 1:1 randomisation ratio. A qualitative component in a sub-set of patients.

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Oesophageal cancer requiring stent for relief of dysphagia

Interventions

Arm A: Self-expanding metal stents (SEMS) (Control Arm)

SEMS insertion will be undertaken in accordance with standard local protocols. Covered or partially covered metal stents will be used and the length type and mode of stent placement will be selected by the clinician. Insertion will occur within two weeks of randomisation.

Arm B: SEMS plus external beam radiotherapy (Intervention Arm)

External beam radiotherapy (EBRT) is routinely available at regional cancer centres across the UK. For palliation of dysphagia in oesophageal cancer, a radiotherapy course delivering a tumour absorbed dose of 20Gy in 5 fractions or 30Gy in 10 fractions within 4 weeks of SEMS insertion.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Assess the impact of radiotherapy in addition to self-expanding metal stent (SEMS) placement on time to progression of patient-reported dysphagia in a patient population unable to undergo surgery.

Key secondary outcome(s))

- 1. Assess the impact of combination treatment on core components of health related quality of life
- 2. Assess the impact of radiotherapy in addition to SEMS placement on overall survival
- 3. Measure morbidity associated with the interventions
- 4. Measure re-intervention rates
- 5. Assess the cost of the addition of radiotherapy to SEMS placement

Completion date

30/11/2018

Eligibility

Key inclusion criteria

- 1. Histological confirmation of oesophageal carcinoma excluding small cell histology
- 2. Not suitable for radical treatment (oesophagectomy or radical chemoradiotherapy) either because of patient choice or medical reasons
- 3. Dysphagia clinically assessed as needing stent as primary treatment of the dysphagia
- 4. Age 16 years or over
- 5. Discussion and treatment decision for SEMS placement made by an Upper GI multi-disciplinary team
- 6. Clinician assessment of ability to attend for radiotherapy
- 7. Expected survival of at least 12 weeks
- 8. Written informed consent

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Total final enrolment

220

Key exclusion criteria

- 1. Histology of small cell carcinoma type
- 2. Tumour length of greater than 12 cm
- 3. Tumour growth within 2 cm of the upper oesophageal sphincter
- 4. Endoscopic treatment of the tumour, other than dilatation, planned in the peri-stent period
- 5. Presence of a tracheo-oesophageal fistula
- 6. Presence of a pacemaker
- 7. Previous radiotherapy to the area of the proposed radiotherapy field
- 8. Planned endoscopic treatment of the tumour (e.g. laser) in the immediate peri-stenting period
- 9. Pregnancy

Date of first enrolment

01/10/2012

Date of final enrolment

31/08/2018

Locations

Countries of recruitment

United Kingdom

Scotland

Study participating centre

Ward 32

Dundee United Kingdom DD1 9SY

Sponsor information

Organisation

Velindre NHS Trust (UK)

ROR

https://ror.org/05ntqkc30

Funder(s)

Funder type

Government

Funder Name

NIHR Health Technology Assessment Programme - HTA (UK), 10/50/49

Results and Publications

Individual participant data (IPD) sharing plan

Not provided at time of registration

IPD sharing plan summary

Not provided at time of registration

Study outputs

| Output type | Details | Date created | Date added | Peer reviewed? | Patient-facing? |
|-------------------------------|-------------------------------|--------------|------------|----------------|-----------------|
| Results article | results | 01/04/2021 | 22/02/2021 | Yes | No |
| Results article | | 01/05/2021 | 28/05/2021 | Yes | No |
| Protocol article | protocol | 22/10/2014 | | Yes | No |
| Participant information sheet | Participant information sheet | 11/11/2025 | 11/11/2025 | No | Yes |
| Plain English results | | | 28/02/2023 | No | Yes |