

Vitamin D supplementation compared to placebo in people presenting with their first episode of psychosis neuroprotection design

Submission date 25/02/2015	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered
Registration date 25/02/2015	Overall study status Completed	<input checked="" type="checkbox"/> Protocol
Last Edited 30/12/2021	Condition category Mental and Behavioural Disorders	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Vitamin D is commonly known as the “sunshine hormone” – if your skin is regularly exposed to sunlight, it makes enough vitamin D to keep your body healthy. However, people living in the UK commonly have low vitamin D, because of our lack of strong sunshine. While vitamin D is known to be important for bone health, there is growing evidence that it may also help people recover from various brain diseases. In other words, having enough vitamin D may protect the brain (i.e. it could be ‘neuroprotective’). At this stage, there is a lack of evidence to clarify if adding vitamin D to the standard treatment of people recovering from their first episode of a psychotic disorder can help recovery. The best way to examine this question is with a Randomized Controlled Trial (RCT). To do this we need to compare vitamin D supplements to placebo. A computer program decides which group you are allocated to, so there is an equal chance (50:50) that you will be allocated to either of the two groups. The results are compared to see if one is better. In a ‘blind trial’ you will not know which treatment group you are in. This trial is ‘double blind trial’, meaning that neither you nor the study team will know in which treatment group you are. If adding vitamin D supplements to standard treatment can help people make a better recovery, then in the future we will be able to provide this type of treatment routinely. The Vitamin D (or placebo) supplement will be given to you once a month by a clinician, alongside the regular treatment that you receive from your mental health team. The aim of this study is to explore if the addition of a vitamin D supplement to standard treatments (which may include medications, general support and talking therapies) can help people recover after having their first episode of a psychotic disorder. To do this, we will compare a group of people who are receiving a vitamin D supplement as well as their standard treatment versus those who receive a placebo supplement (dummy supplement that contains no vitamin D) and standard treatment alone.

What does the study involve

Current as of 11/10/2017:

Study participants are recruited from NHS Trust research sites (including Early Intervention Services (EIS), Home Treatment Teams, inpatient units, community teams and Patient Identification Centres (PICS)). Participation in the study is entirely voluntary. Half of the

participants who join the study are randomly allocated to receiving the vitamin D supplement while half receive a 'placebo' treatment (an identical looking treatment but without vitamin D). Each participant is asked to take six drops (just over one teaspoon) of the vitamin D supplement (or placebo) once a month for a total of 6 months. During their participation in the study (6 months), we ask each participant that they do not use other vitamin D supplements that exceed 400 IU/Daily, either alone or as part of a multivitamin preparation. During the study, the participants take part in a number of assessments. A trained research phlebotomist takes a small sample of blood (the equivalent of a few tablespoons). . The purpose of the blood test is to have a measure of what each participants vitamin D levels are when they enter the study and again at the 6 month assessment points. From the same blood sample we will also check on each participants general blood chemistry, including calcium levels and hormones that are linked to vitamin D. Participants are also asked if they would give another blood sample for genetic testing (in many cases, genes can alter vitamin D levels). Each participant is also asked various questions about symptoms, mood, plus any side effects caused by the treatments. To monitor their physical health we ask each participants permission to measure their weight, height, waist and blood pressure measurements. We will repeat questionnaires administered at the start of the study and at 6 months and will review each participants clinical notes to investigate changes over time. At 3 months we perform a blood test to check each participants calcium and hormone levels.

Previous:

Study participants are recruited from Early Intervention in Psychosis Services. Participation in the study is entirely voluntary. Half of the participants who join the study are randomly allocated to receiving the vitamin D supplement while half receive a 'placebo' treatment (an identical looking treatment but without vitamin D). Each participant is asked to take six drops (just over one teaspoon) of the vitamin D supplement (or placebo) once a month for a total of 12 months.. During their participation in the study (1 year), we ask each participant that they do not use other vitamin D supplements, either alone or as part of a multivitamin preparation. During the study, the participants take part in a number of assessments. A trained research phlebotomist takes a small sample of blood (the equivalent of a few tablespoons) from each . The purpose of the blood test is to have a measure of what each participants vitamin D levels were when they entered the study and again at the 6 and 12 month assessment points.. At the end of the study, with their permission, we will provide clinical team of each participant with their most recent Vitamin D level. From the same blood sample we will also check on each participants general blood chemistry, including calcium levels and hormones that are linked to vitamin D. Participants are also asked if they would give another blood sample for genetic testing (in many cases, genes can alter vitamin D levels). Each participant is also asked various questions about symptoms, mood, plus any side effects caused by the treatments. To monitor their physical health we ask each participants permission to measure their weight, height, waist and blood pressure measurements. We will repeat questionnaires administered at the start of the study at 6 and 12 months and will review each participants clinical notes to investigate changes over time. At 3 and 9 months we perform a blood test to check each participants calcium and hormone levels.

What are the possible benefits and risks of participating?

It is hoped that the participants will benefit from the treatment by having closer contact with the clinic. If they also receive the Vitamin D supplement then this may be good for bone and muscle health.

Where is the study run from?

South London and Maudsley NHS Foundation Trust (UK)

When is the study starting and how long is it expected to run for?
April 2015 to December 2019 (as of 26/10/2018)

Who is funding the study?
Stanley Medical Research Institute (USA)

Who is the main contact?
Dr. Gabriella Wojewodka, gabriella.wojewodka@kcl.ac.uk

Contact information

Type(s)
Scientific

Contact name
Dr Gabriella Wojewodka

Contact details
Department of Psychosis Studies
Institute of Psychiatry
Box PO63
De Crespigny Park
London
United Kingdom
SE5 8AF
-
gabriella.wojewodka@kcl.ac.uk

Type(s)
Public

Contact name
Dr Gabriella Wojewodka

Contact details
Department of Psychosis Studies
Institute of Psychiatry
Box PO63
De Crespigny Park
London
United Kingdom
SE5 8AF
-
gabriella.wojewodka@kcl.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)
2014-002639-32

Protocol serial number

18385

Study information

Scientific Title

A randomised, double blind, placebo controlled parallel group trial of vitamin D supplementation compared to placebo in people presenting with their First Episode of psychosis
Neuroprotection Design

Acronym

DFEND

Study objectives

The aim of this study is to investigate the effects of vitamin D supplementation in people presenting with their first episode of psychosis.

Ethics approval required

Old ethics approval format

Ethics approval(s)

14/LO/1588

Study design

Randomised; Interventional; Design type: Treatment

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Psychosis

Interventions

Vitamin D3: We will use a safe and convenient monthly treatment regimen in the DFEND trial. Under the direct supervision of trial staff, we will administer 120,000 IU of vitamin D3, using the widely used oral Vigantol oil preparation. This potent form of vitamin D has been used often in clinical trials, and comparable strategies are currently underway in large population based randomised controlled trials in the USA, New Zealand, Australia and the United Kingdom (total samples approx. 90,000 individuals).

Follow Up Length: 12 months (Updated 11/10/2017: 6 months)

Study Entry : Single Randomisation only

Intervention Type

Drug

Phase

Phase II

Primary outcome(s)

Current primary outcome measure as of 24/05/2021:

Psychosis symptom severity assessed using the total Positive and Negative Syndrome Scale (PANSS) at 6-month follow-up

Previous primary outcome measure as of 11/10/2017:

PANSS is used at baseline, 3 and 6 months.

Previous primary outcome measure:

PANSS; Timepoint(s): 6 months, 12 months

Key secondary outcome(s)

Current secondary outcome measures as of 24/05/2021:

1. Psychosis symptom severity assessed using the total Positive and Negative Syndrome Scale (PANSS) at 6-month follow-up
 2. Positive symptom severity assessed using the PANSS Positive Scale subscore at 3 and 6 months
 3. Negative symptom severity assessed using the PANSS Negative Scale subscore at 3 and 6 months
 4. Cognitive symptom severity assessed using the PANSS General Psychopathology Scale subscore at 3 and 6 months
 5. Ability to function assessed using Global Assessment of Function (GAF) at 6 months
 6. Depression assessed using the Calgary Depression Scale (CDS) at 6 months
 7. Waist circumference (cm) at 6 months
 8. Body mass index (BMI) (kg/m²) at 6 months
 9. HbA1c (mmol/mol) at 6 months
 10. Total cholesterol (mmol/l) at 6 months
 11. C-reactive protein (CRP)(mg/l) at 6 months
 12. Vitamin D (25(OH)D) concentrations at 6 months
-

Previous secondary outcome measures as of 11/10/2017:

1. Calgary Depression Scale is used at baseline and 6 months
2. GAF scale is used at baseline and six months

Previous secondary outcome measures:

1. Calgary Depression Scale; Timepoint(s): 6, 12 months
2. GAF scale; Timepoint(s): 6, 12 months

Completion date

20/12/2019

Eligibility

Key inclusion criteria

Current inclusion criteria as of 11/10/2017:

1. Aged between 18-65 years including women of child-bearing age
2. Diagnosis of functional psychosis defined according to ICD-10 criteria
3. Willing to refrain from taking multivitamins or non-study vitamin D supplements (including cod liver oil), that exceed 400IU/day of vitamin D throughout the study
4. Patients who are willing to give a vitamin D blood sample
5. Patients who are able to and have given written informed consent

Previous inclusion criteria:

1. Patients experiencing their first episode of psychosis (or FEP, defined as first presentation in the last six months)
2. Attending clinical services run in the South London and Maudsley Hospital NHS Foundation Trust.
3. Aged 18-45 years
4. Must have capacity to provide written informed consent and have sufficient English language skills to complete the study
5. Subjects must agree to refrain from taking multivitamin or non-study vitamin D supplements throughout the study
6. Must be willing to provide a vitamin D blood sample at baseline

Psychosis will be defined according to ICD-10 criteria for psychosis (codes F20-29 and F3033) and confirmed with an OPCRIT (OPERational CRITERia) diagnosis.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

65 years

Sex

All

Total final enrolment

149

Key exclusion criteria

Current exclusion criteria as of 11/10/2017:

1. Known intolerance of Vitamin D2 or D3 or known allergy to any of the trial medications
2. Those who are currently taking vitamin D supplements at a dose exceeding 400IU/day.
3. Those who have taken cardiac glycosides; calcium channel blockers; or oral, intramuscular, or intravenous corticosteroids; bendroflumethiazide; isoniazid, or rifampicin in the past one month
4. Known active tuberculosis, sarcoidosis, hypo or hyperparathyroidism, past or present

nephrolithiasis (renal stones), suspected or diagnosed hepatic or renal dysfunction, any malignancy other than non-melanoma skin cancer not in remission for ≥ 3 years, calcium disorders

5. Baseline corrected serum calcium > 2.6 mmol/L

6. Patients with known history of hypercalcaemia

7. Pregnant or breast-feeding women and women planning a pregnancy

8. Patients lacking the capacity to provide written informed consent

Previous exclusion criteria:

1. Patients whose diagnosis was evaluated retrospectively and found not to fulfil the diagnostic criteria.

2. Individuals who are suicidal at baseline

3. Those with known endocrine disorders, CVS disease, or diabetes

4. Those with contraindications to Vigantol or prescribed cardiac glycosides

5. Pregnant women and women planning a pregnancy

Date of first enrolment

01/01/2016

Date of final enrolment

14/06/2019

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

South London and Maudsley NHS Foundation Trust

King's College London

SLaM R&D and KCL joint office

Institute Of Psychiatry, Psychology and Neuroscience

16 De Crespigny Park

London

United Kingdom

SE5 8AF

Study participating centre

Southern Health NHS Foundation Trust

Tom Rudd Unit

Moorgreen Hospital

Botley Road

West End

Southampton
United Kingdom
SO30 3JB

Study participating centre

Kent and Medway NHS and Social Care Partnership Trust (KMPT)

Clinical Research Network: Kent Surrey Sussex (CRN:KSS)

KMPT Research & Development,

Beech House

Hermitage Lane

Maidstone

United Kingdom

ME16 9PH

Study participating centre

Cheshire and Wirral Partnership NHS Foundation Trust

Research Department

NIHR Clinical Research Network: North West Coast

Churton House

Countess of Chester Health Park

Chester

United Kingdom

CH2 1BQ

Study participating centre

South West London and St George's Mental Health NHS Foundation Trust

Clinical Research Unit

Barnes Hospital

South Worple Way

London

United Kingdom

SW14 8SU

Sponsor information

Organisation

King's College London

ROR

<https://ror.org/0220mzb33>

Funder(s)

Funder type

Research organisation

Funder Name

Stanley Medical Research Institute

Alternative Name(s)

The Stanley Medical Research Institute, SMRI

Funding Body Type

Private sector organisation

Funding Body Subtype

Research institutes and centers

Location

United States of America

Results and Publications

Individual participant data (IPD) sharing plan

Not provided at registration

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		01/12/2021	30/12/2021	Yes	No
Protocol article	protocol	06/01/2020	08/01/2020	Yes	No
HRA research summary			28/06/2023	No	No