

Effects of fermented milk on prevention of antibiotic-associated diarrhoea and Clostridium difficile infection in patients undergoing planned bone surgery

Submission date 26/12/2014	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 18/01/2015	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 19/02/2020	Condition category Digestive System	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Antibiotics are used in many conditions and are often given to patients to protect them against infection at the time of surgery. Diarrhoea can occur after the use of antibiotics and is a common problem that is difficult to avoid. It can delay recovery from surgery and in the worst cases can be a serious problem for the patient's health. Evidence is emerging that probiotics, 'friendly' bacteria usually in special drinks or yoghurts, might reduce this risk. The aim of this study is to investigate whether a daily bottle of Yakult, compared with placebo, can reduce the rate of antibiotic-associated diarrhoea in patients having planned orthopaedic surgery at the Elective Orthopaedic Centre (UK).

Who can participate?

Patients who have been referred for an elective orthopaedic operation at the Elective Orthopaedic Centre (UK)

What does the study involve?

Participants are randomly allocated to one of two groups: Yakult or placebo. They need to refrain from consuming any probiotics in the 8 weeks before their surgery. They might be telephoned around 3 weeks before the operation as a reminder, and to check that nothing has changed and that they meet the criteria for entry into the study. Patients are then provided with a supply of Yakult or placebo and should start drinking one bottle per day for 2 weeks before the surgery and for a further 2 weeks after the surgery, recording this information on the sheets provided. The operation and treatments proceed as normal and patients' care is as standard. After the operation, and for the next 4 weeks, patients need to record their stool appearance using a chart that they are provided with. If they have diarrhoea in the 4 weeks following the operation, as defined by three or more watery stools on two or more consecutive days, they need to inform us by telephone or email and send a stool sample for testing. This testing is part of the patients' clinical care because diarrhoea can be serious in some cases and can be treated if the stool sample shows the cause. Patients also need to complete a questionnaire about

gastrointestinal symptoms, which is short and simple, at the time of providing consent, in the hospital 2 weeks and and 4 weeks after surgery.

What are the possible benefits and risks of participating?

The main possible benefit would be a reduced risk of antibiotic-associated diarrhoea, which could otherwise prolong recovery from surgery and could be detrimental to health. There are no known safety issues surrounding this study.

Where is the study run from?

Epsom General Hospital, Elective Orthopaedic Centre (UK)

When is the study starting and how long is it expected to run for?

April 2010 to August 2014

Who is funding the study?

1. Yakult Ltd (UK)
2. Elective Orthopaedic Centre (UK)
3. St George's Charitable Trust (UK)

Who is the main contact?

Dr Carl Moran

Contact information

Type(s)

Scientific

Contact name

Dr Carl Moran

Contact details

St Richard's Hospital
Spitalfield Lane
Chichester
United Kingdom
PO19 6SE

Additional identifiers

Protocol serial number

v9

Study information

Scientific Title

Effects of Yakult (*Lactobacillus casei* Shirota) on prevention of antibiotic-associated diarrhoea and *Clostridium difficile* infection in patients undergoing elective orthopaedic surgery

Study objectives

Maintenance of a healthy commensal microbiota and the gut's colonisation resistance by the administration of once daily probiotic *Lactobacillus casei* Shirota in the form of fermented milk (Yakult), before and after elective orthopaedic surgery, can help reduce the risk of diarrhoea associated with antibiotics and *Clostridium difficile*.

Ethics approval required

Old ethics approval format

Ethics approval(s)

National Research Ethics Service, North London REC 3, 27/09/2010, ref: 10/H0709/52

Study design

Single-centre double-blind randomised placebo-controlled study

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

Infections

Interventions

Patients will be randomly allocated to one of two groups. One group will receive a daily Yakult drink for 2 weeks before and 2 weeks after the elective operation. The other group will receive a placebo version of the drink for the same period. The placebo drink will not have the active ingredient of Yakult, but will look and taste identical and will be packaged in the same way. The placebo drink will also be manufactured by Yakult.

Intervention Type

Supplement

Primary outcome(s)

Incidence of antibiotic-associated diarrhoea during 4 weeks after surgery. This will be assessed in a hospital and then reported by participants following discharge through direct contact with us or records of their bowel habits over the 4 weeks following the operation. The stool chart will then be sent back to us in a pre-paid envelope.

Key secondary outcome(s)

1. Duration of antibiotic-associated diarrhoea: this will either be recorded directly if the diarrhoea occurs while the patient is still in hospital or by the participant on a stool chart that covers the 4 weeks after the operation. The recorded information is then sent back by the participant in a pre-paid envelope.
2. The incidence of *Clostridium difficile* diarrhoea, as diagnosed with a positive *Clostridium difficile* toxin test and stool culture to exclude other enteric pathogens in all patients with undiagnosed diarrhoea. Any inpatients with diarrhoea will have a stool sample sent for analysis, as per standard hospital care. Participants are asked to inform the trialists of the diarrhoea if it occurs in the 4 weeks after surgery, but following discharge from hospital, so that they can have a stool sample taken (as part of standard care).

3. Gastrointestinal symptoms recorded with the Gastrointestinal Symptom Rating Scale before and after surgery. They are recorded face-to-face at the time of consent and at the time of the operation. Participants record the symptoms themselves at 2 weeks and 4 weeks after the surgery and send the results in pre-paid envelopes.

Completion date

01/08/2014

Eligibility

Key inclusion criteria

1. Undergoing elective joint replacement surgery at the Elective Orthopaedic Centre (UK)
2. Must be able to adhere to the protocol requirements
3. Must be capable of giving informed consent
4. At least 18 years old

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

1. Diarrhoea on pre-admission or within the preceding week
2. Lactose intolerance or intolerance to dairy products
3. Regular probiotic consumption in the previous 8 weeks
4. Antibiotic use in the previous 4 weeks
5. Patients unable to give written informed consent
6. Immunosuppression
7. Active inflammatory bowel disease or bowel surgery less than 6 months ago, cancer, leukaemia or pancreatitis

Date of first enrolment

21/07/2011

Date of final enrolment

15/08/2012

Locations

Countries of recruitment

United Kingdom

England

Study participating centre
Epsom General Hospital
Elective Orthopaedic Centre
Denbies Wing
Surrey
United Kingdom
KT18 7EG

Sponsor information

Organisation
St George's University of London

ROR
<https://ror.org/040f08y74>

Funder(s)

Funder type
Industry

Funder Name
Yakult UK Ltd (UK)

Funder Name
Elective Orthopaedic Centre (UK)

Funder Name
St George's Charitable Trust (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not expected to be made available