

Does the use of sutures reduce the wound complication rate following surgery in hip fractures?

Submission date 13/05/2020	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 18/01/2021	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 18/01/2021	Condition category Surgery	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

The femoral neck is the most common location for a hip fracture. About 75,000 femoral neck are treated by the NHS every year. The reported incidence of wound complications is around 10%. Recent research has suggested an increased risk of infection. The aim of this study is to investigate wound healing and complications related to surgery following neck of femur fracture in the elderly. The aim is to compare wound problems and infection following two different methods of skin closure: subcuticular monocryl suture or metal clips.

Who can participate?

All patients admitted with intracapsular or extracapsular neck of femur fracture and undergoing either hemiarthroplasty or Dynamic Hip Screw operation and with mental capacity.

What does the study involve?

Participants are randomly allocated to be treated with either subcuticular monocryl suture or metal clips for skin closure after surgery. Wound problems, infection and duration of surgery are compared.

What are the possible benefits and risks of participating?

The possible benefits are that the wound healing process can be closely monitored associated with the standard wound closure methods used (sutures and metal clips). The researchers do not anticipate any risks as such.

Where is the study run from?

Medway Maritime Hospital (UK)

When is the study starting and how long is it expected to run for?

April 2012 to January 2018

Who is funding the study?

Investigator initiated and funded

Who is the main contact?

Mr Rajesh Bawale, rajesh.bawale@nhs.net

Prof. Bijayendra Singh, bijayendra.singh@nhs.net

Contact information

Type(s)

Scientific

Contact name

Mr Rajesh Bawale

ORCID ID

<https://orcid.org/0000-0002-3127-0105>

Contact details

Medway Maritime hospital

Windmill road

Gillingham

Gillingham

United Kingdom

ME7 5NY

+44 (0)1634 830000

rajesh.bawale@nhs.net

Type(s)

Scientific

Contact name

Prof Bijayendra Singh

Contact details

Medway Maritime hospital

Windmill road

Gillingham

Gillingham

United Kingdom

ME7 5NY

+44 (0)1634 830000

bijayendra.singh@nhs.net

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

Nil known

Study information

Scientific Title

Use of sutures reduces wound complication rate following surgery in hip fractures - a prospective randomised trial

Study objectives

Orthopaedic surgical site infections are one of the most important and often devastating post-operative complications and can prolong hospital stay by a few days to weeks. Hence choosing the appropriate method of wound closure to prevent wound complications is vital. There is no consensus in the literature as to which skin closure method is better. The aim of this study is to compare the wound problems and infection following two different methods of skin closure: subcuticular sutures and metal clips.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 27/02/2012, NRES Committee London - Queen Square (Room 4W/12, 4th Floor West, Charing Cross Hospital, Fulham, Palace Road, London, W6 8RF, UK; +44 (0)20 3311 7287; queensquare.rec@hra.nhs.uk), REC ref: 11/LO/1452

Study design

Prospective single-centre randomized parallel trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Recovery from surgery for fractured neck of femur

Interventions

The randomisation and the allocation are done by using sequentially numbered opaque and unmarked sealed envelopes.

After completion of the hip hemiarthroplasty and dynamic hip screw surgery, the surgical wounds are closed with either metal clips or the 3-0 absorbable poliglecaprone as allocated following randomisation.

Intervention Type

Procedure/Surgery

Primary outcome(s)

1. Wound problems measured using Singh's Wound Score (SWS) at days 2, 5, 7, 10 and 14 days
2. Wound infection measured using Singh's Wound Score (SWS) at days 7,10 and 14 days

Key secondary outcome(s)

Duration of surgery (min) measured using patient records at the end of surgery

Completion date

15/01/2018

Eligibility**Key inclusion criteria**

1. Admitted with intracapsular or extracapsular neck of femur fracture
2. Undergoing either hemiarthroplasty or dynamic hip screw operation
3. Mental capacity

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Total final enrolment

526

Key exclusion criteria

Does not meet inclusion criteria

Date of first enrolment

24/07/2012

Date of final enrolment

04/03/2017

Locations**Countries of recruitment**

United Kingdom

England

Study participating centre

Medway Maritime Hospital

Windmill Road

Gillingham

Gillingham
United Kingdom
ME7 5NY

Sponsor information

Organisation

Medway Maritime Hospital

ROR

<https://ror.org/02380m508>

Funder(s)

Funder type

Other

Funder Name

Investigator initiated and funded

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Prof. B Singh (bijayendra.singh@nhs.net) and Mr R Bawale (rajesh.bawale@nhs.net).

IPD sharing plan summary

Available on request