# A telephone-based case management intervention for patients at risk of high emergency department utilisation in the English NHS

Submission date	Recruitment status No longer recruiting	[X] Prospectively registered		
23/02/2015		☐ Protocol		
Registration date	Overall study status Completed  Condition category Other	Statistical analysis plan		
05/03/2015		Results		
Last Edited		Individual participant data		
11/06/2019		Record updated in last year		

#### Plain English summary of protocol

Background and study aims

There is a relatively large group of patients with repeated emergency visits and admissions who use a disproportionate amount of health care resources. Several studies have shown that programmes (interventions) that help people to develop the skills, knowledge and confidence to manage their own health (patient activation) can result in heath improvements, increased quality of life and fewer unplanned contacts with healthcare services without any increases in planned care. We want to test a proactive health coaching protocol designed for patients at high risk of being admitted to an emergency healthcare department. It involves regular and individualized contact with a trained nurse and includes care coordination efforts (where a number of healthcare professionals work together to deliver appropriate healthcare to a patient), advice and training. The overall goal of this study is to investigate whether this health coaching protocol can reduce avoidable use of the healthcare system by frequent visitors to emergency departments.

#### Who can participate?

Adults that are considered to be at high risk of hospital admissions in the next 6 months according to a risk prediction model.

#### What does the study involve?

Participants are randomly allocated into one of two groups. Those in group 1 (intervention group) are allocated a designated health coach. Following an initial assessment, the health coach regularly contacts the patient by telephone while also working with others, such as healthcare providers or social services, in order to improve and integrate care for that patient. Those in group 2 (control group) are not allocated a health coach. Participants in both the intervention and the control group are assessed in terms of how much they use healthcare services over a period of four years.

What are the possible benefits and risks of participating? There are no benefits or risks to participating in this study.

Where is the study run from?

The study is conducted in participation with the NHS Vale of York and NHS Harrow Clinical Commissioning Groups and Hospitals.

When is the study starting and how long is it expected to run for? February 2015 to April 2023 (updated 11/06/2019, previously: April 2021)

Who is funding the study? Health Navigator AB (Sweden)

Who is the main contact?

- 1. Joachim Werr (joachim.werr@healthnavigator.se)
- 2. Eleitha Medina (eleitha.medina@health-navigator.co.uk)

#### Contact information

#### Type(s)

Scientific

#### Contact name

Mr Joachim Werr

#### Contact details

Health Navigator AB
Sveavägen 63
Stockholm
Sweden
SE-113 59
+46 70 626 65 71
joachim.werr@healthnavigator.se

#### Type(s)

Scientific

#### Contact name

Ms Eleitha Medina

#### Contact details

One King Street London United Kingdom W6 9HR +44 79 9973 2622 eleitha.medina@health-navigator.co.uk

#### Additional identifiers

#### Clinical Trials Information System (CTIS)

2015-000810-23

#### ClinicalTrials.gov (NCT)

Nil known

#### Protocol serial number

Nil known

### Study information

#### Scientific Title

Implementation of a telephone-based case management intervention for patients at risk of high emergency department utilisation in the English NHS

#### Study objectives

It is possible to reduce the rate of emergency admissions and emergency outpatient visits among patients with a predicted high healthcare utilisation

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

NRES Committee London - Bromley, 07/07/2015, ref: 15/LO/0992

#### Study design

Randomized controlled trial

#### Primary study design

Interventional

#### Study type(s)

Prevention

#### Health condition(s) or problem(s) studied

The study is based among patients deemed to have a high risk of emergency admission

#### **Interventions**

Each participant assigned to the intervention group will be allocated a designated health coach. Following an initial assessment, the health coach will be in regular contact with the patient by telephone while also engaging relevant other stakeholders (e.g. health care providers, or social services) with the objective of improving and integrating care for that patient.

#### Intervention Type

Behavioural

#### Primary outcome(s)

- 1. Emergency hospitalizations, ascertained using hospital electronic medical records
- 2. Emergency hospital outpatient visits, ascertained using hospital electronic medical records

These will be measured at the end of follow-up, i.e. after 2 years.

#### Key secondary outcome(s))

- 1. Non-elective bed days, ascertained using hospital electronic medical records
- 2. Total hospital admissions, ascertained using hospital electronic medical records
- 3. Total bed days, ascertained using hospital electronic medical records
- 4. A&E attendances, ascertained using hospital electronic medical records
- 5. Total health-care costs, ascertained using hospital electronic medical records
- 6. Patient activation, ascertained using PAM-13
- 7. Quality of life, ascertained using SF-12

These will be measured at the end of follow-up, i.e. after 2 years.

#### Completion date

01/04/2023

# Eligibility

#### Key inclusion criteria

- 1. Patients residing in the studied region
- 2. Patients 18 years or older
- 3. Patients considered to be at high risk of non-elective admissions in the next 6 months according to a risk prediction model. The prediction model is based on patient administrative data, and makes an automated risk assessment based on the patient's sex, number of diagnoses (registered diagnoses in the past 2 years), planned and unplanned hospital contacts in the past 6 months and readmissions. The risk assessment is based on a logistic regression model which will be developed and validated specifically for the study.

#### Participant type(s)

**Patient** 

#### Healthy volunteers allowed

No

#### Age group

Adult

#### Lower age limit

18 years

#### Sex

All

#### Key exclusion criteria

- 1. Patients who have had contact with healthcare services within the past six months, and have one or more of the following diagnoses (not necessarily the diagnosis relating to their contacts):
- 1.1. Dementia
- 1.2. Psychotic disorders
- 1.3. Mental disorders caused by misuse
- 1.4. Terminal cancer

- 2. Estimated remaining life expectancy of < 1 year
- 3. Has undergone major surgery in the last six months or planned to undergo major surgery in the foreseeable future
- 4. Severe hearing loss
- 5. Language difficulties that require an interpreter
- 6. Level of cognitive ability which is not sufficient for receiving and responding to telephone counselling
- 7. Does not have access to a telephone connection

#### Date of first enrolment

04/05/2015

#### Date of final enrolment

05/05/2021

#### Locations

#### Countries of recruitment

United Kingdom

England

# Study participating centre NHS Vale of York

York United Kingdom YO30 4GQ

# Study participating centre NHS Harrow

London United Kingdom HA1 3AW

# Sponsor information

#### Organisation

Health Navigator AB

## Funder(s)

#### Funder type

#### Funder Name

Health Navigator AB (Sweden)

# **Results and Publications**

Individual participant data (IPD) sharing plan

#### IPD sharing plan summary

Not expected to be made available

#### **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			28/06/2023	No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes