Creating and maintaining functional residual capacity in preterm neonates with Respiratory Distress Syndrome (RDS)

Submission date	Recruitment status No longer recruiting	Prospectively registered		
20/12/2005		☐ Protocol		
Registration date 20/12/2005	Overall study status Completed	Statistical analysis plan		
		[X] Results		
Last Edited 03/08/2007	Condition category Respiratory	[] Individual participant data		

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Arjan B. Pas, te

Contact details

Department of Pediatrics, Leiden University Medical Center, J6-S, P.O. Box 9600 Leiden Netherlands 2300 RC +31 (0)71 5262909 a.b.te_pas@lumc.nl

Additional identifiers

Protocol serial number N/A

Study information

Scientific Title

Acronym

EFURCI

Study objectives

A sustained inflation with a mechanical device and early nasal continuous positive airway pressure is a more effective management strategy for RDS in preterm infants than conventional intervention with a self-inflating mask and bag, and reduces the requirement for mechanical ventilation and surfactant treatment.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Not Specified

Health condition(s) or problem(s) studied

Respiratory Distress Syndrome, Preterm neonates

Interventions

Resuscitation at birth when necessary:

Group A: Prolonged inflation of 10 seconds with a mechanical device (Neopuff t-piece resuscitator) and early CPAP treatment.

Group B: Inflation with self-inflating mask and bag conform international resuscitation guidelines.

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

Need of endotracheal intubation within 3 days after birth.

Key secondary outcome(s))

- 1. Death
- 2. Bronchopulmonary Dysplasia (BPD)

Completion date

01/07/2007

Eligibility

Key inclusion criteria

- 1. Preterm neonates
- 2. Gestational age range 25 32 weeks (more than 24 weeks and 6 days and not more than 32 weeks and 6 days)

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Neonate

Sex

Not Specified

Key exclusion criteria

Antenatal diagnosed congenital anomalies of the cardio or respiratory system or anomalies incompatible with survival.

Date of first enrolment

01/04/2005

Date of final enrolment

01/07/2007

Locations

Countries of recruitment

Netherlands

Study participating centre

Department of Pediatrics, Leiden University Medical Center, J6-S, P.O. Box 9600

Leiden Netherlands

2300 RC

Sponsor information

Organisation

Leiden University Medical Centre (LUMC) (Netherlands)

ROR

https://ror.org/027bh9e22

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Leiden University Medical Centre (The Netherlands)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	Results	01/08/2007		Yes	No