

Bridging of upper part of small bowel with stomach tubing for facilitating weight loss of morbidly obese patients

Submission date 24/05/2020	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 29/05/2020	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 04/03/2022	Condition category Surgery	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Weight loss surgery, also called bariatric or metabolic surgery, is sometimes used as a treatment for people who are very obese. It can lead to significant weight loss and help improve many obesity-related conditions, such as type 2 diabetes or high blood pressure. But it's a major operation and in most cases should only be considered after trying to lose weight through a healthy diet and exercise.

This study aims to test the safety of a new procedure that combines elements of existing procedures.

Who can participate?

Patients aged over 18 and below 65 years with BMI over 40 kg/m² (without comorbidity related to morbid obesity) or 35 kg/m² (with comorbidity related to morbid obesity, especially glucose metabolism)

What does the study involve?

Patient assessment, performing surgery, and follow up at 1, 3, 6 and 12 months postoperatively.

What are the possible benefits and risks of participating?

Favourable weight loss management is expected due to reducing stomach volume and duodeno-jejunal exclusion but operations may have some risks (gastric wall prolapse, leakage, stricture, bleeding, bile reflux, vomiting, diarrhea, hair loss, nutrient deficiency, hypoproteinaemia and anaemia due to malabsorption)

Where is the study run from?

Kanizsai Dorottya Hospital (Hungary)

When is the study starting and how long is it expected to run for?

October 2018 to October 2020

Who is funding the study?
Kanizsai Dorottya Hospital (Hungary)

Who is the main contact?
Dr Istvan Bence Balint, balint.istvan.bence@kmmk.hu

Contact information

Type(s)
Scientific

Contact name
Dr Istvan Bence Balint

ORCID ID
<https://orcid.org/0000-0002-2066-861X>

Contact details
Erkel Ferenc
1/1/3
Nagykanizsa
Hungary
8800
+36 82501300/6238
balint.istvan.bence@kmmk.hu

Additional identifiers

Clinical Trials Information System (CTIS)
Nil known

ClinicalTrials.gov (NCT)
Nil known

Protocol serial number
KDK-2071-2/2019

Study information

Scientific Title
Laparoscopic Single-Anastomosis duodeno-ileal bypass with Gastric plication (SADI-GP) in the management of morbid obesity (LASAGNE)

Acronym
LASAGNE

Study objectives
Laparoscopic single-anastomosis duodeno-ileal bypass with gastric plication is expected to be a safe procedure and have low complication rates and favourable weight loss outcomes.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 05/07/2019, Kanizsai Dorottya Hospital (Hospital of Nagykanizsa) Institutional Review Board (8800, Nagykanizsa, Szekeres, J.u. 2-8, Hungary; +36 93502092; titkarsag@nkkorhaz), ref: KDK-2071-2/2019

Study design

Interventional non-randomised study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Laparoscopic surgery for severe obesity and related metabolic disorders

Interventions

Patients were operated on to perform laparoscopic single anastomosis duodeno-ileal bypass with gastric plication. Follow-ups were at 1, 3 and 6 months and are planned at 12 months.

Intervention Type

Procedure/Surgery

Primary outcome(s)

Postoperative complications (gastric wall prolapse, leakage, stricture, bleeding, bile reflux, vomiting, diarrhea, hair loss, nutrient deficiency, hypoproteinaemia and anaemia due to malabsorption, venous thromboembolism, pulmonary and cardiovascular events, 30-day hospital readmission) were classified by Clavien-Dindo

Key secondary outcome(s)

At follow up visits (1,3,6 and 12 months), patients were assessed again by physical examination, blood tests (blood count, ionogram, serum protein, glucose, HgbA1C, iron binding capacity, lipid profile, kidney and liver function, hemostasis) and BAROS-Moorehead-Ardelt II and Weiner et al. questionnaires. Weight, BMI adjusted to age-gender-ethnics, ideal weight (at BMI: 25), excess of weight, excess of BMI, weight loss, EWL% and TWL% were measured as weight loss outcomes

Completion date

29/10/2020

Eligibility

Key inclusion criteria

1. Age over 18 and below 65 years
2. BMI over 40 kg/m² (without comorbidity related to morbid obesity) or 35 kg/m² (with

comorbidity related to morbid obesity, especially glucose metabolism)

3. Patient agreement for follow-ups

4. Obtained informed consent

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

36

Key exclusion criteria

1. Bariatric surgery in previous history
2. Severe mental disorders (drug addiction, alcohol consumption, the use of antipsychotics)
3. Socially vulnerable patients
4. Complete immobilization
5. Patients who did not understand the purpose of the study and bariatric surgery

Date of first enrolment

16/10/2018

Date of final enrolment

30/06/2019

Locations

Countries of recruitment

Hungary

Study participating centre

Kanizsai Dorottya Hospital

Szekeres Jozsef street 2-8.

Nagykanizsa

Hungary

8800

Sponsor information

Organisation

Kanizsai Dorottya Hospital

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Kanizsai Dorottya Hospital

Results and Publications

Individual participant data (IPD) sharing plan

All data generated or analysed during this study will be included in the subsequent results publication.

IPD sharing plan summary

Other

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		17/08/2021	04/03/2022	Yes	No
Protocol file		27/05/2020	08/06/2020	No	No