

# Effect of a multivitamin supplement in patients with cervical squamous intraepithelial lesions and human papillomavirus

|  |  |   |
|--|--|---|
| <b>Submission date</b><br>24/07/2025   | <b>Recruitment status</b><br>No longer recruiting        | <input type="checkbox"/> Prospectively registered<br><input type="checkbox"/> Protocol                                  |
| <b>Registration date</b><br>25/07/2025 | <b>Overall study status</b><br>Completed                 | <input type="checkbox"/> Statistical analysis plan<br><input type="checkbox"/> Results                                  |
| <b>Last Edited</b><br>28/07/2025       | <b>Condition category</b><br>Infections and Infestations | <input type="checkbox"/> Individual participant data<br><input checked="" type="checkbox"/> Record updated in last year |

## Plain English summary of protocol

### Background and study aims

This study aimed to find out whether taking a daily multivitamin could help women with early cervical lesions fight the human papillomavirus (HPV) and reduce the chances of these lesions getting worse. Cervical cancer often starts with small changes in the cervix caused by HPV. Strengthening the immune system through vitamins may help the body clear the virus more effectively.

### Who can participate?

Women between 20 and 60 years old, diagnosed with low-grade cervical lesions (early changes in the cervix) confirmed by a colposcopy (a medical exam using a special magnifying device) and a biopsy. Only women who tested positive for HPV and agreed to participate by signing a consent form were included.

### What does the study involve?

Participants were randomly divided into two groups:

One group took a daily multivitamin with vitamins A, C, D, and E for three months, took a three-month break, and then restarted, for a total of six months.

The other group did not take the vitamins during the same period.

Doctors monitored the size and changes in the lesions using colposcopy at the beginning and after six months. HPV viral load (how much virus was present) was measured using a molecular technique called PCR (Polymerase Chain Reaction). This test works like a “copy machine” for DNA, making millions of copies of the virus’s genetic material so it can be detected and measured. The amount of virus was estimated by comparing the intensity of the PCR bands in a gel (like checking how dark a line appears).

### What are the possible benefits and risks of participating?

The vitamins might help strengthen the immune system, lower HPV levels, and reduce the risk of lesions getting worse. Risks were minimal, as the doses were safe and approved, but some vitamins (especially A, D, and E) can build up in the body, so the intake was carefully controlled.

Where is the study run from?

The study was carried out at the Colposcopy Clinic of Sanitary Jurisdiction II (Ministry of Health) in Ciudad Juárez, Chihuahua, Mexico.

When is the study starting and how long is it expected to run for?

The study started in 2018 and lasted six months for each participant.

Who is funding the study?

The study was funded by the Programa para el Desarrollo Profesional Docente (PRODEP) under the project agreement UACJ-PTC-364/Folio 511-6/17-7605.

Who is the main contact?

Dr Arellano-Ortiz, ana.arellano@uacj.mx

## Contact information

### Type(s)

Public, Scientific, Principal Investigator

### Contact name

Dr Ana Lidia Arellano-Ortiz

### ORCID ID

<https://orcid.org/0000-0002-0985-9191>

### Contact details

Av. Plutarco Elías Calles #1210, Fovissste Chamizal

Juárez

Mexico

32310

+52 6566881800

ana.arellano@uacj.mx

## Additional identifiers

### EudraCT/CTIS number

Nil known

### IRAS number

### ClinicalTrials.gov number

Nil known

### Secondary identifying numbers

RIP12019ICB19

## Study information

### Scientific Title

Multivitamin supplementation versus no supplementation for HPV viral load reduction and lesion progression in cervical squamous intraepithelial lesions

## **Study objectives**

To evaluate the effect of a multivitamin supplement on reducing the progression of cervical squamous intraepithelial lesions.

### **Specific Objectives:**

1. To evaluate the nutritional status of a group of patients with LSIL (Low-Grade Squamous Intraepithelial Lesions) from the Colposcopy Clinic of Sanitary Jurisdiction II in Ciudad Juárez, Chihuahua.
2. To supplement a group of patients with a multivitamin to cover more than 75% of the daily requirements for vitamins A, C, D, and E.
3. To determine the serum and dietary concentrations of these specific nutrients at baseline and after six months of supplementation.
4. To monitor lesion progression through colposcopic analysis and HPV presence at baseline, six, and twelve months after diagnosis.
5. To evaluate the effect of multivitamin supplementation on the regression, progression, or persistence of HPV, comparing it with a group of patients who did not receive supplementation.

### **Hypothesis**

Multivitamin supplementation in patients with low-grade cervical intraepithelial lesions will show a reduction in the progression of squamous intraepithelial lesions compared to those who did not receive supplementation.

## **Ethics approval required**

Ethics approval required

## **Ethics approval(s)**

Approved 21/08/2017, Comité Institucional de ética y bioética de la UACJ (Av. Plutarco Elías Calles #1210 Fovissste Chamizal, Juárez, 32310, Mexico; +52 6566881800; investigacionicb@uacj.mx), ref: CIBE-2017-1-43

## **Study design**

Randomized pilot study

## **Primary study design**

Interventional

## **Secondary study design**

Randomised controlled trial

## **Study setting(s)**

Medical and other records, University/medical school/dental school

## **Study type(s)**

Prevention, Treatment

## **Participant information sheet**

Not available in web format, please use contact details to request a participant information sheet

## **Health condition(s) or problem(s) studied**

Prevention of progression in cervical intraepithelial lesions in patients with HPV

## **Interventions**

A randomized pilot study was conducted between 2018 and 2019 to evaluate the effect of multivitamin supplementation in women with LSIL and HPV. Ethical approval was obtained from the Ethics Committee of the Autonomous University of Ciudad Juárez, and the study complied with the Declaration of Helsinki and national health regulations.

A total of 82 women aged 20–60 years with LSIL confirmed by colposcopy at the Colposcopy Clinic of Sanitary Jurisdiction II in Ciudad Juárez, Chihuahua, were initially recruited. Exclusion criteria included HSIL, cervical cancer, condylomas, pregnancy, baseline HPV-negative results, or refusal to sign informed consent. LSIL diagnosis was confirmed through colposcopy with 5% acetic acid and histopathological biopsy. Cervical samples were collected for HPV detection and DNA extraction, performed using phenol-chloroform-isoamyl alcohol, and HPV was identified by endpoint PCR targeting the L1 capsid gene. Viral load was quantified as relative units (RU/PC) using optical density analysis with ImageJ®.

Sixty-three HPV-positive patients were randomly assigned to an experimental group (n = 31), receiving multivitamin supplementation, or a control group (n = 32) without supplementation. Both groups underwent baseline and six-month evaluations, including dietary assessment, gynecological examination, and viral load measurement.

The experimental group received a multivitamin solution (vitamins A, C, and D) and vitamin E capsules to meet >75% of daily requirements: vitamin A 1200 µg/day, vitamin C 291.3 mg/day, vitamin D 18.75 µg/day, and vitamin E 57.1 mg/day, administered for three months with a three-month washout. Colposcopic outcomes were classified as regression, partial regression, persistence, or progression. Statistical analysis included Student's t-tests, ANOVA, and Chi-square tests with significance set at  $\alpha = 0.05$ .

## **Randomisation:**

Patient recruitment was carried out at the Colposcopy Clinic of Sanitary Jurisdiction II in Ciudad Juárez. Participants were evaluated and assessed by the gynecologist-obstetrician in charge of the clinic, Dr Cecilia Díaz Hernández. Once low-grade squamous intraepithelial lesions were diagnosed, patients were personally invited to participate in the study. Detailed information about the project was provided, and participation was entirely voluntary. Written informed consent was obtained from all participants prior to enrolment. Patients were randomly selected and assigned an identification number. Only the principal investigator was aware of the group allocation for each participant. Neither the gynecologist-obstetrician nor the students involved in the study were informed of the group assignments.

## **Intervention Type**

Supplement

## **Primary outcome measure**

Regression, persistence, or progression assessed by colposcopic evaluation before and after six months of supplementation measured using patient records

## **Secondary outcome measures**

Viral load measured by endpoint PCR and optical densitometry before and after six months of supplementation

**Overall study start date**

01/06/2017

**Completion date**

13/06/2019

## **Eligibility**

**Key inclusion criteria**

1. Women diagnosed with low-grade cervical intraepithelial lesions (LSIL) confirmed by colposcopy and histopathological biopsy
2. Age  $\geq 20$  years
3. Positive for human papillomavirus (HPV) infection

**Participant type(s)**

Patient

**Age group**

Adult

**Lower age limit**

20 Years

**Upper age limit**

60 Years

**Sex**

Female

**Target number of participants**

60

**Total final enrolment**

43

**Key exclusion criteria**

1. Patients with High-Grade Squamous Intraepithelial Lesions (HSIL) , Cervical Cancer, and condylomas
2. Pregnant patients
3. Those who only attended at the beginning and did not return
4. HPV-negative patients at baseline
5. Those who did not sign the informed consent

**Date of first enrolment**

08/02/2018

**Date of final enrolment**

10/05/2018

# Locations

## Countries of recruitment

Mexico

## Study participating centre

Colposcopy Clinic of Sanitary Jurisdiction II (Ministry of Health) (Clínica de Colposcopia de la Jurisdicción Sanitaria II ( Secretaría de Salud))

Paseo Triunfo de La República 3530

Juárez

Mexico

32330

# Sponsor information

## Organisation

Universidad Autónoma de Ciudad Juárez

## Sponsor details

Av. Plutarco Elías Calles #1210 Fovissste Chamizal Ciudad Juárez

Juárez

Mexico

32310

+52 6566882100

dgitt@uacj.mx

## Sponsor type

University/education

## Website

<http://www.uacj.mx/>

## ROR

<https://ror.org/05fj8cf83>

# Funder(s)

## Funder type

University/education

## Funder Name

Fondo Programa para el Desarrollo Profesional Docente (PRODEP)

# Results and Publications

## Publication and dissemination plan

Planned publication in a peer- reviewed journal

## Intention to publish date

## Individual participant data (IPD) sharing plan

The datasets generated and/or analysed during the current study will be published as a supplement to the results publication

## IPD sharing plan summary

Published as a supplement to the results publication

## Study outputs

| Output type                                   | Details    | Date created | Date added | Peer reviewed? | Patient-facing? |
|---|------------|--------------|------------|----------------|-----------------|
| <a href="#">Participant information sheet</a> | in Spanish |              | 25/07/2025 | No             | Yes             |