Safety evaluation of low-level chlorine in water aerosol when inhaled by healthy volunteers

Submission date 03/12/2021	Recruitment status No longer recruiting	Prospectively registeredProtocolStatistical analysis planResults					
Registration date 15/12/2021	Overall study status Completed						
Last Edited 15/12/2021	Condition category Respiratory	Individual participant dataRecord updated in last year					
Plain English summary of protocol Background and study aims. Human white blood cells produce aqueous (dissolved) chlorine that has been reproduced as HS4-20 and used to prevent wound infection and inflammation when administered to the skin and other surfaces. It is believed that this would be beneficial when administered to the airways and lungs to treat lung and airway diseases. The study is designed to discover if HS4-20 is safe to administer as an aerosol and breathe. The study is designed to begin at a short exposure of 30 seconds and, if safe to do so, increase exposure to 90 minutes over several days and in up to 18 healthy volunteers.							
Who can participate Healthy volunteers	??						
What does the stud The participants bro to 90 minutes.	=	4-20 over several days and escalating from 30 seconds					

What are the possible benefits and risks of participating?

There are no expected benefits. The possible risk of participation is increased expectoration (spitting) or mild coughing

Where is the study run from? Hypo-Stream (UK)

When is the study starting and how long is it expected to run for? May 2015 to September 2020.

Who is funding the study? Hypo-Stream Ltd (UK)

Who is the main contact? Myles Dakin myles.dakin@hypo-stream.com

Contact information

Type(s)

Public

Contact name

Mr Myles Dakin

ORCID ID

https://orcid.org/0000-0002-0294-4908

Contact details

Hypo-Stream Cambridge
Unit 9 Beech House
Melbourn Science Park
Melbourn
Cambridge
United Kingdom
SG8 6HB
+44 (0)1763 261129
myles.dakin@hypo-stream.com

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

HS4-20 SE

Study information

Scientific Title

Open-label, exposure-escalation study to evaluate the safety and tolerability of nebulised HS4-20 in healthy volunteers

Acronym

SEHS4-20

Study objectives

That inhalation of HS4-20 via a nebuliser is is safe and without adverse events over long periods of time.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics approval not required. An open-label, exposure escalation study was planned following a safety evaluation via formal scientific advice from the HMRA and subsequent formal scientific advice from the EMA. The study was planned and directed by Prof. R Aspinall (translational medicine) and Dr Thomas Kenny (formerly director at NIHR).

Study design

Open-label exposure-escalation study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Safety evaluation of nebulised HS4-20 physiologic aqueous chlorine aerosol

Interventions

Pulmonary administration of HS4-20 as an aerosol via a nebuliser.

The healthy volunteers were divided into three cohorts:

- 1. Two adults. Exposure escalation from 30 seconds increasing to 30 minutes. A subsequent day exposure escalation from 5 minutes escalating to 90 minutes.
- 2. Four adults. Exposure escalation from 30 seconds to 20 minutes. Subsequent day exposure for 20 minutes repeated twice.
- 3. 12 adults and children. An induction phase of 5 minutes escalating to 20 minutes. Subsequent day exposure of 20 minutes.

Intervention Type

Drug

Phase

Phase I

Drug/device/biological/vaccine name(s)

HS4-20

Primary outcome(s)

Treatment-emergent adverse events measured using observation for the absence of abnormality and of coordination by a clinically trained investigator over a 30-minute period following administration. Follow-up was 7 days post-administration self-reporting. The final follow-up was over a 12-month period of self-reporting and investigator interview.

Key secondary outcome(s))

- 1. Blood pressure measured with an automatic upper arm cuff sphygmomanometer (Omron X3) at baseline (pre-administration), 30 seconds, 2, 5, 10, 20, 30 minutes post-administration in first and second cohorts, the third cohort was measured at 30 seconds, 5, 20, 30 minutes
- 2. Oxygen saturation measured by pulse oximeter (Braun Healthcare Pulse Oximeter 1) throughout the periods of exposure and for 30 minutes after cessation of administration

Completion date

20/09/2020

Eligibility

Key inclusion criteria

Healthy volunteers, no exclusions on basis of age or gender

Participant type(s)

Healthy volunteer

Healthy volunteers allowed

No

Age group

All

Sex

All

Total final enrolment

18

Key exclusion criteria

Does not meet the inclusion criteria

Date of first enrolment

01/06/2015

Date of final enrolment

20/09/2020

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

Hypo-Stream Ltd

Unit 9 Beech House Melbourn Science Park Melbourn Cambridge United Kingdom SG8 6HB

Sponsor information

Organisation

Hypo-Stream Ltd

Funder(s)

Funder type

Industry

Funder Name

Hypo-Stream Ltd

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated and/or analysed during the current study will be published as a supplement to the results publication

IPD sharing plan summary

Published as a supplement to the results publication

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet			09/12/2021	No	Yes
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Protocol file	version 1.0		09/12/2021	No	No