

The weight loss and anti-diabetes outcomes of Sleeve Gastrectomy with Transit Bipartition (SG-TB) and Standard Sleeve Gastrectomy (SG) surgery

Submission date 08/07/2020	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 24/07/2020	Overall study status Ongoing	<input type="checkbox"/> Protocol
Last Edited 21/07/2020	Condition category Nutritional, Metabolic, Endocrine	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Weight loss surgery, also called bariatric or metabolic surgery, is sometimes used as a treatment for people who are very obese. It can lead to significant weight loss and help improve many obesity-related conditions, such as type 2 diabetes or high blood pressure.

Sleeve gastrectomy (SG) and Roux-en-Y gastric bypass (RYGB) are the most popular bariatric procedures currently used. With that being said, SG and RYGB both have limitations of their own. gastroesophageal reflux disease (GERD), diabetes relapse, and weight regain are the issues that may be faced following SG. While RYGB might be able to solve the issues of GERD and diabetes relapse related to SG's, these procedures have an equal weight loss effect. Furthermore, the RYGB procedure creates a risk in that gastric cancer screening is no longer possible. This is worrying especially in countries with high gastric cancer prevalence.

Sleeve gastrectomy with transit bipartition (SG-TB) was introduced in 2012 with the notion that this procedure might be able to solve SG's issues. Currently studies on SG-TB are lacking, especially on the randomized controlled study comparing SG-TB and standard SG procedure. Therefore, this study is aimed to provide an answer for whether SG-TB provides additional benefits against the standard SG.

Who can participate?

Patients aged 18 - 65 years with a BMI of 27 kg/m² or above.

What does the study involve?

Participants will either undergo SG-TB or standard SG surgical procedure over the study period. Participants data on their weight and diabetes conditions as well as prevalence of GERD symptoms and incidence of surgical complications preoperatively and postoperatively will be collected at the start of the study and every 6 months for the 5 years following.

What are the possible benefits and risks of participating?

SG-TB patients might receive longer and superior weight loss and anti-diabetic outcomes than the SG patients. Also fewer SG-TB patients might have GERD symptoms than SG patients.

As SG-TB procedure is a relatively new procedure, there are complications that can occur that have never been previously reported, such as marginal ulcer and severe malnutrition. It has been reported previously that SG-TB might have complications such as incisional or internal hernia and mild malnutrition. On the other hand, standard SG might have increased in leakage and postoperative GERD risks due to the high pressured "sleeve".

Where is the study run from?

Affiliated Hospital of Xuzhou Medical University (China)

When is the study starting and how long is it expected to run for?

August 2019 to December 2026

Who is funding the study?

Affiliated Hospital of Xuzhou Medical University (China)

Who is the main contact?

Dr Jason Widjaja

drjwbs@yahoo.com

Contact information

Type(s)

Scientific

Contact name

Dr Jason Widjaja

ORCID ID

<https://orcid.org/0000-0003-0103-6147>

Contact details

No.99 Huaihai West Road

Affiliated Hospital of Xuzhou Medical University

Xuzhou

China

221006

+86 15094357019

drjwbs@yahoo.com

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

XYFY2020-JS003-01

Study information

Scientific Title

Weight loss and diabetes remission outcomes on SG-TB versus SG: a randomized controlled trial

Study objectives

1. Transit bipartition is superior or equal to sleeve gastrectomy in inducing weight loss.
2. Transit bipartition is superior to sleeve gastrectomy in inducing diabetes remission.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 08/06/2020, Ethical Committee of The Affiliated Hospital of Xuzhou Medical University (Department of General Surgery, the Affiliated Hospital of Xuzhou Medical University, Huaihai West Road No. 99, Xuzhou Jiangsu 221002, P. R. China; no telephone number provided; zhuxccf@163.com), ref: XYFY2020-JS003-01

Study design

Single-centered interventional randomized controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Obesity with or without type-2 diabetes mellitus

Interventions

The study implements a randomized controlled trial (RCT) design (1:1 ratio) involving at least 100 patients undergoing bariatric surgery in total with ≥ 50 in the sleeve gastrectomy with transit bipartition (SG-TB) group and ≥ 50 in the standard sleeve gastrectomy group. The study surgery, data collection, analysis, and evaluation that applies to all groups will be performed by the same individuals. And detailed intervention in intervention arms varies from patient to patient due to the personalized approach. The analysis and reporting of the trial will be in accordance with the CONSORT guidelines.

The overall goal of this particular study is to find out the benefits of SG-TB procedure against the standard SG. In overall:

1. Is SG-TB superior to standard SG in terms of the short-term (1 year) and long-term (>5 years) weight loss outcome?
2. Is SG-TB superior to standard SG in terms of the short-term (1 year) and long-term (>5 years) diabetes remission outcome?
3. The prevalence of gastroesophageal reflux disease (GERD) following both procedures.
4. The early and late complications of SG-TB.

To compare the weight loss efficacy between SG-TB and standard SG, variables that are included are as such: weight (kg), height (m), BMI (calculated using patient's weight and height). In terms of the anti-diabetic outcomes: fasting blood glucose (mmol/L), glycosylated hemoglobin (% HbA1C), fasting insulin and C-peptide; Complete remission is defined as HbA1C <6% without anti-diabetic medications and partial remission as 6-6.49%. These variables will be monitored every 6 months since surgery (at 0, 6, 12, 18, 24, 30, 36, 42, 48, 54, 60 months). In terms of GERD, preoperatively patients will undergo endoscopy for diagnosis as well as history taking of using PPI to solve GERD symptoms; postoperatively all patients will be monitor on the use of PPI as well as performing endoscopy every 1 year since surgery. Early complications are defined as complications that occur within 30 days after the surgery, and late complications as complications occurring in the period following 30 days after surgery.

Intervention Type

Procedure/Surgery

Primary outcome(s)

At surgery, 1-year and 5-year follow-up

1. Weight (kg)
2. BMI (weight, kg divided by the square of height, m)
3. Percentage of Excess BMI Loss, %EBMIL ($[(\text{preoperative BMI} - \text{postoperative BMI}) / (\text{Preoperative BMI} - 25)] \times 100\%$)

Key secondary outcome(s)

1. Diabetes Remission measured at surgery, 1-year and 5-year follow-up using:
 - 1.1. Fasting blood glucose, HbA1C (blood test)
 - 1.2. Anti-diabetic drug usage (patient records)

Completion date

31/12/2026

Eligibility

Key inclusion criteria

1. Age 18-65 years old
2. BMI ≥ 27.5 kg/m²

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

65 years

Sex

All

Key exclusion criteria

1. Type-2 diabetes mellitus of longer than 15 years
2. Previous bariatric surgery
3. Patients debilitated with neurological, cardiovascular and/or malignancies before the procedure
4. Patients refusal to sign the consent form

Date of first enrolment

01/01/2020

Date of final enrolment

31/12/2021

Locations**Countries of recruitment**

China

Study participating centre

The Affiliated Hospital of Xuzhou Medical University

No.99 Huaihai West Road

Department of General Surgery

Affiliated Hospital of Xuzhou Medical University

Xuzhou

China

221006

Sponsor information**Organisation**

Affiliated Hospital of Xuzhou Medical College

ROR

<https://ror.org/011xhcs96>

Funder(s)**Funder type**

University/education

Funder Name

Xuzhou Medical University

Alternative Name(s)

, , Xúzhōu Yīkē Dàxué, XZMU

Funding Body Type

Government organisation

Funding Body Subtype

Universities (academic only)

Location

China

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated and/or analysed during the current study during this study will be included in the subsequent results publication.

IPD sharing plan summary

Other

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes