

The efficacy of a movement control exercise programme to reduce injuries in youth rugby

Submission date 25/03/2015	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 10/04/2015	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 10/05/2021	Condition category Injury, Occupational Diseases, Poisoning	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Injuries to youth rugby players have become a prominent public health concern, raising the importance of developing and implementing appropriate injury prevention strategies. There is growing evidence from other youth sports that supports the use of structured pre-activity exercise routines to reduce the risk of injury. However, studies have yet to investigate the effect of such interventions in youth contact sport populations such as rugby union. The aim of this study is to determine whether the regular use of a specific movement control exercise programme by youth rugby players before training and matches is effective in reducing injury risk from participation. The study will also establish the influence of coach attitudes on the uptake of players to such programmes.

Who can participate?

The study will take place within the independent secondary school system in England. School rugby coaches will be the focus of the intervention as they will be trained to deliver the exercise programme. The effects of the exercise programme will be measured in male youth players aged 14-18 in school rugby programmes over the 2015-16 school winter term by collecting injury risk data.

What does the study involve?

Schools will be randomly allocated to either a pre-session movement control exercise programme or a sham exercise programme, both of which will be coach-delivered. Ongoing monitoring of match and training exposure, use of the exercise programmes, and injury occurrence will allow the effect of the exercise programme to be established. Questionnaires will evaluate coach attitudes, knowledge, and behaviours towards injury prevention both before and after the intervention period.

What are the possible benefits and risks of participating?

Staff in the school rugby programmes will be trained and given resources to deliver pre-activity exercises in their training and pre-match sessions which represent current best practice. These exercises are low to moderate intensity and focus on correct execution of the movement pattern so should present no additional risk to players over and above their normal participation in the sport.

Where is the study run from?

Sport, Health & Exercise Science, Department for Health, University of Bath (UK).

When is the study starting and how long is it expected to run for?

From March 2015 to December 2015.

Who is funding the study?

Rugby Football Union and University of Bath (UK).

Who is the main contact?

Dr Grant Trewartha

g.trewartha@bath.ac.uk

Contact information

Type(s)

Scientific

Contact name

Dr Grant Trewartha

ORCID ID

<http://orcid.org/0000-0002-9021-8956>

Contact details

Department for Health

University of Bath

Claverton Down

Bath

United Kingdom

BA2 7AY

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

N/A

Study information

Scientific Title

The efficacy of a movement control exercise programme to reduce injuries in youth rugby: a cluster randomised controlled trial

Acronym

Study objectives

The intended aims of this study are to assess the efficacy of a pre-activity movement control exercise intervention to reduce the incidence and severity of rugby-related injuries, as well as to assess the influence of coach attitudes on compliance to the exercise programmes.

Specific hypotheses are:

1. The incidence of rugby-related injuries will be reduced following 12 weeks of using the intervention exercise programme when compared with the sham exercise programme.
2. The average severity of rugby-related injuries will be reduced following 12 weeks of using the intervention exercise programme when compared with the sham exercise programme.
3. Coach attitudes towards injury prevention will change as a function of using the intervention exercise programme when compared with the sham exercise programme.
4. Coach attitudes towards injury prevention will demonstrate an influence on compliance to their exercise programmes.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Research Ethics Approval Committee for Health (REACH), University of Bath, 20/03/2015, ref: EP 14/15 141

Study design

Single-centre cluster randomised controlled trial

Primary study design

Interventional

Secondary study design

Cluster randomised trial

Study setting(s)

School

Study type(s)

Prevention

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Sport injury

Interventions

Intervention context

Most rugby players will undertake a number of weekly coach-led training sessions within the school rugby environment in order to prepare appropriately for fixtures. These sessions are predominantly composed of rugby-related games and drills with some supplementary strength

and conditioning sessions, whilst a focus on specific preventive exercise activities is rare. These characteristics would be expected to be consistent in schools across both trial arms as the exclusion criteria for this study will preclude the recruitment of any schools that currently engage in specific preventive training practices outside of normal strength and conditioning sessions.

Control arm: normal training + 'sham' exercise programme

In addition to adhering to normal training regimens, schools in the control group will be given a structured 'sham' programme derived from what is currently regarded as best practice in schools rugby. The programme is structurally indistinct from the intervention programme, with each session occurring at the beginning of each training session or match preparation lasting approximately 15 minutes and with exercises being varied at the same integer (4 weeks). However, the exercises chosen are intended to be distinct from those included in the intervention programme. Each session will be led by the coach or associated member of staff, acting as a delivery agent.

Intervention arm: normal training + intervention exercise programme

Schools in the intervention arm will receive the intervention exercise programme in addition to continuing with usual training practices. The intervention exercise programme in this study will consist of a progressive multifaceted exercise programme which incorporates a blend of training methods such as lower limb balance and agility, whole body strengthening, and plyometrics. The intervention exercise programme is designed to be completed as the initial 15 minutes of every training session and prior to every match, though certain exercises will be withdrawn when the programme is performed prior to matches. As with the control arm, the coach or associated member of staff will act as a delivery agent. The programme is progressive with a new phase being introduced every 4 weeks.

Intervention Type

Behavioural

Primary outcome measure

The incidence and severity of injury, as well as the overall burden of injury in the intervention group compared with the control group.

Details (location, nature, mechanism, severity) of all injuries related to participation in school rugby that incur a 24-hour or greater time loss from subsequent school rugby exposure for the 2015/16 winter term will be recoded on a standardised form. The medical diagnosis for each injury will be coded to three levels according to the Orchard Sports Injury Coding System (OSICS) version by an on-site medical professional.

Coaches of all individual teams will be provided an exposure and compliance report form to be completed on a weekly basis, detailing team-level training and individual match exposure, along with a compliance record in relation to the exercise programme.

Injury and exposure information will be prospectively collected for the entire duration of the schools rugby season from late August 2015 to December 2015.

Poisson regression, with adjustment for clustering by school, offset for player-hours of exposure, and adjustment for the aforementioned covariates (i.e., age/maturation status, playing position, playing experience, level of play, coaches' attitude, and previous injury history) will be used to compare injury incidence and injury burden (injury incidence rate x mean absence per injury) between the trial arms. Injury severity will be compared between the trial arms using an identical model, but with a linear response distribution. Separate analyses will be conducted for match and training injuries. Interaction effects (e.g., maturation status x trial arm) will be examined for evidence of intervention effect modification.

Secondary outcome measures

1. The influence of coach attitudes on compliance and injury burden between the trial arms, in addition to changes in these through use of the exercise programmes.
2. The study design and both exercise programmes have been formulated in order to optimise compliance within the schools rugby setting. Nevertheless, this aspect will be monitored throughout the study. The coaches responsible for recording the weekly match and training exposure will also be responsible for recording whether or not the squad completed their allocated exercise programme. These data will be combined with exposure data in order to produce an index of compliance for all included teams (such as a percentage of exposures where the exercise programme was completed).
3. The attitudes, knowledge, and beliefs of coaches involved in leading the exercise programmes both before (August 2015) and after (December 2015) the intervention period. This will be undertaken by two paper-based questionnaires of all coaches involved in the study, one prior to study commencement and one shortly before study conclusion using a standardised questionnaire designed by the authors based upon previous work investigating similar features. The questionnaires will assess whether coaches in the different trial arms have differing baseline views on injuries and injury prevention in youth rugby, how these views are associated with compliance to their allocated exercise programme, and whether these views are altered as a result of implementing their allocated exercise programme.
4. The influence of compliance on injury burden will also be analysed. Compliance will be computed into two entities: coach compliance (% of exposures where exercise programme was completed) and player compliance (% of players that completed the exercise programme during each exposure). The product of these two proportions will be taken forward for analysis as the total compliance. Teams in both trial arms will be stratified into tertiles based on total compliance: low, intermediate, and high. Poisson regression will be used to investigate the influence of compliance to the exercise programme on the risk of injury, with the low compliance tertile in the control arm serving as the reference group.
5. Behavioural, attitudes and knowledge variables will be pre-coded and analysed in a number of different ways. Only coaches that complete the study and provide questionnaire responses for at least the pre-season questionnaire will be included in this analysis. These variables will be compared at baseline between the two trial arms and at the end of the intervention to assess any changes using both categorical and continuous data analysis. To analyse the influence of coach beliefs on compliance, linear regression analysis will be used with team compliance (%) as the dependent variable and questionnaire responses serving as the predictor variables. The responses to the questionnaire will also be compared between the compliance tertile groups using categorical data analyses. To investigate the influence of coach attitudes on injury burden, poisson regression, adjusted for clustering by school, will be used with injury burden (days/1000 player-hours) as the dependent variable and questionnaire responses serving as the predictor variables. The results will be presented as Relative Risk (RR) with 95% confidence intervals and p-values. Both sets of analyses will be adjusted for coaching experience (years) and coaching level, which will be recorded as part of the pre-season questionnaire.

Overall study start date

01/01/2015

Completion date

31/12/2015

Eligibility

Key inclusion criteria

1. Males
2. Aged 14-18 years
3. Participating in secondary school rugby union programmes
4. Consenting to participate
5. Participating school has on-site medical facilities

Participant type(s)

Healthy volunteer

Age group

Child

Lower age limit

14 Years

Upper age limit

18 Years

Sex

Male

Target number of participants

Approximately 1920 participants, derived from 32 clusters and 60 participants per cluster

Total final enrolment

3188

Key exclusion criteria

1. Not male 14-18 years
2. Withheld informed consent
3. Participating in a school rugby programme without on-site medical support
4. Participating in a school rugby programme already engaged in specific injury prevention exercise programme

Date of first enrolment

30/03/2015

Date of final enrolment

30/06/2015

Locations**Countries of recruitment**

England

United Kingdom

Study participating centre

University of Bath
Claverton Down
Bath
United Kingdom
BA2 7AY

Sponsor information

Organisation
University of Bath

Sponsor details
Claverton Down
Bath
England
United Kingdom
BA2 7AY

Sponsor type
University/education

Website
<http://www.bath.ac.uk>

ROR
<https://ror.org/002h8g185>

Funder(s)

Funder type
Other

Funder Name
Rugby Football Union

Results and Publications

Publication and dissemination plan
To be confirmed at a later date

Intention to publish date
30/06/2016

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol article	protocol	19/01/2016		Yes	No
Results article		01/08/2017	10/05/2021	Yes	No