# The effect of starting dialysis earlier than usual in patients with damaged kidneys

Submission date	Recruitment status  No longer recruiting	<ul><li>Prospectively registered</li></ul>		
03/02/2019		[X] Protocol		
Registration date 18/02/2019	Overall study status Completed	Statistical analysis plan		
		[X] Results		
<b>Last Edited</b> 07/08/2023	Condition category Urological and Genital Diseases	[] Individual participant data		

#### Plain English summary of protocol

Background and study aims

More than 1 in 2 critically ill patients in the intensive care unit (ICU) develop acute kidney injury (AKI). A proportion of patients need dialysis treatment, also known as replacement therapy (RRT). The aim of RRT is to remove toxins and excess fluid which may have accumulated because of kidney failure. At present it is not clear whether it is best to start RRT only when patients have evidence of severe AKI or whether it is better to initiate RRT earlier, i.e. as soon as patients have signs of impaired kidney function.

#### Who can participate?

Patients aged over 18 years admitted to the intensive care unit with serious kidney problems.

#### What does the study involve?

RRT consists of having a catheter (a tube or 'line') in a vein in the neck or in the groin through which blood flows to a dialysis machine where it is cleansed and excess water is removed. The cleansed blood is then returned to the patient via the catheter.

#### What are the possible benefits and risks of participating?

RRT can be lifesaving but has side effects, possibly including bleeding and bloodstream infections. Importantly, if at any stage the treating clinician feels that RRT is mandatory or alternatively should not be started, their judgement will override. Also, there will be no change in any clinical management and no extra blood samples will be taken.

#### Where is the study run from?

Guy's & St Thomas Hospital, Westminster Bridge Road, London, SE1 9RT, UK.

When is the study starting and how long is it expected to run for? February 2018 to March 2020.

#### Who is funding the study?

The National Institute for Health Research (United Kingdom) Health Technology Assessment programme.

# Contact information

#### Type(s)

Scientific

#### Contact name

Mrs Marlies Ostermann

#### Contact details

Guy's & St Thomas Hospital
Department of Critical Care
Westminster Bridge Road
London
United Kingdom
SE1 9RT
+44 (0)2071883038
Marlies.Ostermann@gstt.nhs.uk

# Additional identifiers

#### Clinical Trials Information System (CTIS)

Nil known

#### Integrated Research Application System (IRAS)

191390

#### ClinicalTrials.gov (NCT)

NCT02568722

#### Protocol serial number

IRAS 191390

# Study information

#### Scientific Title

STandard versus Accelerated initiation of Renal Replacement Therapy in Acute Kidney Injury (STARRT-AKI): a multi-centre, randomized, controlled trial

#### Acronym

STARRT-AKI

#### Study objectives

In critically ill patients with severe acute kidney injury (AKI), randomization to accelerated initiation of renal replacement therapy (RRT), compared to a conservative strategy consistent

with standard care, leads to:

- 1. improved survival at 90 days; and
- 2. better recovery of kidney function, defined as independence from dialysis at 90 days.

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

Approved 22/05/2017, Camberwell St Giles REC (Institute of Psychiatry, Seminar Room 6, IOP Main Building, SE5 8AF, +44 (0) 207 1048044, NRESCommittee.London-CamberwellStGiles@nhs. net), ref:

#### Study design

Randomised controlled trial

#### Primary study design

Interventional

#### Study type(s)

**Treatment** 

#### Health condition(s) or problem(s) studied

Acute kidney injury

#### **Interventions**

Patients in the ICU with early AKI will be randomised to RRT using standard criteria versus accelerated (earlier) RRT. Patients randomised to the accelerated arm will be started on RRT within 12 hours of eligibility. In patients randomised to standard initiation, RRT will be discouraged until serum potassium >5.9 mmol/L, pH <7.21 or serum bicarbonate <13 mmol/L, respiratory failure due to fluid overload, or persistent AKI stage 2 or 3 for >72 hours.

#### Intervention Type

Procedure/Surgery

#### Primary outcome(s)

Mortality measured at 90 days

#### Key secondary outcome(s))

Recovery of kidney function defined as independence from dialysis at 90 days

#### Completion date

31/05/2021

# **Eligibility**

#### Key inclusion criteria

- 1. Age ≥18 years
- 2. Admission to an intensive care unit (ICU)

- 3. Evidence of kidney dysfunction [serum creatinine  $\geq$ 100 µmol/L in women and  $\geq$  130 µmol/L in men]
- 4. Evidence of severe AKI

#### Participant type(s)

**Patient** 

#### Healthy volunteers allowed

No

#### Age group

Adult

#### Lower age limit

18 years

#### Sex

All

#### Total final enrolment

2927

#### Key exclusion criteria

- 1. Lack of commitment to provide RRT as part of limitation of ongoing life support.
- 2. Presence of a drug overdose that necessitates initiation of RRT.
- 3. Any RRT within the previous 2 months.
- 4. Kidney transplant within the past 365 days.
- 5. Known pre-hospitalization advanced chronic kidney disease, defined by an estimated glomerular filtration rate <20 mL/min/1.73 m2 in a patient who is not on chronic dialysis.
- 6. Presence or clinical suspicion of renal obstruction, rapidly progressive glomerulonephritis, vasculitis, thrombotic microangiopathy (eg, thrombotic thrombocytopenic purpura, hemolytic uremic syndrome, malignant hypertension, scleroderma renal crisis) or acute interstitial nephritis.
- 7. Likelihood that an absolute indication for RRT will arise in the subsequent 24 hours based on the most recent blood work for the following parameters: serum K >5.5
- 8. Likelihood that an absolute indication for RRT will arise in the subsequent 24 hours based on the most recent blood work for the following parameters: serum bicarbonate <15 mmol/L

#### Date of first enrolment

01/02/2018

#### Date of final enrolment

31/05/2020

### Locations

#### Countries of recruitment

United Kingdom

England

# Study participating centre Guy's & St Thomas Hospital

Westminster Bridge Road London United Kingdom SE1 9RT

# Sponsor information

#### Organisation

University of Toronto

#### **ROR**

https://ror.org/03dbr7087

# Funder(s)

#### Funder type

Government

#### **Funder Name**

Health Technology Assessment Programme

#### Alternative Name(s)

NIHR Health Technology Assessment Programme, Health Technology Assessment (HTA), HTA

#### **Funding Body Type**

Government organisation

#### **Funding Body Subtype**

National government

#### Location

**United Kingdom** 

## **Results and Publications**

Individual participant data (IPD) sharing plan

# Not provided at time of registration

# IPD sharing plan summary

Data sharing statement to be made available at a later date

# **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		16/07/2020	18/11/2022	Yes	No
Protocol file	version 3.3	15/12/2020	07/08/2023	No	No