

Coronary Artery Bypass graft surgery in patients with asymptomatic carotid stenosis

Submission date 07/09/2010	Recruitment status Stopped	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 25/10/2010	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 26/10/2022	Condition category Circulatory System	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Coronary artery stenosis is a blockage of the arteries that supply blood to the heart muscle. It can be treated with a coronary artery bypass graft (CABG), which is a surgical procedure that diverts blood around the blocked part of the artery, or a carotid endarterectomy (CEA), which is a surgical procedure to unblock the artery. The aim of this study is to compare the safety and effectiveness of CABG alone versus CABG and CEA together in patients with coronary artery stenosis.

Who can participate?

Patients aged over 18 with coronary artery stenosis

What does the study involve?

Participants are randomly allocated to be treated with CABG either with or without CEA. The rates of non-fatal strokes or deaths are measured in both groups within 30 days after the surgery. Participants are followed-up for 1 year (including a physical and an ultrasound examination) with an additional yearly telephone follow-up for up to 5 years.

What are the possible benefits and risks of participating?

Not provided at time of registration

Where is the study run from?

University Hospital Essen (Germany)

When is the study starting and how long is it expected to run for?

September 2010 to December 2019

Who is funding the study?

Deutsche Forschungsgemeinschaft (Germany)

Who is the main contact?

Prof. Christian Weimar

christian.weimar@uk-essen.de

Contact information

Type(s)

Scientific

Contact name

Prof Christian Weimar

Contact details

Department of Neurology

University Hospital Essen

Hufelandstr. 55

Essen

Germany

45122

+49 (0)201 723 6503

christian.weimar@uk-essen.de

Additional identifiers

Protocol serial number

N/A

Study information

Scientific Title

Coronary Artery Bypass graft surgery in patients with Asymptomatic Carotid Stenosis: a randomised controlled open multicentre group sequential trial with two parallel groups and blinded observers

Acronym

CABACS

Study objectives

The objective of this study is to compare the safety and efficacy of isolated coronary artery bypass graft (CABG) versus synchronous CABG and carotid endarterectomy (CEA) in patients with asymptomatic coronary artery stenosis (CAS) greater than or equal to 80% according to the European Carotid Surgery Trial (ECST) criteria.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Institutional Review Board (IRB) of the Medical Faculty of Essen, 05/08/2010, ref: 10-4325

Study design

Randomised controlled open multicentre blinded parallel-group sequential trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Coronary heart disease, carotid artery stenosis

Interventions

CABG with or without CEA must be performed as soon as possible (max. within 7 days) after randomisation. All patients will be treated with up-to-date medication following national and international guidelines. Recommendations for this treatment are provided by the 'Best Medical Treatment' subcommittee. Standards for surgical treatment have been formulated by the Surgical quality sub-committee. Aspirin (or clopidogrel) has to be applied before CABG as well as after surgery. All participating centres of cardiothoracic surgery are obligated to participate in a national quality management register. All participating surgeons have to meet the following standards:

1. Anonymous confirmation of the last 30 consecutively performed CEA, affirmed by the head of department
2. Anonymous confirmation of the last 150 consecutively performed CABG, affirmed by the head of department

Follow-up duration will be 1 year (including physical and ultrasound examination) with an additional yearly telephone follow-up up to 5 years after the intervention.

Intervention Type

Procedure/Surgery

Primary outcome(s)

The primary efficacy and safety endpoint is the event rate of nonfatal strokes or deaths from any cause (whatever occurs first) within 30 days after the intervention (either isolated CABG or synchronous CABG + CEA).

Key secondary outcome(s)

1. Number of ischemic strokes ipsilateral to the initially higher grade, not occluded stenotic carotid artery within 30 days and 1 year
2. Any stroke or vascular death within 30 days, 1 year and 5 years
3. Deaths from any cause within 30 days, 1 year and 5 years
4. Number of disabling strokes (definition: stroke with resulting impairment >3 on the modified Rankin Scale) within 30 days and 1 year
5. Change of cognitive performance on the Demtec scale from randomization to 30 days and 1 year
6. Technical failure of intervention
7. Number of myocardial infarctions within 30 days, and from 30 days to 1 year
8. Duration of ventilatory support after operation (CABG ± CEA)
9. Total length of hospital stay and German diagnosis related group (G-DRG) for acute hospital stay
10. Total length of ICU stay

Completion date

31/12/2019

Eligibility

Key inclusion criteria

Subjects meeting all of the following criteria will be considered for admission to the trial:

1. Asymptomatic (past 180 days) stenosis greater than or equal to 80% (following criteria of the ECST of the extracranial carotid artery in patients scheduled for CABG)
2. Negative pregnancy test in pre-menopausal women
3. Written informed consent and full legal capacity
4. Carotid stenosis treatable with CEA
5. Ability of the patient to participate in follow-up examinations
6. Aged greater than or equal to 18 years, either sex

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

127

Key exclusion criteria

Current exclusion criteria as of 09/04/2013:

Subjects presenting with any of the following criteria will not be included in the trial:

1. Non-atherosclerotic stenosis (e.g. dissection, floating thrombus, fibromuscular dysplasia, tumor)
2. Complete occlusion of the carotid artery to be treated
3. Prior stenting of the carotid artery to be treated
4. Stenosis following radiotherapy
5. Additional higher grade intracranial or intrathoracic stenosis (tandem stenosis)
6. Recent (past 180 days) ischemic symptoms ipsilateral to carotid stenosis or occlusion
7. Contralateral carotid occlusion or other known indication for carotid revascularization (apart from scheduled CABG)
8. NSTEMI within the past 48 hours, STEMI within the past 7 days or hemodynamically unstable patients
9. Evidence for intracranial bleeding within the past 90 days
10. Modified Rankin Scale score >3 or severe aphasia
11. Patients unlikely to survive more than 1 year due to concomitant diseases
12. Planned combined cardiac valve replacement or any other cardiac surgery beyond CABG (+/-

CEA) during the procedure

13. Major surgery (apart from study procedures) planned within 8 weeks from randomization

14. Participation in another clinical trial

Previous exclusion criteria until 09/04/2013:

Subjects presenting with any of the following criteria will not be included in the trial:

1. Non-atherosclerotic stenosis (e.g. dissection, floating thrombus, fibromuscular dysplasia, tumor)
2. Complete occlusion of the carotid artery to be treated
3. Prior stenting of the carotid artery to be treated
4. Stenosis following radiotherapy
5. Additional higher grade intracranial or intrathoracic stenosis (tandem stenosis)
6. Recent (past 180 days) ischemic symptoms ipsilateral to carotid stenosis or occlusion
7. Contralateral carotid occlusion or other known indication for carotid revascularization (apart from scheduled CABG)
8. Myocardial infarction (NSTEMI or STEMI) within the past 7 days or hemodynamically unstable patients
9. Known high risk for cardiogenic embolism requiring anticoagulation (mechanical heart valve, chronic atrial fibrillation, left ventricular thrombus, left ventricular aneurysm)
10. Evidence for intracranial bleeding within the past 90 days
11. Modified Rankin Scale score >3 or severe aphasia
12. Patients unlikely to survive more than 1 year due to concomitant diseases
13. Planned combined cardiac valve replacement or any other cardiac surgery beyond CABG (+/- CEA) during the procedure
14. Major surgery (apart from study procedures) planned within 8 weeks from randomization
15. Participation in another clinical trial

Date of first enrolment

01/12/2010

Date of final enrolment

19/12/2014

Locations

Countries of recruitment

Germany

Study participating centre

University Hospital Essen

Essen

Germany

45122

Sponsor information

Organisation

University Hospital Essen (Universitätsklinikum Essen) (Germany)

ROR

<https://ror.org/02na8dn90>

Funder(s)

Funder type

Research council

Funder Name

Deutsche Forschungsgemeinschaft

Alternative Name(s)

German Research Association, German Research Foundation, Deutsche Forschungsgemeinschaft (DFG), DFG

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

Germany

Results and Publications

Individual participant data (IPD) sharing plan

Not provided at time of registration

IPD sharing plan summary

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	5-year results	09/09/2022	12/09/2022	Yes	No
Results article		15/09/2017	26/10/2022	Yes	No
Protocol article	protocol	01/06/2012		Yes	No
Study website	Study website	11/11/2025	11/11/2025	No	Yes