

# Comparing survival of teeth after restoration using two types of filling material

<b>Submission date</b> 03/02/2019	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 15/02/2019	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 22/02/2019	<b>Condition category</b> Oral Health	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Atraumatic restorative treatment (ART) is an approach to dental treatment that uses only hand-powered instruments. Electricity, anaesthetics and advanced tools are not needed, which means that this approach can be used in developing countries to treat dental cavities, which are otherwise often treated by removing the tooth. ART can also be used in patients who have a dental phobia. However, ART treatment of teeth with complex cavities can have a higher chance of failing than conventional treatment. This trial aims to investigate compare two types of cement material used to fill cavities using the ART approach.

### Who can participate?

Secondary school students aged 15 or 16 years who have at least one premolar or molar tooth with decay that has resulted in a cavity.

### What does the study involve?

Participants will be randomly allocated to one of two groups. Participants in both groups will have their cavities treated using ART. The cavities will be filled with one filler or the other depending on which group they were allocated to. The participants will come back for a dental examination at 1 week and 6, 12, 18 and 24 months after the treatment and the success of the tooth restoration will be assessed each time.

### What are the possible benefits and risks of participating?

The procedure is the same for both groups and might involve some discomfort when the decay is being cleaned out of the cavity. One of the fillers might result in a lower chance of the filling and the tooth remaining sound.

### Where is the study run from?

Cairo University Dental School (Egypt)

### When is the study starting and how long is it expected to run for?

November 2010 to May 2016

Who is funding the study?  
The researcher is funding the study.

Who is the main contact?  
Enas Mobarak, enasmobarak@hotmail.com

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Prof Enas Mobarak

**Contact details**  
Faculty of Oral and Dental Medicine  
Cairo University  
11 El Saraya Street  
Manial  
Cairo  
Egypt  
12311

**Type(s)**  
Scientific

**Contact name**  
Dr Jo Frencken

**ORCID ID**  
<https://orcid.org/0000-0001-9612-1752>

**Contact details**  
Radboud University College of Dental Science  
Nijmegen  
Netherlands  
6525 XZ  
-  
Jo.Frencken@radboudumc.nl

## Additional identifiers

**Clinical Trials Information System (CTIS)**  
Nil known

**ClinicalTrials.gov (NCT)**  
Nil known

**Protocol serial number**  
RHD-IRB0000687-13oct2014-EM02

# Study information

## Scientific Title

Survival of multiple-surface ART restorations using a zinc-reinforced glass-ionomer restorative after 2 years: A randomized triple-blind clinical trial

## Study objectives

The survival of multiple-surface atraumatic restorative treatment (ART) zinc-reinforced high-viscosity glass-ionomer cement (HVGIC) restorations in permanent posterior teeth is significantly higher than that of comparable restorations using conventional HVGIC after 2 years.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Approved 13/10/2014, ethical committee at Ministry of Health and Population, Government Health Insurance in Egypt, ref: RHD-IRB0000687-13Oct2014-EM02

## Study design

Randomised controlled trial

## Primary study design

Interventional

## Study type(s)

Treatment

## Health condition(s) or problem(s) studied

Tooth cavity in more than one surface

## Interventions

Randomisation:

The test intervention was a new restorative material (zinc-containing glass-ionomer cement [ChemFil Rock]). The control intervention was a commonly used glass-ionomer material, FUJI IX GP. First a computer programme randomised a test/control restoration over the four operators and by gender. Using envelopes with marks for gender, the student chose the envelope of his /her gender and presented it to a third person who, using a randomised list of restorative materials, informed the operator which restorative was to be used. The intervention was masked for operator, evaluation (using replicas) and statistician (test-control group only known by chief investigator).

Intervention:

Tooth cavities were prepared entirely using hand instruments. Soft carious dentine was excavated using ART hand excavators. Then the cavity was wiped with water-soaked cotton pellets, dried using dry cotton pellets and cautiously inspected for the presence of soft dentine, which was then removed when necessary.

In the control arm, a dentine conditioner was applied to the floor and the walls of the prepared cavity for 10 s. The conditioner was rinsed out using a cotton pellet soaked in water until no remnant of the conditioner was visible (about 10 s). Then, the cavity was blotted using dry cotton pellets (5 s). This procedure was not performed in the test arm. A prefabricated matrix

band and a wedge were inserted. A high-viscosity glass-ionomer cement (HVGIC) was the filling material. In the test group, HVGIC Chemfil Rock was inserted in the tooth cavity and in the control arm, HVGIC Fuji IX GP was used. Excess HVGIC was removed with a large excavator and ART contouring instrument and left to harden for 25-30 s. The wooden wedge and matrix band were then removed and the height of the filling was checked using an articulating paper. Participants were instructed neither to eat nor to brush the restored side of the mouth for at least 2 h, and to brush the teeth twice daily with a fluoride containing toothpaste every day thereafter. The participant was also advised to contact the operator in case of having any dental problems or pain.

#### Evaluation:

Using intra-oral photographic mirrors, digital clinical photographs of the occlusal and approximal surfaces were taken preoperatively after cavity preparation and postoperatively at baseline (1 week post-restoration) and at each of the four follow-up time points (6, 12, 18 and 24 months) using a digital camera Nikon D40 and a Macro lens supplied with a ring flash.

A replica of each restored tooth was produced at each of the evaluation points. Two independent evaluators assessed the restorations using ART restoration assessment criteria and those of the FDI (World Dental Federation). In case of disagreement, the case was discussed and consensus reached. Restoration survival curves were estimated and difference between variables were tested with a Wald (chi-square) test.

#### Intervention Type

Procedure/Surgery

#### Primary outcome(s)

Survival of multiple-surface ART restorations after 2 years

#### Key secondary outcome(s)

Main reason for failure of restorations after 2 years

#### Completion date

31/05/2016

## Eligibility

#### Key inclusion criteria

1. Secondary school students
2. Healthy patients without any history of any medical condition that could interfere with the study protocol or affect clinical results
3. Good oral hygiene (Simplified Oral Hygiene Index (S-OHI)  $\leq 0.5$ )
4. Caries prevalence (DMFS)  $< 12$
5. Without adverse oral habits that could affect the study results
6. Presence of natural antagonist tooth that is not restored
7. Presence of at least one tooth with a cavitated or dentine carious lesions including at least occlusal-proximal (multi-surface) surface in first or second permanent premolars/molars situated in any side of the jaw
8. Presence of adjacent tooth to the cavity and being fully erupted, normally positioned (not tilted or rotated) and having natural (not restored) contact
9. Occluso-proximal cavities with size (Site/Stage 2.2, 2.3 and 2.4)
10. Absence of apparent enamel crack or fracture
11. No pulp involvement or symptoms of pulpitis or apical periodontitis

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Child

**Sex**

All

**Total final enrolment**

218

**Key exclusion criteria**

1. Teeth without antagonist teeth or with a prosthetic antagonist
2. Occlusal-proximal cavities limited to enamel layer
3. Deciduous teeth
4. Abnormal oral habits with excessive occlusal function or parafunction, such as pencil chewing or bruxism
5. Daily consumption of citrus juices

**Date of first enrolment**

01/03/2012

**Date of final enrolment**

01/05/2014

**Locations****Countries of recruitment**

Egypt

Netherlands

**Study participating centre**

**Cairo University Dental School**

14 El Ansar Street

Dokki

Cairo

Egypt

12311

**Study participating centre**

**Radboud University Medical Centre**  
P.O. Box 9101  
Nijmegen  
Netherlands  
6500 HB

## Sponsor information

### Organisation

Dental School, University of Cairo

### ROR

<https://ror.org/03q21mh05>

## Funder(s)

### Funder type

Other

### Funder Name

Investigator initiated and funded

## Results and Publications

### Individual participant data (IPD) sharing plan

The data were stored at the Radboud University Medical Centre, Geert Grooteplein Zuid 10, 6525 GA Nijmegen, The Netherlands. The availability of the data is unknown and there is no data sharing plan.

### IPD sharing plan summary

Stored in repository