

Cold compared with room temperature compress on the perineal repair site following normal vaginal delivery in women who have delivered their first child

Submission date 16/02/2022	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 05/04/2022	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 28/12/2023	Condition category Pregnancy and Childbirth	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Vaginal delivery offers many benefits such as a rapid recovery process and an early start to the mother-infant relationship especially in primiparous (first birth) women when it is unfortunately often associated with perineal trauma (any damage to the area between the vagina and the anus during the birth of a baby). Cryotherapy (ice pack application) is a non-pharmacological, non-invasive, low-cost therapy, which reduces local tissue temperature. This localised method results in an anti-inflammatory effect that consequently leads to less swelling and pain.

The objective of this proposed study is to evaluate cold versus room temperature compress (placebo) to the perineal repair site on perineal pain in primiparous women.

Who can participate?

All women aged 18 years old and above, in their first pregnancy and planning on vaginal delivery is eligible to participate in this study. The final inclusion criteria will be determined upon completion of perineal repair.

What does the study involve?

The study involved applying either cold or room temperature compress at the perineal repair site at 3 different points of time: immediately post repair, at 4-hour and 8-hour post delivery. The compress will be placed at the perineum for 20minutes each before it is being discarded.

What are the possible benefits and risks of participating?

The purpose of the trial is to evaluate the trial interventions as there is uncertainty about which is better or they may be equivalent. Information obtained from this study will help guide the management of future patients in similar circumstances. Some women may feel the compress to be uncomfortable. As the compress will be placed at the perineum for 20minutes, patient will have difficulty in sitting or walking but they will be provided with proper disposable panties that can support the compress and make them more comfortable without any extra cost. There is no

evidence to show that applying either of this compress will increase risk of perineal wound breakdown.

Where is the study run from?

Department of Obstetrics and Gynaecology, University Malaya Medical Center (Malaysia)

When is the study starting and how long is it expected to run for?

November 2021 to March 2023

Who is funding the study?

Department of Obstetrics and Gynaecology, University Malaya Medical Center (Malaysia)

Who is the main contact?

Dr. Khaliza Kazrin binti Abd Karim, azaareen84@gmail.com

Prof PC Tan, tanpengchiong@yahoo.com

Contact information

Type(s)

Principal investigator

Contact name

Dr Khaliza Kazrin Abd Karim

ORCID ID

<https://orcid.org/0000-0001-8713-6581>

Contact details

University Malaya Medical centre

Jalan Professor Ungku Aziz

Kuala Lumpur

Malaysia

59100

+60 122710429

azaareen84@gmail.com

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

MECID number 2021118-10761

Study information

Scientific Title

Cold compared with room temperature compress on the perineal repair site following normal vaginal delivery in primiparous women

Acronym

CROCIP

Study objectives

We hypothesized that women randomised to cold compress compared to controls will have

- lower pain score when mobilizing post-delivery
- lower requirement of oral analgesia
- earlier resumption of vagina intercourse

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 10/02/2022, Medical Research Ethics Committee of University Malaya Medical Centre (Level 2, Kompleks Pendidikan Sains Kejururawatan, University of Malaya Medical Centre Jalan Professor Ungku Aziz 59100 Kuala Lumpur, Malaysia; +603-7949 3209/2251; ummc-mrec@ummc.edu.my), ref: 2021118-10761

Study design

Interventional single-centre randomized controlled trial

Primary study design

Interventional

Study type(s)

Quality of life

Health condition(s) or problem(s) studied

Supportive care for perineal wound sustained in primiparous women during normal vaginal delivery.

Interventions

Patient who are agreed to participate in this study, will be randomised to:

1. Cold compress (frozen gel pack) - experimental group, or
2. Room temperature compress (using identical gel pack, soft at room temperature) - control group

The allocated pack will be placed against the sutured site of the perineum for 20 min at 3 different time points: immediately after repair, at 4 h after repair, and 8 h after repair.

Analgesia:

Stat dose of 1 g paracetamol po after perineal repair then 1 g paracetamol po qds prn. If breakthrough pain, escalation to 400 mg ibuprofen po qds. Further escalation at care providers' discretion

At discharge 1 g paracetamol po qds prn or if moderate pain, 400 mg ibuprofen po qds prn with 3 days' supply. Further escalation at care providers' discretion

Randomization:

Randomization sequence generated in random blocks of 4 or 8 with within block randomization using <https://www.sealedenvelope.com/simple-randomiser/v1/lists> by an investigator not involved in recruitment. The randomization sequence is concealed in numbered, sealed opaque envelopes. Randomization is by opening the lowest-numbered envelope remaining for the newest recruit.

Intervention Type

Procedure/Surgery

Primary outcome(s)

Perineal pain score measured using a 0-10 numerical rating scale (NRS) at mobilisation, at 12 h after delivery, and 24 h after delivery

Key secondary outcome(s)

Measured at pre-discharge using patient records:

1. Time to first flatus
2. Time to first satisfactory breastfeeding experience
3. Time to first urination
4. Time to ambulation
5. VNRS: Satisfaction with allocated compress to your perineal repair site (@24 H)
6. Analgesia used

Measured at 14 days after delivery (by telephone):

7. Oral analgesia use for perineal pain
8. VNRS pain score during mobilisation

Measured at 6 weeks after delivery (by telephone):

9. Oral analgesia use for perineal pain
10. VNRS pain score during mobilisation
11. Resumption of vagina intercourse
12. Exclusive breastfeeding
13. Perineal healing (Likert scale)

Completion date

31/03/2023

Eligibility

Key inclusion criteria

1. Nulliparous
2. Planning to have a normal birth
3. ≥ 18 years old
4. ≥ 37 weeks gestation
5. Singleton birth
6. Spontaneous vaginal delivery (SVD)
7. Sutured second degree perineal injury: spontaneous or episiotomy
8. Apgar score at 5 minutes ≥ 9

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Female

Total final enrolment

224

Key exclusion criteria

1. Post-partum haemorrhage (≥ 1000 ml)
2. Indwelling bladder catheter
3. Admission to high dependency or intensive care
4. Extended or multiple vaginal tears
5. Vulva-vaginal haematoma
6. Cold allergy

Date of first enrolment

09/05/2022

Date of final enrolment

07/02/2023

Locations**Countries of recruitment**

Malaysia

Study participating centre

University Malaya Medical Centre

Jalan Prof. DiRaja Ungku Aziz

Kuala Lumpur

Malaysia

59100

Sponsor information**Organisation**

University Malaya Medical Center

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

University Malaya Medical Center

Results and Publications

Individual participant data (IPD) sharing plan

Data sharing statement to be made available at a later date

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		24/12/2023	28/12/2023	Yes	No
Participant information sheet		29/01/2021	21/02/2022	No	Yes