

Does the filling state of the seminal vesicles influence sexual desire?

Submission date 03/04/2014	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 09/05/2014	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 24/07/2020	Condition category Urological and Genital Diseases	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

The seminal vesicles are part of the male inner genitals. It is not known if the seminal vesicles have an impact on sexual interest, i.e. if full seminal vesicles increase sexual interest. This would potentially make sense from a biological/behavioural point of view. The aim of this study is to find out if there is any specific brain area that changes its activity in relation to the filling status of the seminal vesicles. The detection of such a specific brain area would suggest that the filling status of the seminal vesicles changes the activity in the brain.

Who can participate?

Heterosexual men between the ages of 20 and 30

What does the study involve?

On day 1 in the morning, participants provide a blood sample for analysis of sex hormone levels, a conventional MRI scan of the pelvis to assess the size of the seminal vesicles, and a functional MRI of the brain. After these examinations, they masturbate in order to empty their seminal vesicles. On day 2 in the morning, they undergo the same examinations.

What are the possible benefits and risks of participating?

The detection of specific brain areas would potentially have a clinical impact in men undergoing radical prostatectomy for prostate cancer, which nowadays is usually performed with removal of both seminal vesicles. If a relationship is found between seminal vesicle filling status, specific brain areas and sexual interest and confirmed by further clinical studies, the standard removal of the seminal vesicles during radical prostatectomy could be re-thought in order to potentially help to preserve sexual interest. There are no risks involved in this study.

Where is the study run from?

University Hospital Bern (Switzerland)

When is the study starting and how long is it expected to run for?

May 2014 to April 2015

Who is funding the study?
Max and Hedwig Niedermaier Foundation and University Hospital Bern (Switzerland)

Who is the main contact?
Dr Frédéric Birkhäuser
frederic.birkhaeuser@insel.ch

Contact information

Type(s)
Scientific

Contact name
Dr Frédéric Birkhäuser

Contact details
Freiburgstrasse 4
Bern
Switzerland
3010
-
frederic.birkhaeuser@insel.ch

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
N/A

Study information

Scientific Title
Does the filling state of the seminal vesicles influence sexual desire? A pilot study

Study objectives
Filling state of seminal vesicles is an independent factor for the intensity of sexual desire, influencing sexual arousal in healthy men.

Ethics approval required
Old ethics approval format

Ethics approval(s)
Kantonale Ethikkommission Bern (KEK), 05/09/2014, ref: 005/14

Study design

Interventional study

Primary study design

Interventional

Secondary study design

Non randomised study

Study setting(s)

Hospital

Study type(s)

Other

Participant information sheet

Not available in web format, please use the contact details to request a patient information sheet

Health condition(s) or problem(s) studied

The physiological role of healthy seminal vesicles on sexual desire

Interventions

The five participants all undergo the same intervention. On day 1 in the morning, they first undergo a blood analysis (5 min), thereafter they undergo a conventional MRI of the pelvis and a functional MRI of the brain (40 min). After these examinations, they masturbate in order to empty their seminal vesicles. On day 2 in the morning, they undergo exactly the same examinations.

Intervention Type

Other

Phase

Not Applicable

Primary outcome measure

The effect of seminal vesicle filling state on cortical activation and sexual arousal investigated by fMRI, which would be reflected by a reduced activation of specific regions in fMRI of the brain on study after ejaculation.

Secondary outcome measures

1. Changes in sexual arousal score with full/empty seminal vesicles
2. Changes of volume in seminal vesicles during first (before ejaculation) and second (after ejaculation) MRI examination

Overall study start date

01/05/2014

Completion date

30/04/2015

Eligibility

Key inclusion criteria

1. Men between the age of 20 and 30
2. Heterosexuality
3. Written informed consent
4. Ejaculation abstinence of 3-5 days before first fMRI examination

Participant type(s)

Patient

Age group

Adult

Sex

Male

Target number of participants

5

Key exclusion criteria

1. Erectile dysfunction
2. Claustrophobia
3. Depression
4. Status post brain trauma
5. Severely reduced visibility
6. Medication intake with possible side effects on libido

Date of first enrolment

01/05/2014

Date of final enrolment

30/04/2015

Locations

Countries of recruitment

Switzerland

Study participating centre

University Hospital Bern

Bern

Switzerland

3010

Sponsor information

Organisation

University Hospital Bern (Switzerland)

Sponsor details

Freiburgstrasse 4

Bern

Switzerland

3010

-

urology@insel.ch

Sponsor type

Hospital/treatment centre

Website

<http://www.insel.ch/en/>

ROR

<https://ror.org/01q9sj412>

Funder(s)**Funder type**

Charity

Funder Name

Max and Hedwig Niedermaier Foundation (Switzerland)

Funder Name

University Hospital Bern (Switzerland)

Results and Publications**Publication and dissemination plan**

Not provided at time of registration

Intention to publish date**Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration