

Evaluation of early and combined treatment with surfactant and inhaled nitric oxide in newborns with respiratory failure and pulmonary hypertension

Submission date 03/09/2019	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 06/09/2019	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 26/08/2020	Condition category Neonatal Diseases	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Hypoxemic respiratory failure (HRF) is a leading cause of illness and death among term and near term newborns. Few studies have tested surfactant treatment in these patients with conflicting results. The aim of this study is to test whether early use of surfactant in combination with inhaled nitric oxide (iNO) will prevent newborns with moderate hypoxemic respiratory failure from developing severe HRF.

Who can participate?

Term and near-term newborns (35 weeks gestational ages and over) with acute HRF

What does the study involve?

Participants are randomly allocated to one of two groups. The surfactant-iNO group receive the combined treatment of surfactant plus inhaled Nitric Oxide (iNO). The control group receive standard treatment with iNO. Infants of both groups receive standard intensive care, which includes high frequency or conventional mechanical ventilation, inotropic support and sedation. ECMO was considered in patients with a persistent OI>40.

What are the possible benefits and risks of participating?

It is expected that the combined treatment of inhaled Nitric Oxide and surfactant will help to improve oxygenation and prevent the progression to severe HRF. This may reduce mortality and the need for ECMO treatment. Surfactant is a usual therapy for preterms with respiratory distress, occasionally its administration is associated with oxygen desaturation and bleeding, but the incidence of these complications is minimized with careful application and monitoring.

Where is the study run from?

Five level III Neonatal Intensive Care Units in five hospitals in Chile

When is the study starting and how long is it expected to run for?
August 2008 to July 2015

Who is funding the study?
Chilean National Fund for Research in Health (FONIS)

Who is the main contact?
Dr Alvaro González
alvgonza@med.puc.cl

Contact information

Type(s)
Scientific

Contact name
Dr Alvaro González

ORCID ID
<https://orcid.org/0000-0002-0442-9109>

Contact details
Diagonal Paraguay 362, 8° piso
Santiago
Chile
8330077
+56 (0)223543887
alvgonza@med.puc.cl

Additional identifiers

Protocol serial number
FONIS SA07120035

Study information

Scientific Title
Randomized controlled trial to assess the early addition of exogenous surfactant to inhaled nitric oxide in the treatment of newborns with respiratory failure and pulmonary hypertension

Acronym
Surfactant plus Nitric Oxide in Neonatal Hypoxic Respiratory Failure

Study objectives
The early addition of exogenous surfactant to the treatment with inhaled Nitric Oxide of newborns with respiratory failure (OI>20) and pulmonary hypertension will improve oxygenation and prevent them from developing severe hypoxemic respiratory failure (OI > 40).

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 08/07/2007, Research Ethics Committee of Medicine School Pontificia Universidad Catolica de Chile (Comité Etico Científico Facultad de Medicina, Pontificia Universidad Católica de Chile, Diagonal Paraguay 383, Torre 11, Local 4, Santiago, Chile 8330077; Tel: +56 (0)2 23548173; Email: cecmeduc@med.puc.cl), CE# 027807

Study design

Prospective randomized double-blind placebo-controlled clinical trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Neonatal hypoxic respiratory failure and pulmonary hypertension

Interventions

Enrolled infants were randomized into one of the two study groups using sequential sealed opaque envelopes from a computer-generated randomization list. This allocation sequence was generated centrally by an independent statistician, and sequentially sealed envelopes were sent to each center.

Surfactant-iNO group: received treatment with 20 ppm of inhaled Nitric Oxide (INOmax, INO Therapeutics, USA) with up to 2 doses of 100 mg/kg of poractant alfa (Curocerf; Chiesi Farmaceutici, Italy)

Control group: received treatment with 20 ppm iNO + placebo (air) and standard intensive care, which in both groups includes high frequency or conventional mechanical ventilation with 100% FiO₂, inotropic support and sedation

Infants in both groups could receive open-label surfactant if they developed an OI>40 within the first 48 hrs of study. ECMO was considered in patients with a persistent OI>40.

Study surfactant combined with iNitric Oxide treatment were used within first 48 hours after study entry, but infants continue receiving standard intensive care later according to local and physician in charge guidelines. Patients were strictly monitored and followed during their whole hospital stay until discharge home.

Intervention Type

Mixed

Primary outcome(s)

The proportion of infants developing treatment failure during the first 48 hours of study treatment, defined as an oxygenation index (OI) \geq 40

Key secondary outcome(s)

1. Proportion of infants requiring ECMO or death recorded during hospital stay
2. Oxygen and ventilatory requirements recorded during hospital stay

Completion date

31/07/2015

Eligibility

Key inclusion criteria

1. Near term infants (35 weeks gestation) with birth weights > 2000 g and who are less than 72 hours old
2. Infants requiring mechanical ventilation with hypoxemic respiratory failure with an OI > 20, calculated on two consecutive measurements of post ductal arterial blood gases without acidosis (pH > 7.25) and efforts will be done to adjust ventilatory support to reach a PCO₂ < 60
3. A previous echocardiogram will be performed to rule out a cardiac anomaly and evaluate the presence of pulmonary hypertension (a tricuspid insufficiency jet with an estimated systolic pulmonary artery pressure \geq 2/3 of systemic systolic arterial blood pressure and/or evidence of right-to-left shunting through foramen ovale or ductus arteriosus)

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Neonate

Sex

All

Total final enrolment

100

Key exclusion criteria

1. Life-threatening congenital anomalies
2. Congenital heart disease
3. Congenital diaphragmatic hernia
4. Other forms of lung hypoplasia syndromes
5. Suspected or confirmed chromosomal abnormality
6. Postnatal age > 72 hours

Date of first enrolment

01/03/2009

Date of final enrolment

30/04/2014

Locations

Countries of recruitment

Chile

Study participating centre

Hospital Clinico Pontificia Universidad Catolica de Chile

Marcoleta 367

Santiago

Chile

8330077

Study participating centre

Hospital Regional Guillermo Grant Benavente, Concepción

San Martin 1436

Concepción

Chile

4070038

Study participating centre

Hospital Dr Luis Tisné

Avenida Las Torres 5150

Santiago

Chile

8640002

Study participating centre

Hospital Dr Sóteo del Río

Avenida Concha y Toro 3459

Puente Alto

Santiago

Chile

8207257

Study participating centre

Hospital San José

San José 1196

Independencia

Santiago

Chile

8380419

Sponsor information

Organisation

Pontificia Universidad Católica de Chile

ROR

<https://ror.org/04teye511>

Funder(s)

Funder type

Government

Funder Name

Chilean Fund for Health Research (FONIS): SA07I20035

Results and Publications

Individual participant data (IPD) sharing plan

Deidentified individual participant data will be made available, in addition to study protocols, the statistical analysis plan, and the informed consent form. The data will be made available upon publication to researchers who provide a methodologically sound proposal for use in achieving the goals of the approved proposal. Proposals should be submitted to alvgonza@med.puc.cl

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	13/08/2020	26/08/2020	Yes	No