# Exposure to nicotine and tobacco products during pregnancy

Submission date	Recruitment status  No longer recruiting	<ul><li>Prospectively registered</li></ul>		
25/11/2019		Protocol		
Registration date	Overall study status Completed Condition category	Statistical analysis plan		
27/11/2019		Results		
Last Edited		Individual participant data		
17/01/2025	Pregnancy and Childbirth	[X] Record updated in last year		

#### Plain English summary of protocol

Background and study aims

Health professionals and pregnant women are cautious about recommending or using e-cigarettes (ECs) in pregnancy, due to lack of detailed safety information. The main aim of the study is to measure toxins and cancer causing substances in the urine of women who used to smoke and now just use ECs compared with women who just smoke.

#### Who can participate?

Pregnant women who are smokers, users of ECs, nicotine replacement therapy, smoke and use ECs, or have never smoked.

#### What does the study involve?

Pregnant women will be invited to take part when they attend a routine antenatal appointment. If they agree to participate, they will be asked by a researcher to provide a urine sample (or a routinely collected sample will be used if available) and to complete a questionnaire (about themselves and use of nicotine/tobacco products, alcohol, cannabis). They will also be asked to blow into a device that measures exposure to tobacco smoke. They will then be asked to meet with the researcher again at a further routine antenatal visit, in order to repeat the assessments.

What are the possible benefits and risks of participating? None

#### Where is the study run from?

- 1. St George's University Hospitals NHS Foundation Trust, UK
- 2. Lancashire teaching Hospitals NHS Foundation Trust, UK
- 3. Bradford Teaching Hospital NHS Foundation Trust, UK
- 4. The Newcastle Upon Tyne Hospitals NHS Foundation Trust, UK
- 5. Gateshead Health NHS Foundation Trust, UK

When is the study starting and how long is it expected to run for? November 2019 to October 2021 Who is funding the study? Cancer Research UK

Who is the main contact? Prof. Michael Ussher mussher@squl.ac.uk

#### Contact information

#### Type(s)

Scientific

#### Contact name

Prof Michael Ussher

#### **ORCID ID**

https://orcid.org/0000-0002-0995-7955

#### Contact details

Population Health Research Insititute St George's University of London Cranmer Terrace London United Kingdom SW17 ORE +44 (0)20 8725 5605 mussher@squl.ac.uk

#### Additional identifiers

#### Clinical Trials Information System (CTIS)

Nil known

#### Integrated Research Application System (IRAS)

269631

#### ClinicalTrials.gov (NCT)

Nil known

#### Protocol serial number

V4.0; IRAS project ID: 269631

## Study information

#### Scientific Title

Maternal exposure to carcinogens, toxicants and nicotine during pregnancy among e-cigarette users, smokers, nicotine replacement therapy users and among those who have never used nicotine or tobacco products

#### **Study objectives**

During pregnancy, we hypothesise that eletronic cigarette (EC) users will have significantly lower levels of urinary acrolein than smokers. As secondary objectives we will explore levels of a range of biomarkers in exclusive EC users compared with exclusive smokers, dual users of cigarettes and ECs, exclusive nicotine replacement therapy (NRT) users; dual users of cigarettes and NRT, and 'never users of tobacco or nicotine products'. Generally, the hypothesis is that less tobacco smoke exposure will equate to less toxin and carcinogen exposure.

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

Approved 29/10/2019, London - Brighton & Sussex Research Ethics Committee (Health Research Authority, Skipton House, 80 London Road, London, SE1 6LH; +44(0)207 104 8052; NRESCommittee.SECoast-BrightonandSussex@nhs.net), ref: 19/LO/1599

#### Study design

Observational cross-sectional multicentre study

#### Primary study design

Observational

#### Study type(s)

Other

#### Health condition(s) or problem(s) studied

Use of nicotine and tobacco products during pregnancy

#### **Interventions**

Pregnant women will be invited to take part when they attend a routine antenatal appointment. If they agree to participate, they will be asked by a researcher to provide a urine sample (or a routinely collected sample can be used if available) and to complete a questionnaire (about themselves and use of nicotine/tobacco products, alcohol, cannabis). They will also be asked to blow into a device that measures exposure to tobacco smoke. They will then be asked to meet with the researcher again at a further routine antenatal visit, in order to repeat the assessments. Urine samples will be stored in a freezer and at the end of the study will be sent to an external laboratory for testing. The results will provide valuable information for women who are pregnant, for healthcare professionals who care for those women and for regulators, about the potential safety of using ECs during pregnancy.

#### Intervention Type

Other

#### Primary outcome(s)

Current primary outcome measure as of 13/12/2021:

Maternal urinary levels of the volatile organic compound Acrolein (metabolite HPMA) measured using urine test at the time of participation. Acrolein has two metabolites (3HPMA & 2CoEMA); the single primary outcome will be 3HPMA because 3HPMA has a higher ratio in smokers vs non-smokers compared with 2CoEMA. The metabolite 2CoEMA will be analysed as a secondary outcome.

Previous primary outcome measure:

Maternal urinary levels of the volatile organic compound Acrolein (metabolites HPMA and CEMA) measured using urine test at the time of participation

#### Key secondary outcome(s))

Current secondary outcome measures as of 13/12/2021:

All tested in maternal urine at time of participation:

- 1. Tobacco exposure will be assessed with a test for minor tobacco alkaloids (anabasine, anatabine), expired carbon monoxide levels
- 2. Levels of nicotine, cotinine and their metabolites, to derive total nicotine equivalents
- 3. Nicotine metabolite ratio (NMR)
- 4. Two tobacco-specific nitrosamines (TSNAs) (i.e., 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNAL) and N-nitrosonornicotine (NNN)), two heavy metals (i.e., cadmium (Cd) and lead (Pb)) and one polycyclic aromatic hydrocarbon (PAH) (i.e., 2-naphthol)
- 5. 13 volatile organic compounds (VOCs) parent (metabolite): acrolein (2CoEMA), 1,3-butadiene (MHB3), acrylamide (AAMA), acrylonitrile (CYMA), Benzene (PMA), crotonaldehyde (HPMM), cyanide (ATCA), isoprene (IPM3), ethylbenzene (PHGA), propylene oxide (HPM2), styrene (MADA), toluene/benzyl alcohol (BMA), and m,p-xylene (34MH)
- 6. Human chorionic gonadotropin (hCG)
- 7. Pregnancy-associated plasma protein A (PAPPA-A)
- 8. Use of alcohol and cannabinoids

Previous secondary outcome measures:

All tested in maternal urine at time of participation:

- 1. Tobacco exposure will be assessed with a test for minor tobacco alkaloids (anabasine, anatabine), expired carbon monoxide levels
- 2. Levels of nicotine, cotinine and their metabolites, to derive total nicotine equivalents
- 3. Nicotine metabolite ratio (NMR)
- 4. Two tobacco-specific nitrosamines (TSNAs) (i.e., 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNAL) and N-nitrosonornicotine (NNN)), two heavy metals (i.e., cadmium (Cd) and lead (Pb)) and one polycyclic aromatic hydrocarbon (PAH) (i.e., 2-naphthol)
- 5. 12 volatile organic compounds (VOCs) parent (metabolite): 1,3-butadiene (MHB3), acrylamide (AAMA), acrylonitrile (CYMA), Benzene (PMA), crotonaldehyde (HPMM), cyanide (ATCA), isoprene (IPM3), ethylbenzene (PHGA), propylene oxide (HPM2), styrene (MADA), toluene /benzyl alcohol (BMA), and m,p-xylene (34MH)
- 6. Human chorionic gonadotropin (hCG)
- 7. Pregnancy-associated plasma protein A (PAPPA-A)
- 8. Use of alcohol and cannabinoids

#### Completion date

31/10/2021

# **Eligibility**

#### Key inclusion criteria

- 1. Pregnant women
- 2. Aged at least 16 years of age
- 3. Able to speak and read English
- 4. Do not report having had an active infection (chest infection, cold or flu, sore throat, or fever) within 24 hours of either of the two assessments
- 5. Willing and able to give informed consent for participation in the study

- 6. Meet criteria for one of six groups related to level and type of use of nicotine and tobacco products:
- 6.1. Group A. 'Exclusive smokers': Smoked more than 5 cigarettes every day for more than the last six months and have not vaped or used NRT in the last month
- 6.2. Group B. 'ECs only': Not smoked cigarettes for at least a month and have vaped every day for at least the last two weeks. Also, have not used NRT in the last month
- 6.3. Group C. 'NRT only': Not smoked cigarettes for at least a month and have used NRT every day for at least the last two weeks. Also, have not vaped in the last month
- 6.4. Group D. Dual users of cigarettes and ECs: Smoked cigarettes on most days of the week for more than 6 months. Also, have been using e-cigarettes on most days of the week for at least the last two weeks
- 6.5. Group E. Dual users of cigarettes and NRT: Smoked cigarettes on most days of the week for more than 6 months. Also, have been using NRT on most days for at least the last two weeks 6.6. Group F. 'never smokers': Have never smoked or used any nicotine or tobacco products

#### Participant type(s)

Patient

#### Healthy volunteers allowed

No

#### Age group

Mixed

#### Lower age limit

16 years

#### Sex

Female

#### Total final enrolment

149

#### Key exclusion criteria

Current participant exclusion criteria as of 20/10/2021:

- 1. Have used tobacco products other than cigarettes in the last month (e.g., cigars, heat-not-burn)
- 2. Have used both NRT and ECs in the last month
- 3. Have used nicotine free e-cigarettes in the last month
- 4. Report having had an active infection (chest infection, cold or flu, sore throat, or fever) within 24 hours of either of the two assessments
- 5. Reported as not currently smoking with an expired carbon monoxide reading of >3 parts per million (where an expired carbon monoxide reading has been obtained)

#### Previous participant exclusion criteria:

- 1. Have used tobacco products other than cigarettes in the last month (e.g., cigars, heat-not-burn)
- 2. Have used both NRT and ECs in the last month
- 3. Have used nicotine free e-cigarettes in the last month
- 4. Report having had an active infection (chest infection, cold or flu, sore throat, or fever) within 24 hours of either of the two assessments

# Date of first enrolment 14/11/2019

# Date of final enrolment 30/04/2020

#### Locations

#### Countries of recruitment

United Kingdom

England

SW17 0QT

Study participating centre
St George's University Hospitals NHS Foundation Trust
Blackshaw Road
London
United Kingdom

# Study participating centre Lancashire teaching Hospitals NHS Foundation Trust Preston Poyal Hospital

Preston Royal Hospital Sharoe Green Lane Preston United Kingdom PR2 9HT

#### Study participating centre Bradford Teaching Hospital NHS Foundation Trust

Bradford Royal Infirmary Smith Lane Bradford United Kingdom BD9 6RJ

# Study participating centre The Newcastle Upon Tyne Hospitals NHS Foundation Trust The Royal Victoria Infirmary Queen Victoria Road

Newcastle upon Tyne United Kingdom NE1 4LP

# Study participating centre Gateshead Health NHS Foundation Trust

Queen Elizabeth Hospital Queen Elizabeth Avenue Sheriff Hill Gateshead United Kingdom NE9 6SX

## Sponsor information

#### Organisation

St George's University Hospitals NHS Foundation Trust

#### **ROR**

https://ror.org/039zedc16

# Funder(s)

#### Funder type

Charity

#### **Funder Name**

Cancer Research UK

#### Alternative Name(s)

CR\_UK, Cancer Research UK - London, Cancer Research UK (CRUK), CRUK

#### **Funding Body Type**

Private sector organisation

#### **Funding Body Subtype**

Other non-profit organizations

#### Location

**United Kingdom** 

## **Results and Publications**

#### Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are not expected to be made available as consent has not be obtained for this purpose.

#### IPD sharing plan summary

Not expected to be made available

#### **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			28/06/2023	No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Plain English results			17/01/2025	No	Yes