

The addition of ACT or a talking control to treatment as usual for the management of dysfunction in advanced cancer

Submission date 22/07/2015	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 22/07/2015	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 18/08/2023	Condition category Cancer	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

<http://www.cancerresearchuk.org/about-cancer/find-a-clinical-trial/a-study-comparing-2-talking-therapies-for-people-having-treatment-for-cancer-symptoms-canact>

Contact information

Type(s)

Public

Contact name

Dr Marc Serfaty

Contact details

University College London
Department of Mental Health Sciences
Rowland Hill Street
London
United Kingdom
NW3 2PF

Additional identifiers

Protocol serial number

18493

Study information

Scientific Title

The addition of ACT or a Talking Control to treatment as usual for the management of dysfunction in advanced cancer: a feasibility randomised controlled trial

Acronym

CanACT

Study objectives

Aim:

To conduct a feasibility randomised controlled trial of the clinical and cost effectiveness of ACT compared to a talking control (TC) or treatment as usual (TAU) in the management of dysfunction in advanced cancer.

Objectives:

1. To test the feasibility of recruitment to/and attrition in people with advanced cancer into a randomised controlled trial of TAU plus ACT compared to TAU plus TC
2. To explore the feasibility of providing a therapist delivered intervention; individual ACT or TC
3. To assess the usefulness and acceptability of a number of clinical and economic outcomes
4. To determine whether data generated from outcomes in this trial support a larger RCT into the clinical effectiveness of ACT in advanced cancer patients
5. To obtain qualitative data about the experience of receiving ACT and TC

Ethics approval required

Old ethics approval format

Ethics approval(s)

NRES committee London – Riverside, 04/12/2014, ref: 14/LO/0813

Study design

Randomized; Interventional; Design type: Treatment

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Topic: Cancer; Subtopic: All Cancers/Misc Sites; Disease: All

Interventions

There are two intervention arms:

1. Treatment as usual (TAU) plus acceptance commitment therapy (ACT)
2. TAU plus talking control (TC)

Interventions will be described in detailed manuals to ensure core components are well defined and replicable. For ACT and TC, up to 8 sessions (each 1 hour) will be offered weekly and delivered within 3 months. Sessions will usually take place in a palliative care day therapy unit but the final 3 may occur at home if requested. Both ACT and TC will be delivered by the same therapist to reduce the effects of non-specific factors (e.g. warmth, professionalism).

1. ACT: a contextual behavioural approach which uses a collection of techniques aimed at increasing psychological flexibility to change outcomes. Psychological flexibility is the ability to

persist in valued life activities alongside distressing or unwanted private events. There are two main processes

1.1. Mindfulness and acceptance (cognitive fusion and self as context)

1.2. Commitment and behaviour change (being present, defining valued directions and committed action)

We shall follow the work of Hayes et al (1999) and supplement the approach with the manual "Get out of your mind and into your life" (Hayes and Smith, 2005). Therapists will be familiar with the skills training manual "Learning ACT" by Luoma, Hyes and Walser (2005) and "ACT made simple" by Harris (2009). Psychological flexibility mediates outcomes in chronic pain (Wicksell et al, 2010); acceptance, mindfulness and values each mediate outcomes for generalised anxiety.

2. TC: this promotes use of common factors in therapy by encouraging the therapist to be warm and welcoming, allowing people to ventilate their feelings, feel heard and understood. It specifically discourages focusing on problem areas and stipulates that problem solving should not be attempted. The therapist is trained to adhere to guidelines and this approach has been used previously to control for common factors in therapy.

Intervention Type

Other

Primary outcome(s)

FACT-G: A self-report 5 point Likert type scale, consisting of 27 questions taking 5 minutes to complete. It has four well-being domains (physical, social/family, emotional and activity) summed to a total score. With high internal consistency reliability and validity, it is recommended for assessing health-related quality of life in advanced cancer. A low score suggests poorer function. The mean score in a cancer population is 80.9 (SD 17.0). We shall select those with scores below the mean (pilot predicts 80% of population).

Key secondary outcome(s)

1. Psychological well-being: (5 minutes) Kessler-10 (K10, a 10-item self-report global measure of "psychological distress" in previous 4 weeks, taking 5 minutes to complete

2. Physical function: (5 minutes). Two minute walking test: the distance walked in 2 minutes (walking continuously, in a controlled space, at own speed, using an aid if necessary). One minute sit to stand test: number of times a person can stand up and sit down from a standardised chair over one minute.

3. Process of ACT: (10 minutes) Acceptance and Action Questionnaire II (AAQII): A 10 item scale measuring experiential avoidance that assesses willingness to accept undesirable thoughts and feelings, whilst acting in congruence with personal values and goals

4. Economic measures: (10 minutes)

4.1. EuroQol (EQ5D): A generic utility measure of quality of life consisting of 5-domains and a visual analogue scale, used in cost-effectiveness analysis

4.2. ICECAP-Supportive care measure: A seven item capability index for older people

Completion date

31/01/2018

Eligibility

Key inclusion criteria

1. A clinical diagnosis of advanced cancer defined as disease not amenable to curative treatment, those with metastases at diagnosis, those at first or subsequent extensive recurrence and those receiving palliative treatments

2. Total FACTG score of < 81
3. Agreement to be randomised
4. Sufficient understanding of English to engage in ACT

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

42

Key exclusion criteria

- 1 Clinician estimated survival of less than 4 months
2. Age under 18 years

Date of first enrolment

01/09/2015

Date of final enrolment

01/06/2016

Locations**Countries of recruitment**

United Kingdom

England

Study participating centre

Marie Curie Hampstead Hospice

Lyndhurst Gardens

London

United Kingdom

NW3 5NS

Study participating centre

St John's Hospice
60 Grove End Road
London
United Kingdom
NW8 9NH

Study participating centre
St. Joseph's Hospice
Mare St
London
United Kingdom
E8 4SA

Sponsor information

Organisation
Camden and Islington NHS Foundation Trust (UK)

ROR
<https://ror.org/03ekq2173>

Funder(s)

Funder type
Government

Funder Name
National Institute for Health Research

Alternative Name(s)
National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type
Government organisation

Funding Body Subtype
National government

Location
United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		21/12/2018	18/08/2023	Yes	No
Protocol article	protocol	11/02/2016		Yes	No
HRA research summary			28/06/2023	No	No
Other publications	Qualitative analysis of sessions	20/05/2022	18/08/2023	Yes	No