

# Effects of a standardized mother-child intervention programme in adolescent mothers and their children

<b>Submission date</b> 19/09/2012	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
<b>Registration date</b> 07/01/2013	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 26/10/2020	<b>Condition category</b> Other	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

In Germany about 7,500 babies are born each year to mothers below the age of 18, and 22,000 to mothers below age 20. Teenage pregnancy in Germany is more common among individuals with lower socio-economic status and poor education. Childhood neglect, physical and sexual abuse, inconsistent parenting and being a child to a teenage mother themselves are all risk factors for pregnancy in adolescence. Thus, children of adolescent mothers are a high-risk group for child neglect and maltreatment, especially if other sources of stress are present. Their development is often disturbed and they display more aggressive behaviour than children of adult mothers. Based on the results of a previous study, a training programme for young mothers was developed (STEEP, Steps Toward Effective and Enjoyable Parenting). However, so far nothing is known about whether a child's well-being can be increased by such a training programme. We want to find out whether STEEP helps to develop maternal sensitivity and how this effects attachment and child well-being.

### Who can participate?

120 mothers who were pregnant before their 21st birthday (adolescent mothers) and their children aged 3-6 months, and 40 mothers aged 25 or over (adult mothers).

### What does the study involve?

The adolescent mothers are randomly allocated to one of two groups. The intervention group will take part in a programme based on the principles of STEEP. The mothers are visited every 2 weeks by the same adviser until the child is 12 months old. Every second visit a short video of the mother and child interacting will be analyzed together with the mother, which has proven to be a promising way to increase sensitive parenting. In addition, psychiatric diseases in the mother (such as depression) will be treated by a psychiatrist/psychologist if needed. The other group will receive treatment as usual (support from the child welfare office). For further comparisons 40 mothers aged 25 or over (adult mothers) and matched for education will also be assessed. The study involves three assessment points: one at the beginning of the study, one

after 9 months of treatment and a third one after another 6 months. Participants are interviewed, complete questionnaires, and are asked to leave saliva, hair and blood samples and to undergo an MRI scan.

What are the possible benefits and risks of participating?

The STEEP intervention may improve the relationship between the mother and the child. There are no risks of participating.

Where is the study run from?

University Hospital of the Rheinisch-Westfaelische Technische Hochschule Aachen (Germany).

When is the study starting and how long is it expected to run for?

October 2012 to June 2016.

Who is funding the study?

Federal Ministry of Education and Research (Germany).

Who is the main contact?

Prof Beate Herpertz-Dahlmann

b.herpertzdahlmann@ukaachen.de

## Contact information

**Type(s)**

Scientific

**Contact name**

Prof Beate Herpertz-Dahlmann

**ORCID ID**

<https://orcid.org/0000-0001-8450-3323>

**Contact details**

University Hospital of the Rheinisch-Westfaelische Technische Hochschule Aachen

Department of Child and Adolescent Psychiatry and Psychotherapy

Neuenhofer Weg 21

Aachen

Germany

52074

+49 (0)241 808 8737

b.herpertzdahlmann@ukaachen.de

## Additional identifiers

**Protocol serial number**

DRKS00004409

## Study information

**Scientific Title**

Behavioural, developmental and neural effects of a standardized mother-child intervention programme in adolescent mothers and their children

## **Acronym**

TeeMo

## **Study objectives**

Within a RCT design, a parental programme based on STEEP plus psychiatric treatment of the mother in case of mental illness will be compared to TAU (standardized support by the child welfare office) in adolescent, high-risk mothers. The primary outcome variables are the maternal sensitivity and the child's well-being. Moderators of treatment outcome (e.g. child development, genotype, child temperament, neural and hormonal mechanism associated with parental bonding in the mother) as well as treatment effects in the child (cognitive and somatic development, structural brain maturation) will be systematically explored. A control group of adult mothers will be compared with the adolescent group.

On 04/09/2015 the overall trial end date was changed from 31/03/2014 to 30/06/2016.

## **Ethics approval required**

Old ethics approval format

## **Ethics approval(s)**

University Hospital of Aachen, 10/09/2012, ref: 12-006

## **Study design**

Randomised controlled trial with three arms

## **Primary study design**

Intentional

## **Study type(s)**

Quality of life

## **Health condition(s) or problem(s) studied**

Prevention of child neglect in adolescent mothers

## **Interventions**

Three arms:

1. Adolescent intervention group
2. Adolescent treatment as usual (TAU)
3. Adult controls

The active group will take part in a programme based on the principles of STEEP when the children are at least 3 months of age. They are visited every 2 weeks by the same adviser for the duration of 9 months. The topics of the home visits will be built on the manual by Erickson /Egeland (German edition by Sues, 2009). On average mother-child dyads will receive 12-18 home visits. Every second visit a short video-documentation of mother-child interaction will be analyzed together with the mother, which has proven to be a promising tool to increase sensitive parenting. In addition, psychiatric diseases in the mother (such as depression, etc.) will be treated by a psychiatrist/psychologist if needed.

## **Intervention Type**

Other

## **Phase**

Not Applicable

## **Primary outcome(s)**

1. Maternal sensitivity score, assessed by the sensitivity-scale of the Emotional Availability Scale (EAS)
2. Child's "well-being", assessed with the EAS-Scales
3. Child's responsiveness to the parent
4. Child's involvement of the parent

There will be three appointments: first one at the beginning of the intervention right after the inclusion, next one after intervention (9 months after inclusion) and the last one 6 months after the end of intervention (15 Months after inclusion). TAU and the adult control group will be assessed with parallel appointments.

## **Key secondary outcome(s)**

1. BIS-15 Barratt Impulsiveness Scale
2. BDI-II Becks Depression Inventory, Second Edition
3. BSDI-III Bayley Scales of Infant Development
4. CAARS Conners Adult ADHD Rating Scale
5. CAPI Child Abuse Potential Inventory (deutsche Version: EBSK)
6. CECA-Q Childhood Experiences of Care Abuse Questionnaire
7. CFT 20-R Culture Fair Intelligence Test - Scale 2, Revision
8. ECR-R Experiences in Close Relationships Scale-Revised
9. DERS Difficulties in Emotion Regulation Scale
10. IBQ-R Infant Behavior Questionnaire
11. IPDE The International Personality Disorder Examination
12. IRI; EC Interpersonal Reactivity Index
13. M.I.N.I. Mini-International Neuropsychiatric Interview
14. PSI Parental Stress Index (deutsche Version: EBI)
15. SST Strange Situation Test (deutsch FST)
16. STAI State-Trait Anxiety Inventory
17. VASQ Vulnerable Attachment Style Questionnaire
18. Analysis of fMRI-paradigms (Baby Face Incentive Delay Task and Infant Cry Self Distraction Paradigma)

## **Completion date**

30/06/2016

## **Eligibility**

### **Key inclusion criteria**

1. Child 3-6 months of age
2. Mother pregnant before 21st birthday
3. First or second child
4. Verbal intervention possible (no language or intellectual constraint)

5. Caucasian mother
6. Mother and child live together
7. Informed consent

**Participant type(s)**

Other

**Healthy volunteers allowed**

No

**Age group**

Mixed

**Sex**

All

**Total final enrolment**

56

**Key exclusion criteria**

1. Addiction (except for nicotine)
2. Birth of the child before 35th week of pregnancy was fulfilled
3. Syndromes or severe illnesses within the child
4. Severe psychiatric illness or suicidality within the mother
5. Separation of mother and child more than 3 months
6. Another video-intervention is planned

**Date of first enrolment**

01/10/2012

**Date of final enrolment**

01/11/2015

**Locations****Countries of recruitment**

Germany

**Study participating centre**

University Hospital of the Rheinisch-Westfaelische Technische Hochschule Aachen  
Aachen  
Germany  
52074

**Sponsor information**

## Organisation

Federal Ministry of Education and Research [BMBF] (Germany)

## ROR

<https://ror.org/04pz7b180>

## Funder(s)

### Funder type

Government

### Funder Name

Bundesministerium für Bildung und Forschung

### Alternative Name(s)

Federal Ministry of Research, Technology and Space, Bundesministerium für Bildung und Forschung, Federal Ministry of Education and Research, BMBF

### Funding Body Type

Government organisation

### Funding Body Subtype

National government

### Location

Germany

## Results and Publications

### Individual participant data (IPD) sharing plan

#### IPD sharing plan summary

Stored in repository

#### Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	10/07/2020	26/10/2020	Yes	No
<a href="#">Protocol article</a>	protocol	27/05/2015	26/10/2020	Yes	No
<a href="#">Study website</a>	Study website	11/11/2025	11/11/2025	No	Yes