# Effect of XP-endo finisher on microbial root canal flora after using multiple irrigation protocols

Submission date	Recruitment status	<ul><li>Prospectively registered</li></ul>
16/04/2025	No longer recruiting	Protocol
Registration date	Overall study status	Statistical analysis plan
28/04/2025	Completed	Results
Last Edited	Condition category	Individual participant data
28/04/2025	2 2	[X] Record updated in last year

# Plain English summary of protocol

Background and study aims

This study focuses on root canal infections, which are caused by bacteria and can lead to serious dental problems. The goal is to find the best way to clean the root canal using different solutions. Researchers are comparing three methods: using 5.25% sodium hypochlorite alone, combining it with 2% chlorhexidine, and combining it with QMIX. They are also looking at whether using a special tool called the XP-endo Finisher makes a difference.

#### Who can participate?

People who are between 20 and 40 years old, have no preoperative pain, have necrotic teeth, have a periapical radiolucency of 1-5 mm in diameter, and haven't taken antibiotics in the past three months.

#### What does the study involve?

The study involves treating 60 single-rooted teeth in patients and 60 freshly extracted single-rooted teeth in a lab. The teeth are divided into six groups, each using different cleaning methods with or without the XP-endo Finisher.

What are the possible benefits and risks of participating?

The main benefit is improved dental health and appearance. The risks are minimal and almost non-existent.

Where is the study run from? Damascus University (Syria)

When is the study starting and how long is it expected to run for? March 2021 to March 2025.

Who is funding the study? Damascus University (Syria)

# Contact information

## Type(s)

Public, Scientific, Principal investigator

#### Contact name

Dr Farah Estefane

#### **ORCID ID**

https://orcid.org/0009-0007-6488-6735

## Contact details

Damascus University, Mazzah Damascus Syria

\_ -

+963 988704477

farah3.estefan@damascusuniversity.edu.sy

# Type(s)

Public, Scientific, Principal investigator

#### Contact name

Dr Farah Estefane

#### Contact details

Damascus University, Mazzah Damascus Syria

\_\_\_\_\_

+963 988704477 estefanfarah3@gmail.com

# Additional identifiers

# Clinical Trials Information System (CTIS)

Nil known

# ClinicalTrials.gov (NCT)

Nil known

#### Protocol serial number

Nil known

# Study information

#### Scientific Title

A comparative study of several final irrigation protocols and the use of XP-endo finisher in changing of the microbial root canal flora

## **Study objectives**

H<sub>1</sub>: There is no significant difference in bacterial count and antibacterial efficacy against Enterococcus faecalis after root canal irrigation using sodium hypochlorite (NaOCl) without XP-endo Finisher compared to irrigation using sodium hypochlorite with XP-endo Finisher.

H<sub>2</sub>: There is no significant difference in bacterial count and antibacterial efficacy against Enterococcus faecalis after root canal irrigation using sodium hypochlorite (NaOCl) with XP-endo Finisher compared to irrigation using sodium hypochlorite without XP-endo Finisher.

H<sub>3</sub>: There is no significant difference in bacterial count and antibacterial efficacy against Enterococcus faecalis after root canal irrigation using sodium hypochlorite (NaOCl) combined with QMiX without XP-endo Finisher compared to irrigation using sodium hypochlorite with QMiX with XP-endo Finisher.

H<sub>4</sub>: There is no significant difference in bacterial count and antibacterial efficacy against Enterococcus faecalis after root canal irrigation using sodium hypochlorite (NaOCl) combined with QMiX with XP-endo Finisher compared to irrigation using sodium hypochlorite with QMiX without XP-endo Finisher.

H<sub>5</sub>: There is no significant difference in bacterial count and antibacterial efficacy against Enterococcus faecalis after root canal irrigation using sodium hypochlorite (NaOCl) combined with chlorhexidine (CHX) without XP-endo Finisher compared to irrigation using sodium hypochlorite with chlorhexidine with XP-endo Finisher.

H<sub>6</sub>: There is no significant difference in bacterial count and antibacterial efficacy against Enterococcus faecalis after root canal irrigation using sodium hypochlorite (NaOCl) combined with chlorhexidine (CHX) with XP-endo Finisher compared to irrigation using sodium hypochlorite with chlorhexidine without XP-endo Finisher.

# Ethics approval required

Ethics approval required

# Ethics approval(s)

approved 21/02/2022, The Biomedical Research Ethics Committee (BMREC) of Damascus University (Damascus University, Damascus, -, Syria; +963 1133923482; ap. srd@damascusuniversity.edu.sy), ref: DN-210224-185

# Study design

Interventional randomized controlled trial

# Primary study design

Interventional

# Study type(s)

Treatment

Health condition(s) or problem(s) studied

#### Patients with necrotic pulp

#### **Interventions**

Root canal treatment. The study samples (both clinical and laboratory) were divided into six groups: Groups 1 and 2: Irrigation was performed using 5.25% sodium hypochlorite with and without the XP-endo Finisher, respectively. Groups 3 and 4: Irrigation was performed using 5.25% sodium hypochlorite with 2% chlorhexidine, with and without the XP-endo Finisher, respectively. Groups 5 and 6: Irrigation was performed using 5.25% sodium hypochlorite with QMIX, with and without the XP-endo Finisher, respectively.

The randomization process was performed using sealed opaque envelopes managed by an independent party.

#### Intervention Type

Other

#### Primary outcome(s)

To compare the bacterial count before and after clinical root canal preparation using three final irrigation protocols:

- 1. Sodium hypochlorite (NaOCl) alone
- 2. Sodium hypochlorite combined with chlorhexidine (CHX)
- 3. Sodium hypochlorite combined with QMiX

# Key secondary outcome(s))

There are no secondary outcome measures

# Completion date

14/03/2025

# **Eligibility**

## Key inclusion criteria

- 1. Patients medically free from any systemic diseases
- 2. Patients between the age group of 20 and 40 years old
- 3. Absence of preoperative pain
- 4. Necrotic cases only
- 5. Presence of periapical radiolucency 1-5 mm in diameter
- 6. No antibiotics in the past three months

# Participant type(s)

Patient

## Healthy volunteers allowed

No

# Age group

Adult

# Lower age limit

20 years

# Upper age limit

40 years

#### Sex

All

#### Total final enrolment

60

# Key exclusion criteria

- 1. Teeth with symptomatic irreversible pulpitis
- 2. Teeth with open apices
- 3. Consumption of any type of antibiotics in the past three months
- 4. Systemic diseases
- 5. Pregnant and lactating females

#### Date of first enrolment

01/01/2022

## Date of final enrolment

01/01/2023

# Locations

#### Countries of recruitment

Syria

# Study participating centre

# **Damascus University**

Department of Enodontics, Faculty of Dental Medicine, Mazzah Damascus Syria

\_

# Sponsor information

## Organisation

**Damascus University** 

#### **ROR**

https://ror.org/03m098d13

# Funder(s)

# Funder type

University/education

#### **Funder Name**

**Damascus University** 

#### Alternative Name(s)

University of Damascus, , DU

#### **Funding Body Type**

Government organisation

#### **Funding Body Subtype**

Universities (academic only)

#### Location

Syria

# **Results and Publications**

## Individual participant data (IPD) sharing plan

The datasets generated during and/or analyzed during the current study are/will be available upon request from Prof. Dr Kinda Layous (kinda.layous@damascusuniversity.edu.sy). All of the patients' data will be available upon request. Consent was obtained from the participants.

# IPD sharing plan summary

Available on request

# Study outputs

Output type Details Date created Date added Peer reviewed? Patient-facing?

Participant information sheet 11/11/2025 No Yes