

# Does expansion of the upper jaw in cases of posterior crossbite affect the position of the temporomandibular joint and the occlusion of the teeth?

<b>Submission date</b> 01/10/2024	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 04/10/2024	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
<b>Last Edited</b> 03/12/2024	<b>Condition category</b> Oral Health	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

This study aims to compare two different dental devices, the quad-helix and removable expansion plates, to see which is more effective in treating a condition called unilateral posterior crossbite in children. This condition involves a misalignment of the teeth and jaw, causing difficulties in biting and chewing.

### Who can participate?

The study includes 40 children aged between 7 and 10 years who have a unilateral posterior crossbite with a shift in their lower jaw.

### What does the study involve?

Participants are randomly assigned to one of two groups. One group uses the quad-helix device, while the other uses removable expansion plates. Impressions of their teeth are taken at four different times: before treatment starts, after the active treatment phase, after a three-month stabilization phase, and after a six-month monitoring phase.

### What are the possible benefits and risks of participating?

The potential benefit of participating is the correction of the crossbite, which can improve dental function and appearance. Risks might include discomfort from wearing the devices and the usual risks associated with dental treatments.

### Where is the study run from?

Department of Orthodontics, Faculty of Dentistry, Damascus University (Syria)

### When is the study starting and how long is it expected to run for?

The study started on February 20, 2022, and is expected to run until December 10, 2024.

Who is funding the study?  
Damascus University (Syria)

Who is the main contact?  
Dr Aynawi, m.aynawi93@gmail.com

## Contact information

### Type(s)

Public, Scientific, Principal investigator

### Contact name

Dr Mohammad Aynawi

### Contact details

Department of Orthodontics  
Faculty of Dentistry  
Damascus University  
Al-Mazzeah St.  
Damascus  
Syria  
0004  
+963 (0)930 899 344  
Mohammad93.aynawi@damascusuniversity.edu.sy

### Type(s)

Scientific

### Contact name

Prof Ahmad Burhan

### ORCID ID

<https://orcid.org/0000-0002-0727-2653>

### Contact details

Department of Orthodontics  
Faculty of Dentistry  
Damascus University  
Al-Mazzeah St.  
Damascus  
Syria  
0004  
+963 (0)944302075  
Prof.ahmadbouhan@damascusuniversity.edu.sy

## Additional identifiers

### Clinical Trials Information System (CTIS)

Nil known

**ClinicalTrials.gov (NCT)**

Nil known

**Protocol serial number**

5414

## Study information

### Scientific Title

Efficacy of the removable expansion plate and the quad-helix in correcting unilateral posterior crossbite in the mixed dentition: a randomized controlled clinical trial

### Study objectives

1. There is no difference in the effectiveness of the quad-helix appliance and the removable expansion plate in terms of changes in dental alveolar widths
2. There is no difference in the effectiveness of the quad-helix device and the removable expansion plate in the amount of correction of the deviation of the lower dental and mandibular midline

### Ethics approval required

Ethics approval not required

### Ethics approval(s)

No ethics approval is required because the devices used have been previously tested on humans.

### Study design

Randomized controlled clinical trial

### Primary study design

Interventional

### Study type(s)

Treatment

### Health condition(s) or problem(s) studied

Malocclusion, narrow upper jaw

### Interventions

Participants were randomly divided into two groups using a simple computer randomization method at a 1:1 allocation ratio. The patients in the first group were expanded using quad-helix, and the second group was expanded using removable expansion plates.

Impressions were taken at the following evaluation times: (T0): before the expansion begins, (T1): after the end of the active treatment phase, (T2): after the end of the stabilization phase, which lasts 3 months, (T3): after completing the monitoring phase, which lasts 6 months, starting from after the end of the stabilization phase.

### Intervention Type

Device

## Phase

Not Applicable

## Drug/device/biological/vaccine name(s)

Quad-helix or removable expansion device

## Primary outcome(s)

1. Maxillary intermolar distance is measured using a digital caliper at T0, T1, T2, and T3
2. Lower intermolar distance is measured using a digital caliper at T0, T1, T2, and T3
3. Maxillary intercanine distance is measured using a digital caliper at T0, T1, T2, and T3
4. Lower canine distance is measured using a digital caliper at T0, T1, T2, and T3
5. Medial joint space is measured using CBCT at T0 and T1
6. Distal joint space is measured using CBCT at T0 and T1
7. Anterior joint space is measured using CBCT at T0 and T1
8. Posterior joint space is measured using CBCT at T0 and T1
9. Superior joint space is measured using CBCT at T0 and T1
10. Angle of the condyle in the horizontal plane is measured using CBCT at T0 and T1
11. Angle of the condyle in the frontal plane is measured using CBCT at T0 and T1
12. Angle of the condyle in the sagittal plane is measured using CBCT at T0 and T1

(T0): before the expansion begins

(T1): after the end of the active treatment phase.

(T2): after the end of the stabilization phase, which lasts 3 months

(T3): After completing the monitoring phase, which lasts 6 months, starting from the end of the stabilization phase.

## Key secondary outcome(s)

1. Pain is measured using the visual analogue score (VAS) at baseline, 24 hours, 1 week, 2 weeks, and 1 month

## Completion date

10/12/2024

## Eligibility

### Key inclusion criteria

1. Patients with mixed occlusion, aged 7-10 years
2. The presence of a functional unilateral posterior crossbite (associated with lateral slippage)
3. Symmetrical maxillary arch narrowing or symmetrical skeletal narrowing (assessed clinically and then radiographically)
4. The maxillary first molars are erupted and in good condition, with full eruption of the upper and lower molars (at a minimum)
5. Dental and skeletal relationships of the first, light second, or light third category (meaning that the angle ANB falls between 1-5), protrusion between 0.5 to 4 mm, coverage between 0.5 to 4 mm
6. A normal or mild vertical growth model such that the Bjork sum is between 390-406, the Y axis is between 62-72 degrees, and the angle between SN and GoMe is 26-42 degrees
7. There are no general problems, the patient has good oral health, and has not undergone previous orthodontic treatments

## Participant type(s)

Patient

**Healthy volunteers allowed**

No

**Age group**

Child

**Lower age limit**

7 years

**Upper age limit**

10 years

**Sex**

All

**Total final enrolment**

37

**Key exclusion criteria**

1. The presence of periodontal diseases, general diseases, syndromes (cleft lip and palate) or systemic diseases that affect growth
2. Patients who have undergone previous orthodontic treatment
3. If the structural relationship is of the second or third category, moderate or severe
4. Patients with anterior crossbites as well as posterior crossbites
5. Patients with anterior open bites as well as posterior crossbites
6. Patients whose functional posterior crossbites are characterized by anterolateral slippage
7. Patients whose chin deviation on closure is of common structural and functional origin
8. Patients in whom the path of mouth opening from the rest position to the maximum possible position is tortuous or disturbed

**Date of first enrolment**

20/04/2022

**Date of final enrolment**

10/04/2024

## **Locations**

**Countries of recruitment**

Syria

**Study participating centre**

**Damascus University**

Faculty of Dentistry

Department of Orthodontics

Al-Mazzeah St.

Damascus

Syria  
80789

## Sponsor information

### Organisation

Damascus University

### ROR

<https://ror.org/03m098d13>

## Funder(s)

### Funder type

University/education

### Funder Name

Damascus University

### Alternative Name(s)

University of Damascus, , DU

### Funding Body Type

Government organisation

### Funding Body Subtype

Universities (academic only)

### Location

Syria

## Results and Publications

### Individual participant data (IPD) sharing plan

The data-sharing plans for the current study are unknown and will be made available at a later date

### IPD sharing plan summary

Data sharing statement to be made available at a later date