# The ADDapt diet in reducing Crohn's disease inflammation

Submission date	Recruitment status No longer recruiting	[X] Prospectively registered		
29/07/2019		☐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
29/07/2019	Completed  Condition category	☐ Results		
Last Edited		Individual participant data		
17/10/2024	Digestive System	<ul><li>Record updated in last year</li></ul>		

# Plain English summary of protocol

Background and study aims

Crohn's disease (CD) results in chronic intestinal inflammation, is of increasing incidence both in the developed and developing world and has a marked impact on patient quality of life. The prevalence of CD is 10.6 per 100,000 people in the UK and represents a significant annual financial burden of around €16.7 million in Europe.

A wide range of nutrients and food components have been investigated for their role in the pathogenesis and course of CD. A common theme suggests that CD risk is associated with a "Western diet", including high fat, high sugar and processed foods. However, intervention studies that exclude specific aspects of the diet such as sugar or that compare low and high fat diets have failed to show effectiveness in practice. Observational human and experimental animal studies suggest that certain food additives used extensively by the food industry play a role in the pathogenesis and natural history of CD. However, to date no evidence exists for the effectiveness of a diet low in these food additives in CD.

Therefore, the aim of this study is to investigate the effects of a diet low in certain food additives compared to a normal UK diet on CD activity, health-related quality of life, gut bacteria, gut permeability, gut inflammation and dietary intake, in patients with mildly active, stable CD. We will recruit patients with mildly active CD and will randomise them to receive either the diet low in the food additives of interest, or the diet representative of a normal UK diet. Patients will follow the diet for 8 weeks and will attend study visits at the start and end of the trial, at which points questionnaires will be completed and samples will be collected.

# Who can participate?

Adults aged 16 years or older (updated 03/02/2022, previously: aged 18 years or older) with mildly active Crohn's disease.

### What does the study involve?

Patients will be asked to follow a set diet for 8-weeks, during which time they will attend clinic visits to gather information on food intake, general health, and to provide samples for analysis.

# What are the possible benefits and risks of participating?

The results of this study may help to answer scientific questions about whether reducing specific food ingredients improves intestinal inflammation in patients with active Crohn's disease, and

whether there is any beneficial effect on the gut bacteria. The ADDapt diet can be challenging to follow because certain pre-prepared and convenience foods are excluded. There are still plenty of convenience foods, pre-prepared foods and ready-meals that are suitable, and these can still be eaten provided you check their suitability with the dietitian. The ADDapt diet is nutritionally balanced and your dietitian will be able to help answer any questions you have whilst you are on the diet. If you are preparing food for other people, it is safe for them to eat the same meals as you.

Where is the study run from? King's College, London

When is the study starting and how long is it expected to run for? July 2019 to December 2024

Who is funding the study?

- 1. Leona M. and Harry B. Helmsley Charitable Trust
- 2. National Institute for Health Research (NIHR), UK

Who is the main contact? Dr Aaron Bancil aaron.bancil@kcl.ac.uk

# Contact information

Type(s)

Scientific

#### Contact name

Dr Aaron Bancil

#### Contact details

King's College London
4.103 Franklin Wilkins Building
150 Stamford Street
London
United Kingdom
SE1 9NH
+44 (0)207 848 4552
aaron.bancil@kcl.ac.uk

# Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

NCT04046913

Protocol serial number

**CPMS 41991** 

# Study information

#### Scientific Title

The ADDapt diet in reducing Crohn's disease inflammation

#### **Acronym**

**ADDapt** 

# Study objectives

There is a difference in the proportion of patients achieving at least a 70 point reduction in the Crohn's Disease Activity Index (CDAI) between baseline and end of the trial, between the two diet groups.

# Ethics approval required

Old ethics approval format

# Ethics approval(s)

Approved 15/07/2019, East of Scotland Research Ethics Service (EoSRES) (Tayside medical Science Centre, Residency Block Level 3. George Pirie Way, Ninewells Hospital and Medical School, Dundee, DD1 9SY; +441382383878; eosres.tayside@nhs.net), ref: 19/ES/0049

# Study design

Randomized; Interventional; Design type: Treatment, Dietary

# Primary study design

Interventional

# Study type(s)

Quality of life

# Health condition(s) or problem(s) studied

Crohn's disease

#### Interventions

Current intervention as of 03/02/2022:

This will be an 8-week randomised double-blind, placebo-controlled trial in 154 patients with mildly active, stable Crohn's disease. Patients will be recruited from five London inflammatory bowel disease (IBD) centres.

These will be Guy's and St Thomas' NHS Foundation Trust, Barts Health NHS Trust, St Marks Hospital, University College Hospital NHS Foundation Trust and Royal Free Hospital NHS Foundation Trust. Further trusts have been added recently.

In this trial, all patients will follow a low food additive background diet. Patients will be randomised to either the low food additive diet or control diet (mimicking a habitual UK diet) group, in a 1:1 ratio. Block randomisation will be stratified according to Crohn's Disease location (ileal, ileo-colonic, colonic) and whether the patient opts for sigmoidoscopy or not. The randomisation schedule will be constructed using an online program by a researcher not involved in trial planning or recruitment. The researcher randomising participants and measuring trial outcomes will not be aware of diet allocation.

In addition to following the background low food additive diet, patients in the diet groups will receive the following foods:

Low food additive diet (intervention) group:

- Additive-free trial foods formulated specifically for this project, to be eaten in specific portions every day during the trial.

Control group:

- A supply of food additive-containing trial foods formulated specifically for this project.

# Trial procedures:

#### Screening:

Trial investigators will attend gastroenterology clinics at each of the five recruitment sites each week. Gastroenterologists, IBD nurses and IBD pharmacists will refer potentially suitable and willing patients for screening. Patients will be given a participant information sheet (PIS) and offered their hospital's leaflet on sigmoidosopies and will be screened against inclusion and exclusion criteria. Patients will be given at least 24 hours to consider the study and ask questions prior to consent.

Suitable patients wishing to take part will be allowed to consent to the trial on the day of screening if this is more convenient for them than attending a separate consent visit. Otherwise, patients will attend the hospital clinic or the metabolic research unit at King's College London to provide consent.

After consent, eligible patients will be instructed by an experienced research dietitian to complete a 7-day food and symptom diary, which will be used to assess disease activity and dietary intake.

#### Baseline visit:

Patients meeting all of the inclusion criteria and none of the exclusion criteria and having consented will attend a baseline visit at either the hospital clinic or the metabolic research unit at King's College London (whichever is most convenient for the patient) following an overnight fast. The following procedures will be undertaken:

- Demographic characteristics
- Patient-completed questionnaires
- A whole stool sample will be collected within 2 hours of the visit
- A baseline urine sample will be collected for intestinal permeability
- Blood samples will be collected
- Participants opting into the sigmoidoscopy sub study (n=77) will undergo an unprepared flexible sigmoidoscopy
- All participants will be provided with thorough instruction on the low food additive diet by an experienced specialist dietitian.

Patients will be randomized to either the low food additive diet or control diet and will receive a supply of the foods described above.

Monitoring and safety: Patients will be given contact details for the study investigators and will be telephoned once a week to monitor progress with the intervention, compliance to the low food additive diet and consumption of the trial foods. Furthermore, any reported adverse events will be recorded at these contacts, and investigators will address any concerns that patients may have regarding any aspects of the trial.

Compliance: Compliance with the experimental diet allocation and the trial foods will be monitored and encouraged at the weekly telephone contacts.

Follow-up visit: After the 8-week trial, participants will return for an end of trial visit after an overnight fast. Participants will complete a food and symptom diary, identical to that completed during screening, for 7 days prior to this visit. At this visit, the following will take place:

- The Crohn's Disease Activity Index (CDAI) score will be calculated
- Stool, urine and blood samples will be collected using the methods described in the baseline visit. Flexible, unprepared sigmoidoscopies will be performed as described in the baseline visit

# Long-term follow-up:

Following completion of the 8-week trial, patients will have the option of continuing to follow the experimental diet for an additional 16 weeks. At monthly intervals during this period, a telephone visit will be conducted, to investigate diet compliance and acceptability and to provide patient support.

At the end of trial visit (following the 8-week trial), patients will be provided with a blank 7-day food and symptom diary and will be instructed to complete it for the 7 days prior to a long-term visit, which will take place after a total of 24 weeks

At the long-term visit, the following procedures will take place:

- CDAI score will be calculated
- Patient-completed questionnaires
- A whole stool sample and blood samples will be collected using the methods described above. Following completion of the trial (or withdrawal), patients will be instructed to return to their normal diet and will be provided with written general dietary guidance for Crohn's disease (Crohn's and Colitis UK).

#### Previous intervention:

This will be an 8-week randomised double-blind, placebo-controlled trial in 154 patients with mildly active, stable Crohn's disease. Patients will be recruited from five London inflammatory bowel disease (IBD) centres.

These will be Guy's and St Thomas' NHS Foundation Trust, Barts Health NHS Trust, St Marks Hospital, University College Hospital NHS Foundation Trust and Royal Free Hospital NHS Foundation Trust.

In this trial, all patients will follow a low food additive background diet. Patients will be randomised to either the low food additive diet or control diet (mimicking a habitual UK diet) group, in a 1:1 ratio. Block randomisation will be stratified according to Crohn's Disease location (ileal, ileo-colonic, colonic) and whether the patient opts for sigmoidoscopy or not. The randomisation schedule will be constructed using an online program by a researcher not involved in trial planning or recruitment. The researcher randomising participants and measuring trial outcomes will not be aware of diet allocation.

In addition to following the background low food additive diet, patients in the diet groups will receive the following foods:

Low food additive diet (intervention) group:

- Additive-free trial foods formulated specifically for this project, to be eaten in specific portions every day during the trial.

Control group:

- A supply of food additive-containing trial foods formulated specifically for this project.

#### Trial procedures:

#### Screening:

Trial investigators will attend gastroenterology clinics at each of the five recruitment sites each week. Gastroenterologists, IBD nurses and IBD pharmacists will refer potentially suitable and

willing patients for screening. Patients will be given a participant information sheet (PIS) and offered their hospital's leaflet on sigmoidosopies and will be screened against inclusion and exclusion criteria. Patients will be given at least 24 hours to consider the study and ask questions prior to consent.

Suitable patients wishing to take part will be allowed to consent to the trial on the day of screening if this is more convenient for them than attending a separate consent visit. Otherwise, patients will attend the hospital clinic or the metabolic research unit at King's College London to provide consent.

After consent, eligible patients will be instructed by an experienced research dietitian to complete a 7-day food and symptom diary, which will be used to assess disease activity and dietary intake.

#### Baseline visit:

Patients meeting all of the inclusion criteria and none of the exclusion criteria and having consented will attend a baseline visit at either the hospital clinic or the metabolic research unit at King's College London (whichever is most convenient for the patient) following an overnight fast. The following procedures will be undertaken:

- Demographic characteristics
- Patient-completed questionnaires
- A whole stool sample will be collected within 2 hours of the visit
- A baseline urine sample will be collected for intestinal permeability
- Blood samples will be collected
- Participants opting into the sigmoidoscopy sub study (n=77) will undergo an unprepared flexible sigmoidoscopy
- All participants will be provided with thorough instruction on the low food additive diet by an experienced specialist dietitian.

Patients will be randomized to either the low food additive diet or control diet and will receive a supply of the foods described above.

Monitoring and safety: Patients will be given contact details for the study investigators and will be telephoned once a week to monitor progress with the intervention, compliance to the low food additive diet and consumption of the trial foods. Furthermore, any reported adverse events will be recorded at these contacts, and investigators will address any concerns that patients may have regarding any aspects of the trial.

Compliance: Compliance with the experimental diet allocation and the trial foods will be monitored and encouraged at the weekly telephone contacts.

Follow-up visit: After the 8-week trial, participants will return for an end of trial visit after an overnight fast. Participants will complete a food and symptom diary, identical to that completed during screening, for 7 days prior to this visit. At this visit, the following will take place:

- The Crohn's Disease Activity Index (CDAI) score will be calculated
- Stool, urine and blood samples will be collected using the methods described in the baseline visit. Flexible, unprepared sigmoidoscopies will be performed as described in the baseline visit

#### Long-term follow-up:

Following completion of the 8-week trial, patients will have the option of continuing to follow the experimental diet for an additional 16 weeks. At monthly intervals during this period, a telephone visit will be conducted, to investigate diet compliance and acceptability and to provide patient support.

At the end of trial visit (following the 8-week trial), patients will be provided with a blank 7-day food and symptom diary and will be instructed to complete it for the 7 days prior to a long-term visit, which will take place after a total of 24 weeks

At the long-term visit, the following procedures will take place:

- CDAI score will be calculated
- Patient-completed questionnaires
- A whole stool sample and blood samples will be collected using the methods described above. Following completion of the trial (or withdrawal), patients will be instructed to return to their normal diet and will be provided with written general dietary guidance for Crohn's disease (Crohn's and Colitis UK).

#### Intervention Type

Other

# Primary outcome(s)

Reduction in the Crohn's Disease Activity Index (CDAI) between baseline and end of trial (8-weeks)

# Key secondary outcome(s))

- 1. Crohn's disease activity as measured using the Crohn's Disease Activity Index (CDAI) at baseline, week 8 and week 26
- 2. Gastrointestinal inflammation as measured using faecal calprotectin at baseline, week 8 and week 26
- 3. Systemic inflammation as measured using CRP concentration at baseline, week 8 and week 26
- 4. Perceived disease control as measured by the IBD-control questionnaire at baseline, week 8 and week 26
- 5. Health related quality of life as measured by the Inflammatory Bowel Disease Questionnaire (IBD-Q) at baseline, week 8 and week 26.
- 6. Dietary intake as measured by 7-day food diaries at baseline, week 8 and week 26
- 7. Dietary compliance and acceptability as measured by acceptability questionnaire and food-related Quality of Life questionnaire at baseline, week 8 and week 26
- 8. Gastrointestinal permeability as measured by urine analysis at baseline and week 8
- 9. Gastrointestinal microbiota as measured by faecal microbiota composition and mucosal microbiota composition at baseline and week 8 (in a subset of participants)
- 10. Mucosal immune gene expression as measured on immune cells from rectal biopsies at baseline and week 8 (in a subset of participants)

# Completion date

31/12/2024

# **Eligibility**

# Key inclusion criteria

Current participant inclusion criteria as of 03/02/2022:

- 1. Adults aged ≥16 years
- 2. Crohn's disease diagnosis (defined by standard clinical, histological and radiological criteria) of at least 6 months
- 3. Mildly active disease as defined by:
- 3.1 Defined by physician assessment that no change in medication is required
- 3.2 Faecal calprotectin >150  $\mu$ g/g
- 3.3 CDAI between 150-250
- 3.4 Current body weight of ≥50 kg
- 3.5 Individuals able to give informed consent and willingness to participate

Previous participant inclusion criteria:

- 1. Adults aged > = 18 years
- 2. Crohn's disease diagnosis (defined by standard clinical, histological and radiological criteria) of at least 6 months
- 3. Mildly active disease as defined by:
- 3.1 Defined by physician assessment that no change in medication is required
- 3.2 Faecal calprotectin >  $150 \mu g/g$
- 3.3 CDAI between 150-250
- 3.4 Current body weight of > = 50 kg
- 3.5 Individuals able to give informed consent and willingness to participate

# Participant type(s)

Patient

# Healthy volunteers allowed

No

#### Age group

Adult

# Lower age limit

16 years

#### Sex

All

#### Total final enrolment

154

#### Key exclusion criteria

- 1. Changes in dose to azathioprine, 6-mercaptopurine, methotrexate or anti-TNF-a agents or other biologics during the preceding 8 weeks, oral 5-ASA during the preceding four weeks. Currently receiving oral prednisolone/budesonide or discontinued within the last 4 weeks, unless they are on a stable dose of 10 mg/day or less prednisolone (3 mg or less budesonide) for at least 4 weeks with the intention to continue this long term.
- 2. Used rectal 5-ASA or rectal steroids in the preceding 4 weeks
- 3. Previous extensive bowel resection, defined as having had > 2 intestinal resections, a sub-total colectomy or documented short bowel syndrome
- 4. Poorly controlled bile acid malabsorption
- 5. Current stoma
- 6. Recent use of the following treatments: antibiotics, probiotics, prebiotic or fibre supplements in the preceding four weeks, NSAIDs during the preceding week
- 7. Full bowel preparation for a diagnostic procedure in preceding 4 weeks
- 8. Comorbidities including sepsis/fever, diabetes or coeliac disease, or other concomitant serious comorbidity e.g. significant psychiatric, hepatic, renal, endocrine, respiratory, neurological or cardiovascular disease
- 9. Exclusive enteral nutrition in the past 8 weeks
- 10. Assessed as at nutritional risk, as defined by any of the following:
- $10.1 \text{ BMI} < = 18.5 \text{ kg/m}^2$

10.2 Previous or current eating disorder

10.3 Currently receiving prescribed oral nutritional supplements

11. Following a restrictive diet (e.g. multiple restrictions due to numerous self-reported allergies) as judged by the dietitian

12. Reported pregnancy or lactation

#### Date of first enrolment

12/08/2019

#### Date of final enrolment

29/02/2024

# **Locations**

#### Countries of recruitment

**United Kingdom** 

England

# Study participating centre Guy's Hospital

Guy's & St Thomas' NHS Foundation Trust Great Maze Pond London United Kingdom SE1 9RT

# Study participating centre Northwick Park Hospital

Watford Road Harrow United Kingdom HA1 3UJ

# Study participating centre

University College London Hospitals NHS Foundation Trust

235 Euston Road London United Kingdom NW1 2BU

# Study participating centre

# Royal Free Hospital

Pond Street London United Kingdom NW3 2QG

# Study participating centre The Royal London Hospital

Whitechapel Road Whitechapel London United Kingdom E1 1BB

# Study participating centre

Chelsea and Westminster Hospital NHS Foundation Trust

Chelsea & Westminster Hospital 369 Fulham Road London United Kingdom SW10 9NH

# Study participating centre Addenbrookes

Addenbrookes Hospital Hills Road Cambridge United Kingdom CB2 0QQ

# Study participating centre Pennine Acute Hospitals NHS Trust

Trust Headquarters
North Manchester General Hospital
Delaunays Road, Crumpsall
Manchester
United Kingdom
M8 5RB

# Study participating centre

# County Durham and Darlington NHS Foundation Trust

Darlington Memorial Hospital Hollyhurst Road Darlington United Kingdom DL3 6HX

# Study participating centre University Hospital Bristol

Bristol Royal Infirmary Marlborough Street Bristol United Kingdom BS2 8HW

# Study participating centre

The Newcastle upon Tyne Hospitals NHS Foundation Trust

Freeman Hospital Freeman Road High Heaton Newcastle upon Tyne United Kingdom NE7 7DN

# Study participating centre United Leeds Teaching Hospital NHS Trust

Trust Hq Leeds General Infirmary Great George St Leeds United Kingdom LS1 3EX

# Study participating centre

NHS Lothian Crewe Rd S Edinburgh United Kingdom EH4 2XU

Study participating centre

#### East Sussex Healthcare NHS Trust

The Ridge St Leonards-on-Sea United Kingdom TN37 7RD

# Study participating centre Royal Devon and Exeter NHS Foundation Trust

Royal Devon and Exeter Hospital Barrack Road Exeter United Kingdom EX2 5DW

# Study participating centre The Royal Wolverhampton NHS Trust

New Cross Hospital Wolverhampton United Kingdom WV10 0QP

# Study participating centre King's College Hospital NHS Foundation Trust

Denmark Hill London United Kingdom SE5 9RS

# Study participating centre Imperial College Healthcare NHS Trust

St Mary's Hospital Praed Street London United Kingdom W2 1NY

# Study participating centre University Hospital Southampton NHS Trust Foundation Trust

Tremona Road

# Sponsor information

#### Organisation

King's College London

#### **ROR**

https://ror.org/0220mzb33

# Funder(s)

# Funder type

Charity

#### **Funder Name**

Leona M. and Harry B. Helmsley Charitable Trust

#### Alternative Name(s)

Helmsley Charitable Trust, The Leona M. and Harry B. Helmsley Charitable Trust, Leona M. & Harry B. Helmsley Charitable Trust, The Helmsley Charitable Trust, The Leona M and Harry B Helmsley Charitable Trust, Helmsley

# Funding Body Type

Private sector organisation

# Funding Body Subtype

Trusts, charities, foundations (both public and private)

#### Location

United States of America

#### Funder Name

National Institute for Health Research

#### Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

# **Funding Body Type**

Government organisation

# **Funding Body Subtype**

National government

### Location

**United Kingdom** 

# **Results and Publications**

# Individual participant data (IPD) sharing plan

The current data sharing plans for this study are unknown and will be available at a later date

# IPD sharing plan summary

Data sharing statement to be made available at a later date

# **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			28/06/2023	No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes