

Metronidazole versus lactic acid for treating bacterial vaginosis

Submission date 18/09/2017	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 18/09/2017	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 04/10/2024	Condition category Urological and Genital Diseases	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Bacterial vaginosis (BV) is a common condition in women which causes a discharge from the vagina, often with an unpleasant fishy smell. This is embarrassing and often distressing. It is also associated with developing more generalised pelvic infection, the occurrence of complications in pregnancy (including miscarriage and premature delivery) and an increased risk of catching and passing on HIV. It affects 30-50% of women at some time in their lives. Currently an antibiotic called metronidazole is used to treat BV, but often BV comes back again in about a third of women, who then require repeated courses of antibiotic treatment. Antibiotics can also have unwanted side effects and antibiotic resistance can develop if they are used too often. The exact cause of BV is not known but it is associated with a change in the type of bacteria in the vagina, specifically a reduction in 'lactobacilli' and an increase in 'anaerobic bacteria'. Normally, lactobacilli make lactic acid in the vagina, which prevents the growth of anaerobic bacteria. However, in women with BV (where the numbers of lactobacilli have decreased) the vagina is less acid than usual. As a result, the increased number of anaerobic bacteria in the vagina produce various chemicals which cause inflammation and an unpleasant smell. The aim of this study is to see if using an alternative treatment known as lactic acid gel may be effective in treating BV by 'replacing' vaginal acidity in the vagina. Some previous studies have suggested that this approach could be successful but they are not conclusive and current guidelines highlight a need for more information before recommending the use of intravaginal lactic acid gel.

Who can participate?

Women aged 16 or above with symptoms of bacterial vaginosis (BV) and a history of one or more episodes of BV within the previous 2 years which resolved with treatment

What does the study involve?

At their single clinic visit, participants are asked to provide samples for analysis of BV and are then randomly allocated to receive treatment with either intravaginal lactic acid gel used once daily for 7 days (intervention group) or oral metronidazole tablets twice daily for 7 days (control group). Following the initial visit, participants are then followed up for resolution of their symptoms 2 weeks, 3 months and 6 months later with online questionnaires and self-sampling kits, completed at home.

What are the possible benefits and risks of participating?

There are no direct benefits for the participant of taking part in the study. However, the information from this study will help to determine whether topical lactic acid gel is an effective and well tolerated treatment for BV. This study will help to advance understanding and could lead to changes to the current guidelines for recommended BV treatments. Although there is some evidence to suggest that lactic acid gel may be effective in treating BV, we do not know whether it works as well as metronidazole. If it doesn't you may have to have further treatment. As with all medications there is a small risk of side effects, the GP or Nurse will be able to explain these.

Where is the study run from?

Nottingham Clinical Trials Unit (UK)

When is the study starting and how long is it expected to run for?

March 2017 to June 2020 (updated 18/02/2020, previously: November 2019)

Who is funding the study?

Health Technology Assessment Programme (UK)

Who is the main contact?

VITA Trial Team

vita@nottingham.ac.uk

Study website

<https://www.vitastudy.org>

Contact information

Type(s)

Scientific

Contact name

Dr Lindsay Armstrong-Buisseret

Contact details

Nottingham Clinical Trials Unit

University of Nottingham

Queens Medical Centre

Nottingham

United Kingdom

NG7 2RD

+44 115 823 1610

vita@nottingham.ac.uk

Additional identifiers

EudraCT/CTIS number

2016-004483-19

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

CPMS 33038

Study information

Scientific Title

Metronidazole Versus lactic acid for Treating bacterial vAginosis: A randomised controlled trial to assess the clinical and cost effectiveness of topical lactic acid gel for treating second and subsequent episodes of bacterial vaginosis

Acronym

VITA

Study objectives

The aim of this trial is to determine if using lactic acid gel in the vagina to 'replace' vaginal acidity is better than oral metronidazole for symptomatic resolution of recurrent bacterial vaginosis (BV). Some previous studies have suggested that this approach could be successful but they are not conclusive and current guidelines highlight a need for more evidence before recommending the use of intravaginal lactic acid gel. In parallel, the tolerability and acceptability of using lactic acid gel and metronidazole as treatments for BV will be assessed.

Ethics approval required

Old ethics approval format

Ethics approval(s)

London Harrow Research Ethics Committee, 09/09/2017, ref: 17/LO/1245

Study design

Randomized; Interventional; Design type: Treatment, Drug, Device

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Bacterial vaginosis (BV)

Interventions

At their single clinic visit, participants are asked to provide initial samples for analysis of BV and will then be randomised to receive either:

1. Intravaginal lactic acid gel used once daily for 7 days (Intervention)
2. Oral metronidazole tablets, 400mg twice daily for 7 days (control)

Participants are randomised using a remote internet based randomisation system that is developed and maintained by the Nottingham Clinical Trials Unit (NCTU).

Oral metronidazole is currently recommended as first line therapy in the UK national BV treatment guidelines and will be used within its licensed indication. Intravaginal lactic acid gel is a CE-marked medical device, which will be used within its intended purpose. Therefore, the study is classed as a clinical trial of an investigational medicinal product (CTIMP), where oral metronidazole is considered investigational medicinal product (IMP) for the purposes of the trial.

Following the initial visit, participants will then be followed up for resolution of their symptoms 2 weeks, 3 months and 6 months later via participant completed web-based questionnaires and self-sampling kits, completed at home.

Intervention Type

Drug

Pharmaceutical study type(s)

Not Applicable

Phase

Phase III

Drug/device/biological/vaccine name(s)

Metronidazole, lactic acid gel

Primary outcome measure

Resolution of bacterial vaginosis based on participant reported resolution of symptoms at week 2 (14 days from randomisation).

Secondary outcome measures

1. Time to first recurrence of BV is measured using patient reported questionnaires over 6 months
2. Number of participant reported BV episodes is measured using patient reported questionnaires over 6 months
3. Number of participant reported BV treatment courses is measured using patient reported questionnaires over 6 months
4. Microbiological resolution of BV is measured using microscopy of vaginal smears at week 2
5. Comparative tolerability of lactic acid gel and metronidazole assessed by participant reporting of side effects (including nausea, vomiting, taste disturbance, vaginal irritation, diarrhoea and abdominal pain) using patient reported questionnaire at 2 weeks and via participant interviews post intervention
6. Participant reported adherence to treatment is measured using participants reports at 2 weeks
7. Acceptability of treatments via qualitative assessment in a subgroup of participants post

intervention

8. Prevalence of concurrent sexually transmitted infections (gonorrhoea, chlamydia and trichomoniasis) at baseline and at week 2

9. Quality of life is assessed by SF-12 questionnaire at baseline, 2 weeks and 6 months

10. Comparative cost effectiveness measured via NHS Service use questionnaire at week 2, 3 and 6 months

Overall study start date

01/03/2017

Completion date

28/06/2020

Eligibility

Key inclusion criteria

1. Age 16 years or over

2. Clinical diagnosis of bacterial vaginosis – based on patient reported symptoms of discharge with an unpleasant (typically fishy) odour (with or without positive microscopy according to local site practice)

3. History of at least one previous episode of bacterial vaginosis within the past two years (clinically diagnosed or patient reported) which resolved with treatment

4. Willing to use either intravaginal lactic acid gel or oral tablets for the management of BV

5. Willing to take their own vaginal samples

6. Willing to avoid vaginal douching during treatment

7. Willing to provide contact details and be contacted for the purpose of collecting follow-up information

8. Willing to avoid sexual intercourse or use effective contraception for the 7-day duration of study treatment (condoms are not considered to be effective contraception due to a potential interaction with lactic acid gel)

9. Access to the internet, email and willing to complete web based follow up questionnaires in English

10. Written informed consent

Participant type(s)

Patient

Age group

Adult

Lower age limit

16 Years

Sex

Female

Target number of participants

Planned Sample Size: 1,900; UK Sample Size: 1,900

Total final enrolment

Key exclusion criteria

1. Contra-indications or allergy to lactic acid gel or metronidazole tablets
2. Pregnant or breastfeeding
3. Patients currently trying to conceive and not willing to avoid sexual intercourse or use effective contraception for the 7-day duration of study treatment
4. Use of oral antibiotics (other than the study treatment) or antifungal agents; concurrently, within the last 2 weeks or planned use within the next 2 weeks
5. Use of topical vaginal antibiotics, antifungals or acidifying products (other than the study treatment) concurrently, within the last 2 weeks, or planned use within the next 2 weeks
6. Previous participation in this study
7. Current participation in another clinical trial involving an investigational medicinal product

Date of first enrolment

25/09/2017

Date of final enrolment

28/06/2019

Locations**Countries of recruitment**

England

United Kingdom

Study participating centre**Nottingham Clinical Trials Unit (Coordinating Centre)**

The University of Nottingham
Nottingham Health Science Partners
South Block, Queen's Medical Centre
Nottingham
United Kingdom
NG7 2UH

Sponsor information**Organisation**

University Hospitals Birmingham NHS Foundation Trust

Sponsor details

Research and Development (Governance)
1st Floor, Institute of Translational Medicine (ITM)
University Hospitals Birmingham NHS Foundation Trust
Heritage Building, Queen Elizabeth Hospital, Mindelsohn Way

Birmingham
England
United Kingdom
B15 2TH
+44 (0)121 371 8006
Deborah.Popoola@uhb.nhs.uk

Sponsor type

Hospital/treatment centre

Website

<http://www.uhb.nhs.uk/>

ROR

<https://ror.org/014ja3n03>

Funder(s)

Funder type

Government

Funder Name

National Institute for Health Research

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Publication and dissemination plan

Planned publication in a high-impact peer reviewed journal.

Intention to publish date

31/03/2022

Individual participant data (IPD) sharing plan

The data sharing plans for the current study are unknown and will be made available at a later date.

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Protocol article	protocol	27/11/2019	02/12/2019	Yes	No
Results article		01/01/2022	24/01/2022	Yes	No
Results article		09/05/2023	10/05/2023	Yes	No
HRA research summary			28/06/2023	No	No
Results article	Qualitative findings	27/03/2023	20/07/2023	Yes	No
Other publications		15/11/2019	04/10/2024	Yes	No
Other publications	Side effects	03/07/2023	04/10/2024	Yes	No