

# Predrainage administration of Dopamine in the Renal function of selected patients with Obstructive Jaundice

<b>Submission date</b> 08/01/2010	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
<b>Registration date</b> 15/01/2010	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
<b>Last Edited</b> 15/01/2010	<b>Condition category</b> Digestive System	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Prof Javier Padillo

**Contact details**  
Imperio Argentina 43  
Cordoba  
Spain  
14014

## Additional identifiers

**EudraCT/CTIS number**

**IRAS number**

**ClinicalTrials.gov number**

**Secondary identifying numbers**  
N/A

## Study information

**Scientific Title**

Predrainage administration of dopamine associated with fluid administration improved renal function in selected patients with obstructive jaundice: a prospective randomised study

**Acronym**

DROJ

**Study objectives**

No studies have been carried out to evaluate the effects of dopamine in obstructive jaundice (OJ) patients on the renal and endocrine derangements observed in these patients before biliary drainage or surgery. The present study was therefore designed to analyse the effect of dopamine associated with fluid administration on extracellular water (ECW), water and sodium regulating hormones and renal function alterations in patients with OJ undergoing endoscopic internal biliary drainage.

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

Hospital Reina Sofia Clinical Trials and Ethics Committee approved in March 2005 (ref: PIO-20155)

**Study design**

Prospective randomised controlled study

**Primary study design**

Interventional

**Secondary study design**

Randomised controlled trial

**Study setting(s)**

Hospital

**Study type(s)**

Treatment

**Participant information sheet**

Not available in web format, please use the contact details below to request a patient information sheet

**Health condition(s) or problem(s) studied**

Obstructive jaundice

**Interventions**

Two treatment groups were created according to whether patients received dopamine at 3 mg /Kg/min or not, associated with 3000 ml of saline solution for 48 hours before biliary drainage in addition to their regular hospital ward diet. For the post-drainage study, patients treated with dopamine prior to biliary drainage, will be randomised according to whether they continue with dopamine for 72 hours or not. All patients will be kept under the same conditions. After

endoscopic internal drainage, patients will fast for the first 12 hours and 2500 ml of glucosaline solution containing 150 mEq of NaCl will be administrated until the following morning. On the second day, the intravenous infusion will be stopped and patients will receive a 2000 kcal/day diet for up to 72 hours when the study protocol will finish. Billiary drainage is considered successful if total bilirubin decreases by at least 30% and aerobilia with reduction of the common bile duct diameter demonstrated by ultrasound 72 hours after the procedure.

### **Intervention Type**

Drug

### **Phase**

Not Applicable

### **Drug/device/biological/vaccine name(s)**

Dopamine

### **Primary outcome measure**

Renal function (CrCl), assessed at admission, before drainage, as well as 24 hours and 72 hours after biliary drainage.

### **Secondary outcome measures**

Assessed at admission, before drainage, as well as 24 hours and 72 hours after biliary drainage:

1. Extracellular water (ECW) volume
2. Serum leves of aldosterone, renin, atrial natriuretic peptide (ANP), antidiuretic hormone (ADH), endothelin-1 (ET-1)
3. Urine prostaglandin E2 (PgE2)

### **Overall study start date**

10/01/2007

### **Completion date**

21/06/2009

## **Eligibility**

### **Key inclusion criteria**

1. Malignant OJ with serum bilirubin higher than 6 mg/dl
2. Ultrasound evidence of extrahepatic and intrahepatic bile duct dilation (more than 8 mm and 4 mm respectively)
3. Feasibility of internal endoscopic biliary drainage
4. Aged 35 to 76 years, either sex

### **Participant type(s)**

Patient

### **Age group**

Adult

### **Sex**

Both

**Target number of participants**

100

**Key exclusion criteria**

1. Cholangitis
2. Acute pancreatitis
3. Heart disease
4. Arterial hypertension
5. Chronic lung disease
6. Use of diuretics
7. Chronic renal failure

**Date of first enrolment**

10/01/2007

**Date of final enrolment**

21/06/2009

**Locations****Countries of recruitment**

Spain

**Study participating centre**

**Imperio Argentina 43**

Cordoba

Spain

14014

**Sponsor information****Organisation**

Reina Sofia University Hospital (Spain)

**Sponsor details**

Avda. Menéndez Pidal S/N

Cordoba

Spain

14004

**Sponsor type**

Hospital/treatment centre

**Website**

<http://www.juntadeandalucia.es/servicioandaluzdesalud/hrs2/>

**ROR**

<https://ror.org/02vtd2q19>

## **Funder(s)**

**Funder type**

Government

**Funder Name**

Spanish Ministry of Health (Spain) - Health Research Fund (Fondo de Investigaciones Sanitarias [FIS]) (ref: PIO-20155)

## **Results and Publications**

**Publication and dissemination plan**

Not provided at time of registration

**Intention to publish date**

**Individual participant data (IPD) sharing plan**

**IPD sharing plan summary**

Not provided at time of registration