

Carvedilol Treatment in Patients with Transfusion-Dependent Thalassemia

Submission date 18/07/2018	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 24/07/2018	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 08/11/2019	Condition category Circulatory System	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Thalassemia is a blood disorder that causes the body to make an abnormal form of hemoglobin, a protein that carries oxygen in the blood. Severe thalassemia can lead to hemolysis (bursting of red blood cells) and ineffective production of red blood cells. This can lead to symptoms such as anemia, jaundice, swelling of the liver and spleen, changes to the skeleton and delayed growth. For patients with such symptoms, the main treatment is regular blood transfusion and removal of excess iron from the body (iron chelation therapy).

Transfusion-dependent thalassemia is a form of thalassemia that relies on this treatment. Iron chelation therapy is also essential for patients with this, as the most common cause of death in patients with transfusion-dependent thalassemia is iron overload cardiomyopathy, where iron deposits in the heart and leads to heart dysfunction (ventricular dysfunction). Therefore, iron chelation therapy is required to remove this iron from the heart.

Carvedilol is a drug used to treat ventricular dysfunction after a heart attack. The aim of this study was to determine whether carvedilol is an effective treatment for transfusion-dependent thalassemia patients with heart dysfunction.

Who can participate?

Patients with transfusion-dependent thalassemia who have heart dysfunction

What does the study involve?

All patients will be given carvedilol to take daily for 6 months, and will have their heart function assessed before taking the drug and 3 and 6 months after.

What are the possible benefits and risks of participating?

The possible benefit to participants is that taking carvedilol may improve their heart function. The possible risks of participating are side effects of the drug, such as bradycardia and hypotension.

Where is the study run from?

Department of Pediatrics, Faculty of Medicine, Chiang Mai, Thailand

When does the study start and how long is it expected to run for?
October 2011 to December 2017

Who is funding the study?
Thai Research Fund (Thailand)

Who is the main contact?
Dr Suchaya Silvilairat
asilvilairat@gmail.com

Contact information

Type(s)
Scientific

Contact name
Dr Suchaya Silvilairat

ORCID ID
<http://orcid.org/0000-0001-7045-1817>

Contact details
Department of Pediatrics, Faculty of Medicine, Chiang Mai University
Chiang Mai
Thailand
50200

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
CTDT-2016

Study information

Scientific Title
Carvedilol improves left ventricular diastolic dysfunction in patients with Transfusion-Dependent Thalassemia

Acronym
CTDT

Study objectives
Carvedilol can attenuate LV diastolic dysfunction due to iron overload in patients with transfusion-dependent thalassemia who had ventricular dysfunction.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Research Ethics Committee 2, Faculty of Medicine, Chiang Mai University, 23/01/2012, PED-11-863-FB

Study design

Interventional prospective single-centre single-arm non-randomised cohort study

Primary study design

Interventional

Secondary study design

Non randomised study

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

No participant information sheet available

Health condition(s) or problem(s) studied

Transfusion-dependent thalassemia with ventricular diastolic dysfunction

Interventions

All patients were initially given carvedilol in doses of 0.1 mg/kg/day, the medication being divided into 2 separate doses. The dosage was doubled every two weeks until the target dose of 0.8 mg/kg/day was reached and continued for 6 months. The maximum dose was 50 mg/day. All patients had their cardiac function assessed by echocardiography at 3 and 6 months and the level of iron in the heart was also evaluated at 3 and 6 months using cardiac magnetic resonance imaging (MRI). Side effects of the drug were monitored though the study.

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Carvedilol

Primary outcome measure

Left ventricular diastolic function, measured by echocardiography at the baseline and 3 and 6 months after treatment. Echocardiographic data included pulse wave Doppler assessment of

mitral and pulmonary venous flows (ventricular diastolic filling analysis) and tissue Doppler imaging. Diastolic left ventricular dysfunction was graded according to pulse wave Doppler of mitral and pulmonary venous flows and tissue Doppler imaging.

Secondary outcome measures

Side effects of carvedilol use, assessed at the baseline and 3 and 6 months after treatment:

1. Blood pressure
2. Heart rate
3. Symptoms such as dizziness, chest pain and swelling

Overall study start date

01/10/2011

Completion date

01/12/2017

Eligibility

Key inclusion criteria

1. Severe forms of homozygous beta-thalassemia and beta-thalassemia/hemoglobin E disease
2. Transfusion-dependent thalassemia
3. Aged 8-25 years
4. Left ventricular diastolic dysfunction

Participant type(s)

Patient

Age group

Other

Sex

Both

Target number of participants

15

Key exclusion criteria

N/A

Date of first enrolment

01/01/2014

Date of final enrolment

31/12/2016

Locations

Countries of recruitment

Thailand

Study participating centre
Chiang Mai University
110 Tambon. Sripoom
Inthawaroraj Road
A.Muang
Chiang Mai
Thailand
50200

Sponsor information

Organisation
Thailand Research Fund

Sponsor details
14th Floor, SM Tower,
979/17-21 Phaholyothin Road, Samsan Nai, Phyathai
Bangkok
Thailand
10400
6622788200
callcenter@trf.or.th

Sponsor type
Government

Website
<https://www.trf.or.th>

Funder(s)

Funder type
Not defined

Funder Name
Thailand Research Fund

Results and Publications

Publication and dissemination plan
Planned publication in a high-impact peer reviewed journal.

Intention to publish date

31/08/2018

Individual participant data (IPD) sharing plan

Only the final conclusion of analysis of the results of the study will be available upon request from asilvilairat@gmail.com.

IPD sharing plan summary

Available on request