

Prolonging muscle paralysis after surgery: a comparative age-matched study

Submission date 16/04/2017	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 02/06/2017	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 10/08/2017	Condition category Nervous System Diseases	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and purpose of study

Neuromuscular blockade (NM) is a technique which involves interrupting nerve signals with medication in order to paralyse muscles. This technique is commonly used during surgery, as they can assist the effects of general anaesthesia (putting someone to sleep for an operation). Although this technique works very well, it can be associated with undesirable side effects, such as prolonged muscle paralysis after surgery (residual muscle paralysis). In some cases, this can lead to the muscles of the body not getting enough oxygen and difficulties with the airways. Currently, there is little research looking at residual muscle paralysis and age. The aim of this study is to find out how common residual muscle paralysis is in patients aged between 2 and 10 years, 18 and 64 years, and over 65 years.

Who can participate?

Patients aged between 2-10, 18-64 and 65+ years who have been scheduled for surgery under general anesthesia with NM.

What does the study involve?

Patients are assigned to one of three groups based on their age. During their surgery, patients undergo the same treatment with general anaesthesia and muscle relaxants. Every 15 minutes after surgery for the five hours they are in the recovery room, participants are monitored to find out what (if any) degree of muscle paralysis is present. After being discharged from the recovery room, patients are followed up every three hours until 24 hours after surgery.

What are the possible benefits and risks of participating?

All participants will benefit from regular monitoring of muscle paralysis as part of the study. There are no notable risks involved with participating.

Where is the study run from?

First Affiliated Hospital of Dalian Medical University (China)

When is the study starting and how long is it expected to run for?

December 2016 to December 2017

Who is funding the study?
First Affiliated Hospital of Dalian Medical University (China)

Who is the main contact?
Dr Wen Qingping
yabasin@dmu.edu.cn

Contact information

Type(s)
Scientific

Contact name
Dr Wen Qingping

Contact details
First Affiliated Hospital of Dalian Medical University
Department of Anesthesiology
222 Zhongshan Road
Dalian
China
16011
+86 13252991927
yabasin@dmu.edu.cn

Additional identifiers

Protocol serial number
KY2017

Study information

Scientific Title
Residual neuromuscular blockade following surgery and anesthesia: A prospective age-matched cohort study

Study objectives
40% of the patients in the age bracket 2-10 years and 65+ years cohort will develop postoperative residual neuromuscular blockade (TOF ratio 0.9).

Ethics approval required
Old ethics approval format

Ethics approval(s)
Human Research Ethics Committee of the First Affiliated Hospital of Dalian Medical University, 10/01/2017, ref: KY2017

Study design
Prospective observational cohort study

Primary study design

Observational

Study type(s)

Other

Health condition(s) or problem(s) studied

Post-operative residual neuromuscular blockade

Interventions

Participants are allocated to one of three study cohorts based on their age (2-10 years, 18-64 years and 65+ years), which each undergo the following:

Neuromuscular monitoring

The same standard of general anesthesia protocol will be employed for all the study participants during the surgery. Management of muscle blockade will be standardized in all the three study groups. On admission at the post-anesthesia care unit (PACU), surface ECG electrodes for neuromuscular monitoring will be applied on the cleansed skin over the ulnar nerve close to the wrist. To determine quantitative train-of-four (TOF) ratio measurement in all study participants, the leads of train-of-four (TOF) Watch SX will be connected to the surface ECG electrode on the patient arm. The transducer of the TOF Watch-SX that measures the acceleration of the patients' muscles will be applied to the thumb of patients' hand on which the ECG surface electrode will be placed. Based on standard guidelines, the hand adapter will be applied to the anterior surface of the four fingers of the hand. Two serial TOF measurements at 15 minutes intervals will be obtained in all participants, averaged and documented. TOF ratios 0.7 and 0.8 will be defined as moderate neuromuscular block, while TOF ratio of less than 0.7 will be defined as severe residual neuromuscular block. Participants with various grades of TOF ratios (TOF ratio less than 0.7, 0.8 and 0.9) in each study cohort will be summed up and compared. In addition to TOF ratio data to be collected, study participants will be assessed at 15 minutes interval for clinical features of neuromuscular blockade on PACU admission. In this regard a structured examination sheet will be employed to capture data about whether or there is muscle weakness in various muscle groups.

Potential pulmonary complications associated with residual neuromuscular block

Arterial oxygen saturation (SpO₂) will be continuously monitored through pulse oximetry in all study participants during PACU admission. While in PACU, study participants of each cohort will be put on 5 litters face mask airway and oxygenation monitored continuously and data saved. This will enable investigators determine early signs of airway obstruction.

Duration of PACU Hospitalization

The influence of residual neuromuscular block on duration of PACU admission will assessed. Therefore, the time to discharge criteria and the actual discharge time will be recorded.

Duration of participant observation

The maximum duration of observation of patients in PACU will be 5 hours. After discharge from PACU, 3 hourly follow ups in the ward will be observed until the 24 hours postoperative phase elapsed. This will enable investigators detect residual neuromuscular block and its related pulmonary or respiratory complications that may occur on the ward.

Intervention Type

Procedure/Surgery

Primary outcome(s)

Residual neuromuscular blockade (RNMB), defined as train-of-four (TOF) ratio of less than 0.9. Moderate (TOF ratio ≥ 0.7 but < 0.9) and severe (TOF ratio < 0.7) PRNMB, is assessed 5 minutes to administration of cisatracurium at induction of general anesthesia, 1 minute interval after cisatracurium injection until 6 minutes and every 15 minutes in the postanesthetic care unit (PACU) until discharge.

Key secondary outcome(s)

1. Demographic and patient characteristics such as age, weight, sex and co-morbidities, airway obstruction, indication of surgery, anesthesia duration, time (minutes) to PACU discharge and respiratory complications during hospital admission and length of hospital admission (days)
2. Total crystalloids and colloids volume (mls), total cisatracurium (mg) and total neostigmine (mg)
3. Perioperative and PACU temperature ($^{\circ}$) monitoring during perioperative and PACU periods
4. Peripheral oxygen saturation (SpO₂) monitoring during perioperative and PACU periods
5. End tidal carbon dioxide (EtCO₂) monitoring during perioperative and PACU periods
6. Heart rate, systolic, diastolic and mean arterial pressure (mmHg) monitoring during perioperative and PACU periods.
7. Bispectral index (BIS) monitoring perioperative and PACU periods

Completion date

30/12/2017

Eligibility

Key inclusion criteria

1. Patients with American Society of Anesthesiologists (ASA) physical status I-III scheduled to undergo elective surgery undergo general with muscle relaxants
2. Patients within the age groups 2-10, 18-64 and 65+ years

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Mixed

Lower age limit

2 years

Sex

All

Key exclusion criteria

1. Patients with muscular disorders
2. Patients with neurological deficit
3. Patient refusal to consent
4. Patients scheduled to undergo prolong surgical procedures

5. Patients with severe impaired liver and renal function
6. Infants under 2 years
7. Patients with asthma
8. Patients with diabetes mellitus

Date of first enrolment

01/05/2017

Date of final enrolment

30/10/2017

Locations

Countries of recruitment

China

Study participating centre

First Affiliated Hospital of Dalian Medical University

Department of Anesthesiology

222 Zhongshan Road

Liaoning, Dalian

China

116011

Sponsor information

Organisation

First Affiliated Hospital of Dalian Medical University

ROR

<https://ror.org/055w74b96>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

First Affiliated Hospital of Dalian Medical University

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Wen Qingping (yabasin@dmu.edu.cn)

IPD sharing plan summary

Available on request