# Motivating structured walking activity in intermittent claudication

Submission date
31/07/2017

Registration date
02/08/2017

Recruitment status
No longer recruiting
[X] Protocol
[X] Protocol
[X] Statistical analysis plan
[X] Results

Last Edited Condition category Individual participant data
19/09/2023 Circulatory System

#### Plain English summary of protocol

Background and study aims

Peripheral arterial disease can cause leg pain or discomfort (called intermittent claudication (IC)), which limits the ability to walk and carry out everyday activities. Lifestyle changes, like increasing walking, can lead to improvements, but can be a challenge to begin and then continue. The aim of this study is to investigate if a physiotherapist-led, behaviour change treatment effects walking in people with IC. The treatment is designed to build an understanding of IC and walking exercise and help individuals develop strategies to increase regular walking.

Who can participate?
Adults aged 50 years or older who have IC

#### What does the study involve?

Participants attend a study visit to complete two short walking tests and answer questionnaires about their daily activities, beliefs about their symptoms and treatment, quality of life and the costs of having IC. Participants are then randomly allocated to one of two groups. Those in the first group receive the new treatment which involves two 60-minute face-to-face sessions (at the participant's home or local hospital) and two 20-minute telephone calls with a physiotherapist to discuss participants' understanding and beliefs about IC, walking and help participants to set goals and plans to increase walking over 12 weeks. Participants are provided with a step counter (pedometer) and walking record. Those in the second group receive their normal care. After 3 months, all participants attend a second appointment where they will repeat the walking tests and fill out another set of questionnaires. A final set of questionnaires are completed by all participants after 6 months (by post or electronically). Some participants will be invited to provide feedback on their experience of the treatment and trial by telephone or a face-to-face interview with a researcher.

What are the possible benefits and risks of participating?

Participants may benefit from receiving consultations with a physiotherapist, and receiving a step counter and walking diary which can help them increase their walking ability. Participants may benefit from completing a walk test and learning about their own walking ability and walking as a treatment for intermittent claudication. Participants who engage in the walking

activity may experience an increase in pain or discomfort. The treatment involves education and reassurance that these symptoms are unpleasant but do not indicate harm.

Where is the study run from? Guy's Hospital (UK)

When is the study starting and how long is it expected to run for? July 2017 to July 2020

Who is funding the study?
The Dunhill Medical Trust (UK)

Who is the main contact?

Dr Lindsay Bearne, Lindsay.bearne@kcl.ac.uk
(updated 18/11/2020, previously: Dr Julie Bieles, julie.bieles@kcl.ac.uk)

## Contact information

#### Type(s)

Scientific

#### Contact name

Dr Lindsay Bearne

#### Contact details

King's College London
Faculty of Life Sciences & Medicine
School of Population Health & Environmental Sciences
Addison House
Guy's Campus
London
United Kingdom
SE1 1UL
+44 (0)2078486283
Lindsay.bearne@kcl.ac.uk

## Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Integrated Research Application System (IRAS)

215024

ClinicalTrials.gov (NCT)

NCT03238222

Protocol serial number

CPMS 34022, IRAS 215024

## Study information

#### Scientific Title

A brief physiotherapist-led behaviour-change intervention to facilitate walking in older people with peripheral arterial disease: a randomised controlled trial

#### **Acronym**

**MOSAIC** 

#### **Study objectives**

people with IC?

The aim of this study is to answer the questions:

- 1. Does MOSAIC improve walking ability (measured by the 6 Minute Walking Distance [6MWD]) at 3 months compared to usual NHS care in older people with intermittent claudication (IC)?
- 2. Does MOSAIC improve activities of daily living and QoL at 3 months and walking ability, activities of daily living and QoL at 6 months compared to usual NHS care in people with IC?

  3. Is it feasible to collect the measures required to estimate cost utility in future phase 3 trials in
- 4. What are the Mean Clinically Important Difference (MCID) thresholds for the assessment measures in this population?

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

London-Bloomsbury Research Ethics Committee, 27/04/2017, 17/LO/0568

#### Study design

Randomised; Interventional; Design type: Treatment, Education or Self-Management, Psychological & Behavioural, Complex Intervention, Physical, Rehabilitation

#### Primary study design

Interventional

#### Study type(s)

**Treatment** 

#### Health condition(s) or problem(s) studied

Intermittent claudication

#### **Interventions**

Participants visit the study site to complete two short walking tests and answer questionnaires about their daily activities, beliefs about their symptoms and treatment, quality of life and the costs of having IC.

Following informed consent and completion of the baseline assessment, the participant is randomised to one of two arms. Participants are randomised at the level of the individual, stratified within recruitment site, to receive either MOSAIC plus usual NHS care or usual NHS care alone in a ratio of 1:1. The King's College London Clinical Trials Unit provides randomisation. The system is online and can be accessed 24 hours a day. The RA logs in, enters key information about the participant, and the randomisation allocation occurs instantly. Confirmation emails are

generated automatically and sent to the CI (and delegated members of the research team) and trial physiotherapists delivering MOSAIC. The RA, who is the outcome assessor, is blind to group allocation. Once the participant is randomised they are given a unique trial identifying number.

Arm 1 MOSAIC Treatment: Participants in this group receive two 60-minute individual face-to-face consultations (on weeks one & two) and two 20-minute follow-up telephone calls (weeks six & 12) delivered at a convenient time and location of participant's choice (local NHS Trust or participant's home). All sessions are delivered by a trained Band 6/7 physiotherapist. A checklist outlining the components for each session is provided to each physiotherapist. All participants randomized to this arm are provided with a pedometer and a patient manual which include information on intermittent claudication (IC), risk factors, walking guidelines, goal setting, problem solving and action planning worksheets and a walking diary.

Arm 2 Usual Care comparison: Participants randomized to the comparison group continue to receive usual NHS care for intermittent claudication which typically consists of an initial assessment, drug therapy and simple advice to walk provided by a vascular specialist and delivered in the vascular outpatient clinic.

After 3 months, all participants attend a second appointment where they will repeat the walking tests and fill out another set of questionnaires. A final set of questionnaires are completed by all participants after 6 months (by post or electronically). Some participants are invited to provide feedback on their experience of the treatment and trial by telephone or a face-to-face interview with a researcher.

#### **Intervention Type**

Behavioural

#### Primary outcome(s)

The 6 Minute Walking Distance (in metres) at 3 months is measured during a self-paced, standardised 6 Minute Walk Test conducted around a level, 100-foot circuit.

#### Key secondary outcome(s))

- 1. Pain-free and maximal walking ability measured during a 6 Minute Walk Test at baseline, 3 and 6 months
- 2. Self-reported maximal walking distance measured by a global single-item at baseline, 3 and 6 months
- 3. Self-reported walking ability measured by the Walking Estimated-Limitation Calculated by History [WELCH] questionnaire at baseline, 3 and 6 months
- 4. Activities of Daily Living measured by the Nottingham Extended Activities of Daily Living [NEADL] questionnaire at baseline, 3 and 6 months
- 5. Quality of Life measured by the Vascular Quality of Life Questionnaire-6 [VascuQol-6] at baseline, 3 and 6 months

#### Process and feasibility outcomes:

- 1. Treatment cognitions measured by a Theory of Planned Behaviour Questionnaire (TPBQ) at baseline, 3 and 6 months
- 2. Illness cognitions measured by the Brief Illness Perceptions Questionnaire (BIPQ) at baseline, 3 and 6 months
- 3. Walking adherence measured by the Exercise Adherence Rating Scale (EARS) at baseline, 3

and 6 months

- 4. Resource use measured by the Client Services Receipt Inventory at baseline 3 and 6 months
- 5. Quality of Life measured by the EuroQuol 5D-5L (EQ-5D-5L) at baseline 3 and 6 months

#### Completion date

09/01/2021

## **Eligibility**

#### Key inclusion criteria

- 1. ≥50 years of age
- 2. Established PAD (Ankle Brachial Pressure Index ≤0.90 and/or radiographic evidence [added 21 /02/2019: or clinician reported diagnosis]) and IC (presence of symptoms reported on the San Diego Claudication Questionnaire (SDCQ))
- 3. Able to participate in MOSAIC
- 4. Able and willing to provide informed consent

#### Participant type(s)

Patient

#### Healthy volunteers allowed

No

#### Age group

Adult

#### Sex

All

#### Total final enrolment

190

#### Key exclusion criteria

- 1. Unstable IC (self-reported change in symptoms in previous 3 months)
- 2. Walking more than 90 minutes per week (reported via Brief International Physical Activity Questionnaire
- 3. Contraindications to walking exercise (e.g. unstable angina) confirmed by their vascular specialist
- 4. Have completed any prescribed exercise sessions in the previous 6 months or been offered prescribed exercise sessions in the next 6 months.

#### Date of first enrolment

10/10/2017

#### Date of final enrolment

20/03/2020

## Locations

#### Countries of recruitment

#### United Kingdom

#### England

## Study participating centre Guy's Hospital

Guy's & St Thomas' NHS Foundation Trust Great Maze Pond London United Kingdom SE1 9RT

## Sponsor information

### Organisation

King's College London

#### Organisation

Guy's & St Thomas' NHS Foundation Trust

## Funder(s)

#### Funder type

Hospital/treatment centre

#### **Funder Name**

**Dunhill Medical Trust** 

#### Alternative Name(s)

The Dunhill Medical Trust, Dunhill Medical Trust, DunhillMedical, DMT

#### **Funding Body Type**

Private sector organisation

#### **Funding Body Subtype**

Trusts, charities, foundations (both public and private)

#### Location

**United Kingdom** 

## **Results and Publications**

#### Individual participant data (IPD) sharing plan

Current IPD sharing statement as of 14/04/2022:

Requests for sharing deidentified data should be addressed to Lindsay Bearne at lindsay. bearne@kcl.ac.uk.

#### Previous IPD sharing statement:

The data sharing plans for the current study are unknown and will be made available at a later date.

#### IPD sharing plan summary

Available on request

#### **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		12/04/2022	13/04/2022	Yes	No
Protocol article	protocol	24/08/2019	22/11/2019	Yes	No
HRA research summary			28/06/2023	No	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Protocol file	version 6	28/08/2018	19/09/2023	No	No
Statistical Analysis Plan	version 1	08/01/2019	19/09/2023	No	No