# Cognitive behaviour therapy for excessive worry over health in medical patients

Submission date	Recruitment status No longer recruiting	[X] Prospectively registered		
17/06/2008		[X] Protocol		
Registration date	Overall study status	Statistical analysis plan		
20/06/2008	Completed	[X] Results		
Last Edited	Condition category	Individual participant data		
17/05/2016	Mental and Behavioural Disorders			

### Plain English summary of protocol

Background and study aims

Hypochondriasis, or health anxiety, is a significant problem in all parts of clinical practice, but has been largely ignored by services until recently. Our studies, and those of others, suggest that between 1 in 4 and 1 in 10 of all those attending medical clinics have significant health anxiety and this accounts for both considerable suffering and unnecessary use of resources. One of the characteristics of those with severe health anxiety is that it persists and leads to frequent medical consultations in both primary and secondary care. This represents a significant drain on health service resources at a time of considerable cost pressures, and, in addition, the symptoms of health anxiety are highly troubling and disturbing, are associated with much time off work, and adversely affect social functioning. In a study of treatment in a genitourinary medicine clinic, we found that a short course of cognitive behaviour therapy (CBT) was highly effective at reducing health anxiety, and that these benefits are maintained over a period of one year. We also found that the number of consultations in both general practice and the clinics fell in those in the active treatment group over one year but the savings made were small as the costs of investigations are relatively low. We are now planning a larger study of CBT to treat health anxiety in cardiology, respiratory medicine, gastroenterology and endocrinology clinics, where such anxiety is more common than in genuitourinary medicine.

#### Who can participate?

Patients aged between 16 and 75 attending four medical specialty clinics (in cardiology, respiratory medicine, gastroenterology and endocrinology) who have significant health anxiety

#### What does the study involve?

Participants are randomly allocated to one of two groups. One group is treated with CBT adapted for health anxiety in the form of between 5 and 10 one-hour sessions, which will address abnormal worries about health and ways of overcoming them. The other group is treated with a single 45-minute explanatory interview describing the nature of health anxiety and how it tends to be perpetuated. We measure patients' functioning, anxiety and quality of life, as well as health service and employment costs, at the start of the study and at 6 monthly intervals for two years, to see whether the costs of treatment are offset by savings on attendance and investigations over 2 years.

What are the possible benefits and risks of participating? Not provided at time of registration

Where is the study run from? Imperial College London (UK)

When is the study starting and how long is it expected to run for? October 2008 to September 2012

Who is funding the study? Health Technology Assessment Programme (UK)

Who is the main contact? Prof. Peter Tyrer p.tyrer@imperial.ac.uk

# Contact information

# Type(s)

Scientific

#### Contact name

**Prof Peter Tyrer** 

#### Contact details

Department of Psychological Medicine St Dunstan's Road London United Kingdom W6 8RP +44 (0)207 386 1237 p.tyrer@imperial.ac.uk

# Additional identifiers

Protocol serial number HTA 07/01/26

# Study information

#### Scientific Title

Cognitive behaviour therapy for Health Anxiety in Medical Patients

#### **Acronym**

**CHAMP** 

# Study objectives

Cognitive behaviour therapy adapted for health anxiety is a cost-effective way of treating patients with significant health anxiety in medical clinics.

More details can be found at: http://www.nets.nihr.ac.uk/projects/hta/070126 Protocol can be found at: http://www.nets.nihr.ac.uk/\_\_data/assets/pdf\_file/0017/51731/PRO-07-01-26.pdf

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

Nottingham Research Ethics Committee 1, 13/06/2008, ref: 08/H0403/56

#### Study design

Randomised controlled trial

## Primary study design

Interventional

#### Study type(s)

Treatment

#### Health condition(s) or problem(s) studied

Hypochondriasis/mental health

#### Interventions

Arm 1: Cognitive behaviour therapy adapted for health anxiety (Warwick/Salkovskis model). Treatment will be given for between 5 and 10 one hour sessions for each patient, and will address abnormal worries about health and ways of overcoming them.

Arm 2: Single explanatory interview of the nature of health anxiety. This will be a single interview of 45 minutes describing the nature of health anxiety and how it tends to be perpetuated.

Duration of interventions: up to 6 months

Duration of follow-up: 2 years

## Intervention Type

Other

#### Phase

**Not Specified** 

#### Primary outcome(s)

Reduction in health anxiety scores 1 year after randomisation.

## Key secondary outcome(s))

- 1. Reduction in health anxiety scores 2 years after randomisation
- 2. Presence or absence of diagnosis of hypochondriasis at 1 and 2 years after randomisation
- 3. Reduction in generalised anxiety and depression, assessed using the Hospital Anxiety and Depression Scale (HADS) at 1 and 2 years after randomisation
- 4. Improvement in social function and quality of life, assessed using the Social Functioning

Questionnaire (SFQ) and Eurogol EQ-5D at 1 and 2 years after randomisation

- 5. Total costs of health care (primary and secondary) 1 and 2 years after randomisation
- 6. Change in presenteeism and absenteeism at work 1 and 2 years after randomisation

#### Completion date

30/09/2012

# Eligibility

#### Key inclusion criteria

Patients attending four medical specialty clinics (in cardiology, respiratory medicine, gastroenterology and endocrinology) who:

- 1. Have significant health anxiety (score of 20 or more on the Health Anxiety Inventory)
- 2. Are aged between 16 and 75 (both males and females)
- 3. Are living independently
- 4. Are permanent residents in the area
- 5. Have sufficient understanding of English to read and complete the questionnaires
- 6. Give written consent for the interviews, including audiotaping of half the sessions
- 7. Give written consent for access to their medical records

#### Participant type(s)

Patient

# Healthy volunteers allowed

No

## Age group

Adult

#### Sex

All

## Key exclusion criteria

Patients who are:

- 1. Considered too ill in medical terms by their consultants to be considered for the study
- 2. In the process of being investigated for significant pathology and for whom cognitive behaviour therapy might confuse or cause distress
- 3. Those who have significant cognitive impairment

#### Date of first enrolment

01/10/2008

#### Date of final enrolment

30/09/2012

# Locations

#### Countries of recruitment

United Kingdom

Study participating centre Imperial College London London United Kingdom W6 8RP

# Sponsor information

#### Organisation

Imperial College London (UK)

#### **ROR**

https://ror.org/041kmwe10

# Funder(s)

# Funder type

Government

#### **Funder Name**

Health Technology Assessment Programme

#### Alternative Name(s)

NIHR Health Technology Assessment Programme, Health Technology Assessment (HTA), HTA

#### **Funding Body Type**

Government organisation

#### **Funding Body Subtype**

National government

#### Location

**United Kingdom** 

# **Results and Publications**

Individual participant data (IPD) sharing plan

# IPD sharing plan summary

# Study outputs

Output type	Details	Date created Date added	Peer reviewed?	Patient-facing?
Results article	results	18/01/2014	Yes	No
<u>Protocol article</u>	protocol	14/06/2011	Yes	No
Participant information sheet	Participant information sheet	11/11/2025 11/11/2025	No	Yes